

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
9 December 2004 (09.12.2004)

PCT

(10) International Publication Number
WO 2004/105765 A1

(51) International Patent Classification⁷: **A61K 31/529**,
C07D 498/08, A61P 35/00, C07D 239/88, 239/93, 239/94

(21) International Application Number:
PCT/EP2004/005621

(22) International Filing Date: 25 May 2004 (25.05.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PCT/EP03/05723 27 May 2003 (27.05.2003) EP
PCT/EP03/10266 15 September 2003 (15.09.2003) EP
PCT/EP03/51061 18 December 2003 (18.12.2003) EP

(71) Applicant (for all designated States except US):
JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **FREYNE**, Eddy, Jean, Edgard [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). **PERERA**, Timothy, Pietro, Suren [GB/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). **BUIJNSTERS**, Peter, Jacobus, Johannes, Antonius [NL/NL]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). **WILLEMS**, Marc [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). **DIELS**, Gaston, Stanislas, Marcella [BE/BE];

Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). **EMBRECHTS**, Werner, Constant, Johan [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). **TEN HOLTE**, Peter [NL/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE).

(74) Common Representative: **JANSSEN PHARMACEUTICA N.V.**; Turnhoutseweg 30, B-2340 Beerse (BE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,

[Continued on next page]

(54) Title: MACROCYCLIC QUINAZOLINE DERIVATIVES AS ANTIPROLIFERATIVE AGENTS



(I)

(57) Abstract: The present invention concerns the compounds of formula (I) the N-oxide forms, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein Z represents O, CH₂, NH or S; in particular Z represents NH; Y represents -C₃₋₉alkyl-, -C₃₋₉alkenyl-, -C₃₋₉alkynyl-, -C₃₋₇alkyl-CO-NH- optionally substituted with amino, mono- or di(C₁₋₄alkyl)amino or C₁₋₄alkyloxycarbonylamino-, -C₃₋₇alkenyl-CO-NH- optionally substituted with amino, mono- or di(C₁₋₄alkyl)amino- or C₁₋₄alkyloxycarbonylamino-, C₁₋₃alkyl-oxy-C₁₋₃alkyl-, -C₁₋₃alkyl NR¹³-, -C₁₋₃alkyl-, -C₁₋₃alkyl-NR¹⁴-CO-C₁₋₃alkyl-, -C₁₋₃alkyl-CO NR¹⁵-C₁₋₃alkyl-, -C₁₋₆alkyl-CO-NH-, -C₁₋₆alkyl-NH-CO-, -C₁₋₃alkyl-NH-CS-Het²⁰-, -C₁₋₃alkyl-NH-CO-Het²⁰-, -C₁₋₂alkyl-CO-Het²¹-CO-, -Het²²-CH₂-CO-NH-C₁₋₃alkyl-, -CO-NH-C₁₋₆alkyl-, -NH-CO-C₁₋₆alkyl-, -CO-C₁₋₇alkyl-, -C₁₋₇alkyl-CO-, -C₁₋₂alkyl-CO-NH-CR¹⁶R¹⁹-CO-, -C₁₋₂alkyl-CO-NR²⁰-C₁₋₃alkyl-CO-, C₁₋₂alkyl-NR²¹-CH₂-CO-NH-C₁₋₃alkyl-, or -NR²²-CO-C₁₋₃alkyl-NH-; X¹ represents a direct bond, O or -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹¹-, -NR¹¹-C₁₋₂alkyl-, CH₂-, -O-N=CH- or -C₁₋₂alkyl-; X² represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹²-, -NR¹²-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or -C₁₋₂alkyl-. The growth inhibitory effect anti-tumour activity of the present compounds has been demonstrated *in vitro*, in enzymatic assays on the receptor tyrosine kinase EGFR.

WO 2004/105765 A1

BEST AVAILABLE COPY



CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- of inventorship (Rule 4.17(iv)) for US only

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

MACROCYCLIC QUINAZOLINE DERIVATIVES AS ANTIPROLIFERATIVE AGENTS

5 This invention relates to quinazoline derived macrocycles that have been found to possess anti-proliferative activity, such as anti-cancer activity and are accordingly useful in methods of treatment of the human or animal body, for example in the manufacture of medicaments for use in hyper proliferative disorders such as atherosclerosis, restenosis and cancer. The invention also relates to processes for the manufacture of said quinazoline derivatives, to pharmaceutical compositions containing
10 them and to their use in the manufacture of medicaments of use in the production of anti-proliferative effect.

In particular, the compounds of the present invention were found to inhibit tyrosine kinase enzymes, also called tyrosine kinases. Tyrosine kinases are a class of enzymes, which catalyse the transfer of the terminal phosphate of adenosine
15 triphosphate to the phenolic hydroxyl group of a tyrosine residue present in the target protein. It is known, that several oncogenes, involved in the transformation of a cell into a malignant tumour cell, encode tyrosine kinase enzymes including certain growth factor receptors such as EGF, FGF, IGF-1R, IR, PDGF and VEGF. This family of receptor tyrosine kinases and in particular the EGF family of receptor tyrosine kinases
20 are frequently present in common human cancers such as breast cancer, non-small cell lung cancers including adenocarcinomas and squamous cell cancer of the lung, bladder cancer, oesophageal cancer, gastrointestinal cancer such as colon, rectal or stomach cancer, cancer of the prostate, leukaemia and ovarian, bronchial or pancreatic cancer, which are examples of cell proliferation disorders.

25 Accordingly, it has been recognised that the selective inhibition of tyrosine kinases will be of value in the treatment of cell proliferation related disorders. Support for this view is provided by the development of Herceptin® (Trastuzumab) and Gleevec™ (imatinib mesylate) the first examples of target based cancer drugs.
30 Herceptin® (Trastuzumab) is targeted against Her2/*neu*, a receptor tyrosine kinase found to be amplified up to 100-fold in about 30% of patients with invasive breast cancer. In clinical trials Herceptin® (Trastuzumab) proved to have anti-tumour activity against breast cancer (Review by L.K. Shawer *et al*, "Smart Drugs: Tyrosine kinase inhibitors in cancer therapy", 2002, Cancer Cell Vol.1, 117), and accordingly provided
35 the proof of principle for therapy targeted to receptor tyrosine kinases. The second example, Gleevec™ (imatinib mesylate), is targeted against the abelson tyrosine kinase (Bcr-Abl), a constitutively active cytoplasmic tyrosine kinase present in virtually all patients with chronic myelogenous leukaemia (CML) and 15% to 30% of adult patients

-2-

with acute lymphoblastic leukaemia. In clinical trials Gleevec™ (imatinib mesylate) showed a spectacular efficacy with minimal side effects that led to an approval within 3 months of submission. The speed of passage of this agent through clinical trials and regulatory review has become a case study in rapid drug development (Drucker B.J. & Lydon N., "Lessons learned from the development of an Abl tyrosine kinase inhibitor for chronic myelogenous leukaemia.", 2000, J.Clin.Invest. 105, 3).

Further support is given by the demonstration that EGF receptor tyrosine kinase inhibitors, specifically attenuates the growth in athymic nude mice of transplanted carcinomas such as human mammary carcinoma or human squamous cell carcinoma (Review by T.R. Burke Jr., *Drugs of the Future*, 1992, 17, 119). As a consequence, there has been considerable interest in the development of drugs to treat different cancers that target the EGFR receptor. For example, several antibodies that bind to the extra-cellular domain of EGFR are undergoing clinical trials, including Erbitux™ (also called C225, Cetuximab), which was developed by Imclone Systems and is in Phase III clinical trials for the treatment of several cancers. Also, several promising orally active drugs that are potent and relatively specific inhibitors of the EGFR tyrosine kinase are now well advanced in clinical trials. The AstraZeneca compound ZD1839, which is now called IRESSA® and approved for the treatment of advanced non-small-cell lung cancer, and the OSI/Genentech/Roche compound OSI-774, which is now called Tarceva™ (erlotinib), have shown marked efficacy against several cancers in human clinical trials (Morin M.J., "From oncogene to drug: development of small molecule tyrosine kinase inhibitors as anti-tumour and anti-angiogenic agents, 2000, *Oncogene* 19, 6574).

25

In addition to the above, EGF receptor tyrosine kinases has been shown to be implicated in non-malignant proliferative disorders such as psoriasis (Elder *et al.*, *Science*, 1989, 243; 811). It is therefore expected that inhibitors of EGF type receptor tyrosine kinases will be useful in the treatment of non-malignant diseases of excessive cellular proliferation such as psoriasis, benign prostatic hypertrophy, atherosclerosis and restenosis.

30

It is disclosed in International Patent Application WO96/33980 and in *J. Med. Chem.*, 2002, 45, 3865 that certain 4 anilino substituted quinazoline derivatives may be useful as inhibitors of tyrosine kinase and in particular of the EGF type receptor tyrosine kinases. Unexpectedly it was found that Quinazoline derivatives of the present formula (I) that are different in structure, show to have tyrosine kinase inhibitory activity.


35

-3-

It is accordingly an object of the present invention to provide further tyrosine kinase inhibitors useful in the manufacture of medicaments in the treatment of cell proliferative related disorders.

5

This invention concerns compounds of formula (I)



10

the *N*-oxide forms, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

Z represents O, CH₂, NH or S; in particular Z represents NH;

15 Y represents -C₃₋₉alkyl-, -C₃₋₉alkenyl-, -C₃₋₉alkynyl-,

-C₃₋₇alkyl-CO-NH- optionally substituted with amino, mono- or di(C₁₋₄alkyl)amino or C₁₋₄alkyloxycarbonylamino-, -C₃₋₇alkenyl-CO-NH- optionally substituted with amino, mono- or di(C₁₋₄alkyl)amino or C₁₋₄alkyloxycarbonylamino-,

20 -C₃₋₇alkynyl-CO-NH- optionally substituted with amino, mono- or di(C₁₋₄alkyl)amino or C₁₋₄alkyloxycarbonylamino-,

-C₁₋₅alkyl-oxy-C₁₋₅alkyl-, -C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-,

-C₁₋₅alkyl-NR¹⁴-CO-C₁₋₅alkyl-,

-C₁₋₅alkyl-CO-NR¹⁵-C₁₋₅alkyl-, -C₁₋₆alkyl-CO-NH-, -C₁₋₆alkyl-NH-CO-,

-C₁₋₃alkyl-NH-CS-Het²⁰-, -C₁₋₃alkyl-NH-CO-Het²⁰-, C₁₋₂alkyl-CO-Het²¹-CO-,

25 -Het²²-CH₂-CO-NH-C₁₋₃alkyl-, -CO-NH-C₁₋₆alkyl-, -NH-CO-C₁₋₆alkyl-,

-CO-C₁₋₇alkyl-, -C₁₋₇alkyl-CO-, -C₁₋₆alkyl-CO-C₁₋₆alkyl-,

-C₁₋₂alkyl-NH-CO-CR¹⁶R¹⁷-NH-, -C₁₋₂alkyl-CO-NH-CR¹⁸R¹⁹-CO-,

-C₁₋₂alkyl-CO-NR²⁰-C₁₋₃alkyl-CO-, -C₁₋₂alkyl-NR²¹-CH₂-CO-NH-C₁₋₃alkyl-, or

-NR²²-CO-C₁₋₃alkyl-NH-;

-4-

- X^1 represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹¹,
-NR¹¹-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or -C₁₋₂alkyl-;
- X^2 represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹²,
-NR¹²-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or -C₁₋₂alkyl-;
- 5 R^1 represents hydrogen, cyano, halo, hydroxy, formyl, C₁₋₆alkoxy-, C₁₋₆alkyl-,
halo-phenyl-carbonylamino-,
C₁₋₆alkoxy- substituted with halo,
C₁₋₄alkyl substituted with one or where possible two or more substituents selected
from hydroxy or halo;
- 10 R^2 represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-, Het¹⁶-carbonyl-,
C₁₋₄alkyloxycarbonyl-, C₁₋₄alkylcarbonyl-, aminocarbonyl-, mono-or
di(C₁₋₄alkyl)aminocarbonyl-, Het¹, formyl, C₁₋₄alkyl-, C₂₋₆alkynyl-,
C₃₋₆cycloalkyl-, C₃₋₆cycloalkyloxy-, C₁₋₆alkoxy-, Ar⁵, Ar¹-oxy-, dihydroxyborane ,
C₁₋₆alkoxy- substituted with halo,
15 C₁₋₄alkyl substituted with one or where possible two or more substituents selected
from halo, hydroxy or NR⁵R⁶,
C₁₋₄alkylcarbonyl- wherein said C₁₋₄alkyl is optionally substituted with one or
where possible two or more substituents selected from hydroxy or
C₁₋₄alkyl-oxy-;
- 20 R^3 represents hydrogen, C₁₋₄alkyl, or C₁₋₄alkyl substituted with one or more
substituents selected from halo, C₁₋₄alkyloxy-, amino-, mono-or
di(C₁₋₄alkyl)amino-, C₁₋₄alkyl-sulfonyl- or phenyl-;
- R^4 represents hydrogen, hydroxy, Ar³-oxy, Ar⁴-C₁₋₄alkyloxy-, C₁₋₄alkyloxy-,
C₂₋₄alkenyloxy- optionally substituted with Het¹² or R^4 represents C₁₋₄alkyloxy
25 substituted with one or where possible two or more substituents selected from
C₁₋₄alkyloxy-, hydroxy, halo, Het²-, -NR⁷R⁸-, -carbonyl- NR⁹R¹⁰ or
Het³-carbonyl-;
- R^5 and R^6 are each independently selected from hydrogen or C₁₋₄alkyl-;
- R^7 and R^8 are each independently selected from hydrogen, C₁₋₄alkyl, Het⁸,
30 aminosulfonyl-, mono- or di (C₁₋₄alkyl)-aminosulfonyl, hydroxy-C₁₋₄alkyl-,
C₁₋₄alkyl-oxy-C₁₋₄alkyl-, hydroxycarbonyl-C₁₋₄alkyl-, C₃₋₆cycloalkyl,
Het⁹-carbonyl-C₁₋₄alkyl-, Het¹⁰-carbonyl-, polyhydroxy-C₁₋₄alkyl-,
Het¹¹-C₁₋₄alkyl- or Ar²-C₁₋₄alkyl-;
- R^9 and R^{10} are each independently selected from hydrogen, C₁₋₄alkyl, C₃₋₆cycloalkyl,
35 Het⁴, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl- or polyhydroxy-C₁₋₄alkyl-;
- R^{11} represents hydrogen, C₁₋₄alkyl, Het⁵, Het⁶-C₁₋₄alkyl-, C₂₋₄alkenylcarbonyl-
optionally substituted with Het⁷-C₁₋₄alkylaminocarbonyl-, C₂₋₄alkenylsulfonyl-,

-5-

C₁₋₄alkyloxyC₁₋₄alkyl- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;

- 5 R¹² represents hydrogen, C₁₋₄alkyl, C₁₋₄alkyl-oxy-carbonyl-, Het¹⁸-C₁₋₄alkyl-, phenyl-C₁₋₄alkyl-oxy-carbonyl-, Het¹⁷, C₂₋₄alkenylcarbonyl- optionally substituted with Het¹⁹-C₁₋₄alkylaminocarbonyl-, C₂₋₄alkenylsulfonyl-, C₁₋₄alkyloxyC₁₋₄alkyl- or R¹² represents phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;
- 10 R¹³ represents hydrogen, C₁₋₄alkyl, Het¹³, Het¹⁴-C₁₋₄alkyl- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;
- R¹⁴ and R¹⁵ are each independently selected from hydrogen, C₁₋₄alkyl, Het¹⁵-C₁₋₄alkyl- or C₁₋₄alkyloxyC₁₋₄alkyl-;
- 15 R¹⁶ and R¹⁷ each independently represents hydrogen or C₁₋₄alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl, aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;
- R¹⁸ and R¹⁹ each independently represents hydrogen or C₁₋₄alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl,
- 20 aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;
- R²⁰ and R²² each independently represents hydrogen or C₁₋₄alkyl optionally substituted with hydroxy or C₁₋₄alkyloxy;
- R²¹ represents hydrogen, C₁₋₄alkyl, Het²³-C₁₋₄alkylcarbonyl- or R²¹ represents mono- or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-carbonyl- optionally
- 25 substituted with hydroxy, pyrimidinyl, dimethylamine or C₁₋₄alkyloxy;
- Het¹ represents a heterocycle selected from piperidinyl, morpholinyl, piperazinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het¹ is optionally substituted with amino, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl-, phenyl, phenyl-C₁₋₄alkyl-,
- 30 C₁₋₄alkyl-oxy-C₁₋₄alkyl- mono- or di(C₁₋₄alkyl)amino- or amino-carbonyl-;
- Het² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl, thiomorpholinyl or dithianyl wherein said Het² is optionally substituted with one or where possible two or more substituents selected from hydroxy, halo, amino, C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl-,
- 35 hydroxy-C₁₋₄alkyl-oxy-C₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino-, mono- or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-, aminoC₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino-sulfonyl-, aminosulfonyl-;

-6-


Het³, Het⁴ and Het⁸ each independently represent a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het³, Het⁴ or Het⁸ is optionally substituted with one or where possible two or
 5 more substituents selected from hydroxy-, amino-, C₁₋₄alkyl-, C₃₋₆cycloalkyl-C₁₋₄alkyl-, aminosulfonyl-, mono- or di(C₁₋₄alkyl)aminosulfonyl or amino-C₁₋₄alkyl-;

Het⁵ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from
 10 C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het⁶ and Het⁷ each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl,
 15 hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het⁹ and Het¹⁰ each independently represent a heterocycle selected from furanyl, piperidinyl, morpholinyl, piperazinyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het⁹ or Het¹⁰ is optionally substituted C₁₋₄alkyl, C₃₋₆cycloalkyl-C₁₋₄alkyl- or
 20 amino-C₁₋₄alkyl-;

Het¹¹ represents a heterocycle selected from indolyl or



Het¹² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl, thiomorpholinyl or dithianyl wherein said Het¹² is optionally substituted with one or where possible two or more substituents selected from
 25 hydroxy, halo, amino, C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-oxy-C₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino- or mono- or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-;

Het¹³ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from
 30 C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het¹⁴ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-,
 35 C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

-7-

- Het¹⁵ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;
- 5 Het¹⁶ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl, 1,3,2-dioxaborolane or piperidinyl wherein said heterocycle is optionally substituted with one or more substituents selected from C₁₋₄alkyl; and
- Het¹⁷ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from
- 10 C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;
- Het¹⁸ and Het¹⁹ each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl,
- 15 C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;
- Het²⁰, Het²¹ and Het²² each independently represent a heterocycle selected from pyrrolidinyl, 2-pyrrolidinonyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from hydroxy,
- 20 C₁₋₄alkyl, hydroxy-C₁₋₄alkyl- or polyhydroxy-C₁₋₄alkyl-;
- Het²³ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;
- 25 Ar¹, Ar², Ar³, Ar⁴ and Ar⁵ each independently represent phenyl optionally substituted with cyano, C₁₋₄alkylsulfonyl-, C₁₋₄alkylsulfonylamino-, aminosulfonylamino-, hydroxy-C₁₋₄alkyl, aminosulfonyl-, hydroxy-, C₁₋₄alkyloxy- or C₁₋₄alkyl.
- 30 As used in the foregoing definitions and hereinafter,
- halo is generic to fluoro, chloro, bromo and iodo;
 - C₁₋₂alkyl defines methyl or ethyl;
 - C₁₋₃alkyl defines straight and branched chain saturated hydrocarbon radicals having from 1 to 3 carbon atoms such as, for example, methyl, ethyl, propyl and the like;
 - 35 - C₁₋₄alkyl defines straight and branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as, for example, methyl, ethyl, propyl, butyl, 1-methylethyl, 2-methylpropyl, 2,2-dimethylethyl and the like;

-8-

- C₁₋₅alkyl defines straight and branched chain saturated hydrocarbon radicals having from 1 to 5 carbon atoms such as, for example, methyl, ethyl, propyl, butyl, pentyl, 1-methylbutyl, 2,2-dimethylpropyl, 2,2-dimethylethyl and the like;
- C₁₋₆alkyl is meant to include C₁₋₅alkyl and the higher homologues thereof having 6 carbon atoms such as, for example hexyl, 1,2-dimethylbutyl, 2-methylpentyl and the like;
- C₁₋₇alkyl is meant to include C₁₋₆alkyl and the higher homologues thereof having 7 carbon atoms such as, for example 1,2,3-dimethylbutyl, 1,2-methylpentyl and the like;
- C₃₋₉alkyl defines straight and branched chain saturated hydrocarbon radicals having from 3 to 9 carbon atoms such as propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl and the like;
- C₂₋₄alkenyl defines straight and branched chain hydrocarbon radicals containing one double bond and having from 2 to 4 carbon atoms such as, for example vinyl, 2-propenyl, 3-butenyl, 2-butenyl and the like;
- C₃₋₉alkenyl defines straight and branched chain hydrocarbon radicals containing one double bond and having from 3 to 9 carbon atoms such as, for example 2-propenyl, 3-butenyl, 2-butenyl, 2-pentenyl, 3-pentenyl, 3-methyl-2-butenyl, 3-hexenyl and the like;
- C₂₋₆alkynyl defines straight and branched chain hydrocarbon radicals containing one triple bond and having from 2 to 6 carbon atoms such as, for example, 2-propynyl, 3-butylnyl, 2-butylnyl, 2-pentylnyl, 3-pentylnyl, 3-methyl-2-butylnyl, 3-hexynyl and the like;
- C₃₋₆cycloalkyl is generic to cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl;
- C₁₋₄alkyloxy defines straight or branched saturated hydrocarbon radicals such as methoxy, ethoxy, propyloxy, butyloxy, 1-methylethyloxy, 2-methylpropyloxy and the like;
- C₁₋₆alkyloxy is meant to include C₁₋₄alkyloxy and the higher homologues such as methoxy, ethoxy, propyloxy, butyloxy, 1-methylethyloxy, 2-methylpropyloxy and the like;
- polyhydroxy-C₁₋₄alkyl is generic to a C₁₋₄alkyl as defined hereinbefore, having two, three or were possible more hydroxy substituents, such as for example trifluoromethyl.

As used in the foregoing definitions and hereinafter, the term formyl refers to a radical of formula -CH(=O) . When X¹ represent the divalent radical -O-N=CH- , said radical is attached with the carbon atom to the R³, R⁴ bearing cyclic moiety of the compounds of formula (I) and when X² represents the divalent radical -O-N=CH- ,

-9-

said radical is attached with the carbon atom to the R^1 , R^2 bearing phenyl moiety of the compounds of formula (I).

5 The heterocycles as mentioned in the above definitions and hereinafter, are meant to include all possible isomeric forms thereof, for instance pyrrolyl also includes 2*H*-pyrrolyl; triazolyl includes 1,2,4-triazolyl and 1,3,4-triazolyl; oxadiazolyl includes 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl and 1,3,4-oxadiazolyl; thiadiazolyl includes 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl and 1,3,4-thiadiazolyl; pyranyl includes 2*H*-pyranyl and 4*H*-pyranyl.

10 Further, the heterocycles as mentioned in the above definitions and hereinafter may be attached to the remainder of the molecule of formula (I) through any ring carbon or heteroatom as appropriate. Thus, for example, when the heterocycle is imidazolyl, it may be a 1-imidazolyl, 2-imidazolyl, 3-imidazolyl, 4-imidazolyl and 5-imidazolyl; when it is thiazolyl, it may be 2-thiazolyl, 4-thiazolyl and 5-thiazolyl; when it is
15 triazolyl, it may be 1,2,4-triazol-1-yl, 1,2,4-triazol-3-yl, 1,2,4-triazol-5-yl, 1,3,4-triazol-1-yl and 1,3,4-triazol-2-yl; when it is benzothiazolyl, it may be 2-benzothiazolyl, 4-benzothiazolyl, 5-benzothiazolyl, 6-benzothiazolyl and 7-benzothiazolyl.

20 The pharmaceutically acceptable addition salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic acid addition salt forms which the compounds of formula (I) are able to form. The latter can conveniently be obtained by treating the base form with such appropriate acid. Appropriate acids comprise, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic
25 acid; sulfuric; nitric; phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic, malonic, succinic (i.e. butane-dioic acid), maleic, fumaric, malic, tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, pamoic and the like acids.

30

The pharmaceutically acceptable addition salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic base addition salt forms which the compounds of formula (I) are able to form. Examples of such base addition salt forms are, for example, the sodium, potassium, calcium salts, and also the salts with
35 pharmaceutically acceptable amines such as, for example, ammonia, alkylamines, benzathine, *N*-methyl-*D*-glucamine, hydrabamine, amino acids, e.g. arginine, lysine.

-10-

Conversely said salt forms can be converted by treatment with an appropriate base or acid into the free acid or base form.

- 5 The term addition salt as used hereinabove also comprises the solvates which the compounds of formula (I) as well as the salts thereof, are able to form. Such solvates are for example hydrates, alcoholates and the like.

- 10 The term stereochemically isomeric forms as used hereinbefore defines the possible different isomeric as well as conformational forms which the compounds of formula (I) may possess. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture of all possible stereochemically and conformationally isomeric forms, said mixtures containing all diastereomers, enantiomers and/or conformers of the basic molecular structure. All stereochemically
15 isomeric forms of the compounds of formula (I) both in pure form or in admixture with each other are intended to be embraced within the scope of the present invention.

- Some of the compounds of formula (I) may also exist in their tautomeric forms. Such forms although not explicitly indicated in the above formula are intended to be
20 included within the scope of the present invention.

- The *N*-oxide forms of the compounds of formula (I) are meant to comprise those compounds of formula (I) wherein one or several nitrogen atoms are oxidized to the so-called *N*-oxide.

- 25 A first group of compounds according to the present invention consists of those compounds of formula (I) wherein one or more of the following restrictions apply;

Z represents O, NH or S;

- Y represents -C₃₋₉alkyl-, -C₃₋₉alkenyl-, -C₁₋₅alkyl-oxy-C₁₋₅alkyl-,
30 -C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-, -C₁₋₅alkyl-NR¹⁴-CO-C₁₋₅alkyl-,
-C₁₋₅alkyl-CO-NR¹⁵-C₁₋₅alkyl-, -C₁₋₆alkyl-CO-NH-, -C₁₋₆alkyl-NH-CO-,
-CO-NH-C₁₋₆alkyl-, -NH-CO-C₁₋₆alkyl-, -CO-C₁₋₇alkyl-, -C₁₋₇alkyl-CO-,
-C₁₋₆alkyl-CO-C₁₋₆alkyl-, -C₁₋₂alkyl-NH-CO-CHR¹⁶-NH-;

- X¹ represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹¹,
35 -NR¹¹-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or -C₁₋₂alkyl-;

X² represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹²,
-NR¹²-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or -C₁₋₂alkyl-;

-11-

- R^1 represents hydrogen, cyano, halo, hydroxy, formyl, C_{1-6} alkoxy-, C_{1-6} alkyl-,
 C_{1-6} alkoxy- substituted with halo,
 C_{1-4} alkyl substituted with one or where possible two or more substituents selected
from hydroxy or halo;
- 5 R^2 represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-, Het^{16} -carbonyl-,
 C_{1-4} alkyloxycarbonyl-, C_{1-4} alkylcarbonyl-, aminocarbonyl-, mono-or
di(C_{1-4} alkyl)aminocarbonyl-, Het^1 , formyl, C_{1-4} alkyl-, C_{2-6} alkynyl-,
 C_{3-6} cycloalkyl-, C_{3-6} cycloalkyloxy-, C_{1-6} alkoxy-, Ar^5 , Ar^1 -oxy-, dihydroxyborane,
 C_{1-6} alkoxy- substituted with halo,
10 C_{1-4} alkyl substituted with one or where possible two or more substituents selected
from halo, hydroxy or NR^5R^6 ,
 C_{1-4} alkylcarbonyl- wherein said C_{1-4} alkyl is optionally substituted with one or
where possible two or more substituents selected from hydroxy or
 C_{1-4} alkyl-oxy-;
- 15 R^3 represents hydrogen, C_{1-4} alkyl, or C_{1-4} alkyl substituted with one or more
substituents selected from halo, C_{1-4} alkyloxy-, amino-, mono-or
di(C_{1-4} alkyl)amino-,
 C_{1-4} alkyl-sulfonyl- or phenyl;
- R^4 represents hydrogen, hydroxy, Ar^3 -oxy, Ar^4 - C_{1-4} alkyloxy-, C_{1-4} alkyloxy-,
20 C_{2-4} alkenyloxy- optionally substituted with Het^{12} or R^4 represents C_{1-4} alkyloxy
substituted with one or where possible two or more substituents selected from
 C_{1-4} alkyloxy-, hydroxy, halo, Het^2 -, $-NR^7R^8$ -, -carbonyl- NR^9R^{10} or
 Het^3 -carbonyl-;
- R^5 and R^6 are each independently selected from hydrogen or C_{1-4} alkyl;
- 25 R^7 and R^8 are each independently selected from hydrogen, C_{1-4} alkyl, Het^8 ,
aminosulfonyl-, mono- or di (C_{1-4} alkyl)-aminosulfonyl, hydroxy- C_{1-4} alkyl-,
 C_{1-4} alkyl-oxy- C_{1-4} alkyl-, hydroxycarbonyl- C_{1-4} alkyl-, C_{3-6} cycloalkyl,
 Het^9 -carbonyl- C_{1-4} alkyl-, Het^{10} -carbonyl-, polyhydroxy- C_{1-4} alkyl-,
 Het^{11} - C_{1-4} alkyl- or Ar^2 - C_{1-4} alkyl-;
- 30 R^9 and R^{10} are each independently selected from hydrogen, C_{1-4} alkyl, C_{3-6} cycloalkyl,
 Het^4 , hydroxy- C_{1-4} alkyl-, C_{1-4} alkyloxy- C_{1-4} alkyl- or polyhydroxy- C_{1-4} alkyl-;
- R^{11} represents hydrogen, C_{1-4} alkyl, Het^5 , Het^6 - C_{1-4} alkyl-, C_{2-4} alkenylcarbonyl-
optionally substituted with Het^7 - C_{1-4} alkylaminocarbonyl-, C_{2-4} alkenylsulfonyl-,
 C_{1-4} alkyloxy- C_{1-4} alkyl- or phenyl optionally substituted with one or where
35 possible two or more substituents selected from hydrogen, hydroxy, amino or
 C_{1-4} alkyloxy-;


-12-

- R¹² represents hydrogen, C₁₋₄alkyl, C₁₋₄alkyl-oxy-carbonyl-, Het¹⁷, Het¹⁸-C₁₋₄alkyl-, C₂₋₄alkenylcarbonyl- optionally substituted with Het¹⁹-C₁₋₄alkylaminocarbonyl-, C₂₋₄alkenylsulfonyl-, C₁₋₄alkyloxyC₁₋₄alkyl- or R¹² represents phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;
- R¹³ represents hydrogen, C₁₋₄alkyl, Het¹³, Het¹⁴-C₁₋₄alkyl- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;
- R¹⁴ and R¹⁵ are each independently selected from hydrogen, C₁₋₄alkyl, Het¹⁵-C₁₋₄alkyl- or C₁₋₄alkyloxyC₁₋₄alkyl-;
- R¹⁶ represents hydrogen or C₁₋₄alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl, aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;
- Het¹ represents a heterocycle selected from piperidinyl, morpholinyl, piperazinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het¹ is optionally substituted with amino, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl-, phenyl, phenyl-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl- mono- or di(C₁₋₄alkyl)amino- or amino-carbonyl-;
- Het² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl, thiomorpholinyl or dithianyl wherein said Het² is optionally substituted with one or where possible two or more substituents selected from hydroxy, halo, amino, C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-oxy-C₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino-, mono- or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-, aminoC₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino-sulfonyl-, aminosulfonyl-;
- Het³, Het⁴ and Het⁸ each independently represent a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het³, Het⁴ or Het⁸ is optionally substituted with one or where possible two or more substituents selected from hydroxy-, amino-, C₁₋₄alkyl-, C₃₋₆cycloalkyl-C₁₋₄alkyl-, aminosulfonyl-, mono- or di(C₁₋₄alkyl)aminosulfonyl or amino-C₁₋₄alkyl-;
- Het⁵ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

-13-

Het⁶ and Het⁷ each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

- 5 Het⁹ and Het¹⁰ each independently represent a heterocycle selected from furanyl, piperidinyl, morpholinyl, piperazinyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het⁹ or Het¹⁰ is optionally substituted C₁₋₄alkyl, C₃₋₆cycloalkyl-C₁₋₄alkyl- or amino-C₁₋₄alkyl-;



- 10 Het¹¹ represents a heterocycle selected from indolyl or
 Het¹² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl, thiomorpholinyl or dithianyl wherein said Het¹² is optionally substituted with one or where possible two or more substituents selected from hydroxy, halo, amino, C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl-,
 15 hydroxy-C₁₋₄alkyl-oxy-C₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino- or mono- or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-;

- Het¹³ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or
 20 polyhydroxy-C₁₋₄alkyl-;

- Het¹⁴ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

- 25 Het¹⁵ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

- 30 Het¹⁶ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl, 1,3,2-dioxaborolane or piperidinyl wherein said heterocycle is optionally substituted with one or more substituents selected from C₁₋₄alkyl; and

- Het¹⁷ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or
 35 polyhydroxy-C₁₋₄alkyl-;

-14-

- Het¹⁸ and Het¹⁹ each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl,
- 5 hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxy-C₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;
- Ar¹, Ar², Ar³, Ar⁴ and Ar⁵ each independently represent phenyl optionally substituted with cyano, C₁₋₄alkylsulfonyl-, C₁₋₄alkylsulfonylamino-, aminosulfonylamino-, hydroxy-C₁₋₄alkyl, aminosulfonyl-, hydroxy-, C₁₋₄alkyloxy- or C₁₋₄alkyl.
- 10 An interesting group of compounds consists of those compounds of formula (I) wherein one or more of the following restrictions apply :
- Z represents NH;
- Y represents -C₃₋₉alkyl-, -C₂₋₉alkenyl-, -C₁₋₅alkyl-oxy-C₁₋₅alkyl-, -C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-, -C₁₋₅alkyl-NR¹⁴-CO-C₁₋₅alkyl-, -C₁₋₆alkyl-NH-CO-,
- 15 -NH-CO-C₁₋₆alkyl-, -CO-C₁₋₇alkyl-, -C₁₋₇alkyl-CO-, C₁₋₆alkyl-CO-C₁₋₆alkyl-, -C₁₋₂alkyl-NH-CO-CR¹⁶R¹⁷-NH-, -C₁₋₂alkyl-CO-NH-CR¹⁸R¹⁹-CO-, -C₁₋₂alkyl-CO-NR²⁰-C₁₋₃alkyl-CO-, -C₁₋₂alkyl-NR²¹-CH₂-CO-NH-C₁₋₃alkyl-, -NR²²-CO-C₁₋₃alkyl-NH-, -C₁₋₃alkyl-NH-CO-Het²⁰-, C₁₋₂alkyl-CO-Het²¹-CO-, or -Het²²-CH₂-CO-NH-C₁₋₃alkyl-;
- 20 X¹ represents O, -O-C₁₋₂alkyl-, -O-N=CH-, NR¹¹ or -NR¹¹-C₁₋₂alkyl-; in a particular embodiment X¹ represents -NR¹¹-, -O- or -O-CH₂-;
- X² represents a direct bond, O, -O-C₁₋₂alkyl-, -O-N=CH-, NR¹² or NR¹²-C₁₋₂alkyl-; in a particular embodiment X² represents a direct bond, -C₁₋₂alkyl-, -O-C₁₋₂alkyl-, -O- or -O-CH₂-;
- 25 R¹ represents hydrogen, cyano, halo or hydroxy, preferably halo;
- R² represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-, C₁₋₄alkyloxycarbonyl-, Het¹⁶-carbonyl-, C₁₋₄alkyl-, C₂₋₆alkynyl-, Ar⁵ or Het¹;
- In a further embodiment R² represents hydrogen, cyano, halo, hydroxy, C₂₋₆alkynyl- or Het¹; in particular R² represents hydrogen, cyano, halo, hydroxy,
- 30 or Ar⁵;
- R³ represents hydrogen;
- R⁴ represents hydrogen, hydroxy, C₁₋₄alkyloxy- or R⁴ represents C₁₋₄alkyloxy substituted with one or where possible two or more substituents selected from C₁₋₄alkyloxy- or Het²-;
- 35 R¹² represents hydrogen, C₁₋₄alkyl- or C₁₋₄alkyl-oxy-carbonyl-;
- R¹³ represents hydrogen or Het¹⁴-C₁₋₄alkyl, in particular morpholinyl-C₁₋₄alkyl;
- R¹⁴ and R¹⁵ represents hydrogen;

-15-

- R¹⁶ represents hydrogen or C₁₋₄alkyl substituted with hydroxy;
 R¹⁷ represents hydrogen or C₁₋₄alkyl, in particular hydrogen or methyl;
 R¹⁸ represents hydrogen or C₁₋₄alkyl optionally substituted with hydroxy or phenyl;
 R¹⁹ represents hydrogen or C₁₋₄alkyl, in particular hydrogen or methyl, even more
 5 particular hydrogen;
 R²⁰ represents hydrogen or C₁₋₄alkyl, in particular hydrogen or methyl;
 R²¹ represents hydrogen, C₁₋₄alkyl, Het²³-C₁₋₄alkylcarbonyl- or
 R²¹ represents mono-or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-carbonyl- optionally
 substituted with hydroxy, pyrimidinyl, dimethylamine or C₁₋₄alkyloxy;
 10 R²² represents hydrogen or C₁₋₄alkyl optionally substituted with hydroxy or
 C₁₋₄alkyloxy;
 Het¹ represents thiazolyl optionally substituted with amino, C₁₋₄alkyl, hydroxy-C₁₋₄
 alkyl-, phenyl, phenyl-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl- mono- or di(C₁₋₄
 alkyl)amino- or amino-carbonyl-;
 15 Het² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or
 pyrrolidinyl wherein said Het² is optionally substituted with one or where
 possible two or more substituents selected from hydroxy, amino or C₁₋₄alkyl-;
 In a further embodiment Het² represents a heterocycle selected from morpholinyl
 or piperidinyl optionally substituted with C₁₋₄alkyl-, preferably methyl;
 20 Het³ represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or
 pyrrolidinyl wherein said Het³ is optionally substituted with one or where
 possible two or more substituents selected from hydroxy, amino or C₁₋₄alkyl-;
 Het¹² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or
 pyrrolidinyl wherein said Het¹² is optionally substituted with one or where
 25 possible two or more substituents selected from hydroxy, amino or C₁₋₄alkyl-;
 Het¹⁶ represents a heterocycle selected from piperidinyl or pyrrolidinyl;
 Het²⁰ represents pyrrolidinyl, 2-pyrrolidinonyl, piperidinyl or hydroxy-pyrrolidinyl,
 preferably pyrrolidinyl or hydroxy-pyrrolidinyl;
 Het²¹ represents pyrrolidinyl or hydroxy-pyrrolidinyl;
 30 Het²² represents pyrrolidinyl, piperazinyl or piperidinyl.

A particular group of compounds consists of those compounds of formula (I) wherein
 one or more of the following restrictions apply :

Z represents NH;

- 35 Y represents -C₃₋₉alkyl-, -C₂₋₉alkenyl-, -C₃₋₇alkyl-CO-NH optionally substituted with
 amino, mono - or di(C₁₋₄alkyl)amino or C₁₋₄alkyloxycarbonylamino-,
 -C₃₋₇alkenyl-CO-NH- optionally substituted with amino, mono- or

-16-

- di(C₁₋₄alkyl)amino- or C₁₋₄alkyloxycarbonylamino-,
 C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-, -C₁₋₅alkyl-NR¹⁴-CO-C₁₋₅alkyl-, -C₁₋₆alkyl-CO-NH-,
 -C₁₋₅alkyl-CO-NR¹⁵-C₁₋₅alkyl-, -C₁₋₃alkyl-NH-CO-Het²⁰-,
 -C₁₋₂alkyl-CO-Het²¹-CO-, -C₁₋₂alkyl-NH-CO-CR¹⁶R¹⁷-NH-,
 5 -C₁₋₂alkyl-CO-NH-CR¹⁸R¹⁹-CO-, -C₁₋₂alkyl-CO-NR²⁰-C₁₋₃alkyl-CO-, or
 -NR²²-CO-C₁₋₃alkyl-NH-;
 even more particular Y represents -C₃₋₉alkyl-, -C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-,
 -C₁₋₅alkyl-NR¹⁴-CO-C₁₋₅alkyl-, -C₁₋₃alkyl-NH-CO-Het²⁰-,
 -C₁₋₂alkyl-CO-Het²¹-CO-, or -C₁₋₂alkyl-NH-CO-CR¹⁶R¹⁷-NH-;
 10 X¹ represents a direct bond, O or -O-C₁₋₂alkyl-;
 X² represents a direct bond, -CO-C₁₋₂alkyl-, NR¹²-, -NR¹²-C₁₋₂alkyl-, -O-N=CH- or
 -C₁₋₂alkyl-; even more particular X² represents -CO-C₁₋₂alkyl- or
 NR¹²-C₁₋₂alkyl-;
 R¹ represents hydrogen or halo, preferably hydrogen, chloro, fluoro or bromo;
 15 R² represents hydrogen or halo, preferably hydrogen, chloro, fluoro or bromo;
 R³ represents hydrogen;
 R⁴ represents hydrogen or C₁₋₄alkyloxy, preferably C₁₋₄alkyloxy, even more
 preferably methoxy;
 R¹² represents hydrogen or C₁₋₄alkyl, preferably hydrogen or methyl;
 20 R¹³ represents hydrogen or C₁₋₄alkyl;
 R¹⁴ represents hydrogen;
 R¹⁵ represents hydrogen;
 R¹⁶ and R¹⁷ each independently represent hydrogen or C₁₋₄alkyl;
 R¹⁸ and R¹⁹ each independently represent hydrogen or C₁₋₄alkyl optionally substituted
 25 with phenyl or hydroxy;
 R²⁰ and R²¹ each independently represent hydrogen or C₁₋₄alkyl optionally substituted
 with C₁₋₄alkyloxy;
 Het²⁰, Het²¹ and Het²² each independently represent a heterocycle selected from the
 group consisting pyrrolidinyl, 2-pyrrolidinonyl or piperidinyl optionally
 30 substituted with hydroxy.

A preferred group of compounds consists of those compounds of formula (I) wherein
 one or more of the following restrictions apply :

Z represents NH;

- 35 Y represents -C₃₋₉alkyl-, -C₂₋₉alkenyl-, -C₁₋₅alkyl-oxy-C₁₋₅alkyl-,
 -C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-, -C₁₋₆alkyl-NH-CO-, -CO-C₁₋₇alkyl-, -C₁₋₇alkyl-CO- or
 C₁₋₆alkyl-CO-C₁₋₆alkyl;alkyl-NH-CO-Het²⁰-, -C₁₋₂alkyl-CO-Het²¹-CO-,

-17-

$-C_{1-2}alkyl-NH-CO-CR^{16}R^{17}-NH-$, $-C_{1-2}alkyl-CO-NH-CR^{18}R^{19}-CO-$, $-C_{1-2}alkyl-CO-NR^{20}-C$

X^1 represents O, $-O-C_{1-2}alkyl-$, $-O-N=CH-$, NR^{11} or $-NR^{11}-C_{1-2}alkyl-$; in a particular embodiment X^1 represents $-NR^{11}-$, $-O-$ or $-O-CH_2-$;

5 X^2 represents a direct bond, O, $-O-C_{1-2}alkyl-$, $-O-N=CH-$, NR^{12} or $NR^{12}-C_{1-2}alkyl-$; in a particular embodiment X^2 represents a direct bond, $-C_{1-2}alkyl-$, $-O-C_{1-2}alkyl-$, $-O-$ or $-O-CH_2-$;

R^1 represents hydrogen, cyano, halo or hydroxy, preferably halo;

R^2 represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-,

10 $C_{1-4}alkyloxycarbonyl-$, $Het^{16}-carbonyl-$, $C_{1-4}alkyl-$, $C_{2-6}alkynyl-$, Ar^5 or Het^1 ;

In a further embodiment R^2 represents hydrogen, cyano, halo, hydroxy,

$C_{2-6}alkynyl-$ or Het^1 ; in particular R^2 represents hydrogen, cyano, halo, hydroxy, or Ar^5 ;

R^3 represents hydrogen;

15 R^4 represents hydrogen, hydroxy, $C_{1-4}alkyloxy-$ or R^4 represents $C_{1-4}alkyloxy$ substituted with one or where possible two or more substituents selected from $C_{1-4}alkyloxy-$ or Het^2- ;

R^{12} represents hydrogen, $C_{1-4}alkyl-$ or $C_{1-4}alkyl-oxy-carbonyl-$;

R^{13} represents $Het^{14}-C_{1-4}alkyl$, in particular morpholinyl- $C_{1-4}alkyl$;

20 Het^1 represents thiazolyl optionally substituted with amino, $C_{1-4}alkyl$, hydroxy- $C_{1-4}alkyl-$, phenyl, phenyl- $C_{1-4}alkyl-$, $C_{1-4}alkyl-oxy-C_{1-4}alkyl-$ mono- or di($C_{1-4}alkyl$)amino- or amino-carbonyl-;

Het^2 represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or pyrrolidinyl wherein said Het^2 is optionally substituted with one or where

25 possible two or more substituents selected from hydroxy, amino or $C_{1-4}alkyl-$;

In a further embodiment Het^2 represents a heterocycle selected from morpholinyl

or piperidinyl optionally substituted with $C_{1-4}alkyl-$, preferably methyl;

Het^3 represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or

30 pyrrolidinyl wherein said Het^3 is optionally substituted with one or where possible two or more substituents selected from hydroxy, amino or $C_{1-4}alkyl-$;

Het^{12} represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or

pyrrolidinyl wherein said Het^{12} is optionally substituted with one or where possible two or more substituents selected from hydroxy, amino or $C_{1-4}alkyl-$;

Het^{16} represents a heterocycle selected from piperidinyl or pyrrolidinyl.

35

A further group of compounds consists of those compounds of formula (I) wherein one or more of the following restrictions apply :

-18-

Z represents NH;

Y represents -C₃₋₉alkyl-, -C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-, -C₁₋₆alkyl-NH-CO-,
-CO-C₁₋₇alkyl- or -C₁₋₇alkyl-CO-;

X¹ represents -NR¹¹-, -O- or -O-CH₂-;

5 X² represents a direct bond, -NR¹²-, -NR¹²-C₁₋₂alkyl-, -CO-, -O- or -O-CH₂-;

R¹ represents halo; in particular R¹ represents chloro, fluoro or bromo and is at
position 5';

R² represents hydrogen, cyano, halo, hydroxy, or Ar⁵;

R³ represents hydrogen;

10 R⁴ represents C₁₋₄alkyloxy substituted with one or where possible two or more
substituents selected from C₁₋₄alkyloxy- or Het²-;

R¹² represents C₁₋₄alkyl or R¹² represents C₁₋₄alkyl-oxy-carbonyl;

R¹³ represents Het¹⁴-C₁₋₄alkyl;

15 Het² represents a heterocycle selected from morpholinyl or piperidinyl optionally
substituted with C₁₋₄alkyl-;

Het³ represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or
pyrrolidinyl wherein said Het³ is optionally substituted with one or where
possible two or more substituents selected from hydroxy, amino or C₁₋₄alkyl-;

20 Het¹² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or
pyrrolidinyl wherein said Het¹² is optionally substituted with one or where
possible two or more substituents selected from hydroxy, amino or C₁₋₄alkyl-;

Het¹⁴ represents morpholinyl.

25 In a further embodiment of the present invention the compounds of formula (I) are
selected from the group consisting of ;

4,6-ethanediylidenepyrimido[4,5-b][6,1,12]benzoxadiazacyclopentadecine, 17-
bromo-8,9,10,11,12,13,14,19-octahydro-20-methoxy-

4,6-ethanediylidenepyrimido[4,5-b][6,1,12]benzoxadiazacyclopentadecine, 17-
bromo-8,9,10,11,12,13,14,19-octahydro-20-methoxy-13-methyl-

benzamide, 4-fluoro-N-(8,9,10,11,12,13-hexahydro-20-methoxy-4,6-
ethanediylidene-19H-pyrimido[4,5-b][6,13,1]benzodioxazacyclopentadecin-
16-yl)-

4,6-ethanediylidene-8H,14H-pyrimido[4,5-
b][6,12,1]benzodioxazacyclohexadecine, 18-chloro-9,10,11,12,15,20-
hexahydro-21-methoxy-

4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,11]benzoxadiazacyclohexadecin-

-19-

11(12H)-one, 18-chloro-9,10,13,14,15,20-hexahydro-21-methoxy-
4,6-ethanediylidene-14H-pyrimido[4,5-b][6,9,12,1]benzotrioxaazacyclohexadecine,
18-chloro-8,9,11,12,15,20-hexahydro-21-methoxy-
4,6-ethanediylidenepyrimido[4,5-b]pyrrolo[2,1-
k][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 19-chloro-
9,10,11a,12,13,14,16,21-octahydro-22-methoxy-
4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-
one, 17-chloro-9,10,12,13,14,19-hexahydro-20-methoxy-13-methyl-
4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-
one, 17-chloro-9,10,12,13,14,19-hexahydro-20-methoxy-
4,6-ethanediylidene-12H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclohexadecin-
12-one, 18-chloro-8,9,10,11,13,14,15,20-octahydro-21-methoxy-14-methyl-
4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,13]benzoxatriazacyclohexadecin-
11(12H)-one, 18-chloro-9,10,13,14,15,20-hexahydro-21-methoxy-
4,6-ethanediylidenepyrimido[4,5-b][6,1,10,14]benzoxatriazacycloheptadecin-
12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-22-methoxy-
4,6-ethenopyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8H,13H)-
dione, 17-chloro-10,11,14,19-tetrahydro-20-methoxy-
4,6-etheno-8H-pyrimido[4,5-b]pyrrolo[2,1-
l][6,1,10,13]benzoxatriazacyclohexadecine-12,15(14H)-dione, 20-chloro-
9,10,11,12a,13,17,22-heptahydro-23-methoxy-
4,6-ethanediylidene-12H-pyrimido[4,5-b]pyrrolo[2,1-
l][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 20-chloro-
8,9,10,11,12a,13,14,15,17,22-decahydro-23-methoxy-
4,6-ethenopyrimido[4,5-b][6,1,9,14]benzoxatriazacycloheptadecine-9,14(8H,15H)-
dione, 19-chloro-10,11,12,13,16,21-hexahydro-22-methoxy-
4,6-etheno-8H-pyrimido[4,5-b][6,1,9,13]benzoxatriazacyclohexadecine-
9,13(10H,14H)-dione, 18-chloro-11,12,15,20-tetrahydro-21-methoxy-
4,6-ethenopyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine-
11,14(8H,15H)-dione, 19-chloro-9,10,12,13,16,21-hexahydro-22-methoxy-
4,6-ethenopyrimido[4,5-b][6,1,11,16]benzoxatriazacyclononadecine-
11,16(8H,17H)-dione, 21-chloro-9,10,12,13,14,15,18,23-octahydro-24-
methoxy-
4,6-etheno-8H-pyrimido[4,5-b][6,1,11,15]benzoxatriazacyclooctadecine-
11,15(12H,16H)-dione, 20-chloro-9,10,13,14,17,22-hexahydro-23-methoxy-
4,6-ethenopyrimido[4,5-b][6,1,12]benzoxadiazacyclopentadecine, 17-bromo-16-
fluoro-8,9,10,11,12,13,14,19-octahydro-20-methoxy-

-20-

- 4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8H,13H)-dione, 17-chloro-10,11,14,19-tetrahydro-20-methoxy-11-methyl-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8H,13H)-dione, 17-chloro-10,11,14,19-tetrahydro-20-methoxy-11-(1-methylethyl)-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8H,13H)-dione, 17-chloro-10,11,14,19-tetrahydro-20-methoxy-11-(phenylmethyl)-
- 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-11,14-dione, 18-chloro-9,10,12,13,15,20-hexahydro-21-methoxy-12-(1-methylethyl)-
- 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-11,14-dione, 18-chloro-9,10,12,13,15,20-hexahydro-21-methoxy-12,12-dimethyl-
- 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-11,14-dione, 18-chloro-9,10,12,13,15,20-hexahydro-21-methoxy-12-(2-methylpropyl)-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,10,13]benzoxatriazacycloheptadecine-12,15-dione, 19-chloro-8,9,10,11,13,14,16,21-octahydro-22-methoxy-13-(2-methylpropyl)-
- 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-11,14-dione, 18-chloro-9,10,12,13,15,20-hexahydro-21-methoxy-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,10,13]benzoxatriazacycloheptadecine-12,15-dione, 19-chloro-8,9,10,11,13,14,16,21-octahydro-22-methoxy-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8H,13H)-dione, 10,11,14,19-tetrahydro-20-methoxy-11-methyl-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8H,13H)-dione, 10,11,14,19-tetrahydro-20-methoxy-11-(1-methylpropyl)-
- 9,11-ethanediylidenepyrimido[4,5-b]pyrrolo[1,2-i][6,1,9,12]benzoxatriazacyclopentadecine-14,19(5H,13H)-dione, 16,17,18,18a,20,21-hexahydro-22-methoxy-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8H,13H)-dione, 10,11,14,19-tetrahydro-20-methoxy-
- 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,13]benzoxatriazacyclohexadecine-9,13(10H,14H)-dione, 11,12,15,20-tetrahydro-21-methoxy-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,9,14]benzoxatriazacycloheptadecine-9,14(8H,15H)-dione, 10,11,12,13,16,21-hexahydro-22-methoxy-

-21-

- 4,6-ethanediylidenepyrimido[4,5-b]pyrrolo[2,1-k][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 19-chloro-18-fluoro-9,10,11a,12,13,14,16,21-octahydro-22-methoxy-
- 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclohexadecine, 18-chloro-9,10,11,12,13,14,15,20-octahydro-21-methoxy-14-methyl-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 17-chloro-16-fluoro-9,10,12,13,14,19-hexahydro-20-methoxy-13-methyl-
- 4,6-ethanediylidene-12H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 18-chloro-17-fluoro-8,9,10,11,13,14,15,20-octahydro-21-methoxy-14-methyl-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 17-chloro-16-fluoro-9,10,12,13,14,19-hexahydro-20-methoxy-
- 4,6-ethanediylidene-12H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 18-chloro-17-fluoro-8,9,10,11,13,14,15,20-octahydro-21-methoxy-
- 4,6-ethanediylidene-12H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 18-chloro-8,9,10,11,13,14,15,20-octahydro-21-methoxy-
- 9,11-ethanediylidenepyrimido[4,5-b]pyrrolo[1,2-i][6,1,9,12]benzoxatriazacyclopentadecine-14,19(5H,13H)-dione, 3-chloro-16,17,18,18a,20,21-hexahydro-22-methoxy-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8H,13H)-dione, 17-chloro-10,11,14,19-tetrahydro-20-methoxy-10-methyl-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8H,13H)-dione, 17-chloro-10,11,14,19-tetrahydro-11-(1-hydroxyethyl)-20-methoxy-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine-11,14(8H,15H)-dione, 19-chloro-9,10,12,13,16,21-hexahydro-22-methoxy-13-(1-methylpropyl)-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8H,13H)-dione, 17-chloro-10,11,14,19-tetrahydro-11-(hydroxymethyl)-20-methoxy-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine-11,14(8H,15H)-dione, 19-chloro-9,10,12,13,16,21-hexahydro-13-(hydroxymethyl)-22-methoxy-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine-11,14(8H,15H)-dione, 19-chloro-9,10,12,13,16,21-hexahydro-22-methoxy-13-methyl-
- 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-

-22-

9,12-dione, 18-chloro-10,11,13,14,15,20-hexahydro-21-methoxy-4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,14]benzoxatriazacyclooctadecine-9,14-dione, 20-chloro-10,11,12,13,15,16,17,22-octahydro-23-methoxy-4,6-ethanediylidenepyrimido[4,5-b]pyrrolo[2,1-k][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 19-chloro-9,10,11a,12,13,14,16,21-octahydro-22-methoxy-4,6-ethanediylidene-12H-pyrimido[4,5-b]pyrrolo[2,1-l][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 20-chloro-8,9,10,11,12a,13,14,15,17,22-decahydro-23-methoxy-4,6-ethanediylidene-12H-pyrimido[4,5-b]pyrrolo[2,1-l][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 20-chloro-19-fluoro-8,9,10,11,12a,13,14,15,17,22-decahydro-23-methoxy-4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecin-13(8H)-one, 19-chloro-9,10,11,12,14,15,16,21-octahydro-22-methoxy-4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-22-methoxy-10-methyl-4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-12(13H)-one, 19-chloro-10-ethyl-8,9,10,11,14,15,16,21-octahydro-22-methoxy-1,22-ethanediylidene-5H,17H-pyrimido[4,5-b]pyrrolo[2,1-h][6,1,9,12]benzoxatriazacycloheptadecin-14(15H)-one, 7-chloro-10,11,12,13,18,19,19a,20-octahydro-24-methoxy-4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclooctadecin-13(14H)-one, 20-chloro-9,10,11,12,15,16,17,22-octahydro-23-methoxy-14H-4,6-ethanediylidene-9,13-methano-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacycloeicosin-15(16H)-one, 22-chloro-9,10,11,12,17,18,19,24-octahydro-26-methoxy-13H-4,6-ethanediylidene-9,12-ethanopyrimido[4,5-b][6,1,11,14]benzoxatriazacyclononadecin-14(15H)-one, 21-chloro-8,9,10,11,16,17,18,23-octahydro-26-methoxy-14H-4,6-ethanediylidene-10,13-ethano-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacycloeicosin-15(16H)-one, 22-chloro-9,10,11,12,17,18,19,24-octahydro-27-methoxy-4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-10-[[[2-hydroxy-1-(hydroxymethyl)ethyl]amino]acetyl]-22-methoxy-4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-

-23-

- 12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-10-[[2-(hydroxymethyl)-4-morpholinyl]acetyl]-22-methoxy-4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-
- 12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-22-methoxy-4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-
- 12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-10-[[2-(hydroxyethyl)methylamino]acetyl]-22-methoxy-4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-
- 12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-22-methoxy-10-[[[2-(4-pyridinyl)ethyl]amino]acetyl]-4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-
- 12(13H)-one, 19-chloro-10-[[[2-(dimethylamino)ethyl]methylamino]acetyl]-8,9,10,11,14,15,16,21-octahydro-22-methoxy-4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-
- 12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-22-methoxy-10-[[[2-methoxyethyl]amino]acetyl]-4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-
- 12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-22-methoxy-10-[[[3-methoxypropyl]amino]acetyl]-4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-
- 12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-22-methoxy-10-(4-morpholinylacetyl)-4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-
- 12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-22-methoxy-10-[(4-methyl-1-piperazinyl)acetyl]-4,6-ethenopyrimido[4,5-b][6,1,12]benzoxadiazacyclopentadecine-13(8H)-carboxylic acid, 17-bromo-9,10,11,12,14,19-hexahydro-20-methoxy-, phenylmethyl ester
- 4,6-ethanediylidenepyrimido[4,5-b]pyrrolo[2,1-m][6,1,11,14]benzoxatriazacycloheptadecin-13(8H)-one, 21-chloro-9,10,11,12,13a,14,15,16,18,23-decahydro-24-methoxy-4,6-ethanediylidenepyrimido[4,5-b]pyrrolo[2,1-m][6,1,11,14]benzoxatriazacycloheptadecin-13(8H)-one, 21-chloro-20-fluoro-9,10,11,12,13a,14,15,16,18,23-decahydro-24-methoxy-4,6-ethanediylidenepyrimido[4,5-b][6,1,16]benzoxadiazacyclononadecin-16(17H)-one, 21-chloro-8,9,10,13,14,15,18,23-octahydro-24-methoxy-4,6-ethanediylidenepyrimido[4,5-b][6,1,16]benzoxadiazacyclononadecin-16(17H)-

-24-

one, 21-chloro-8,9,10,13,14,15,18,23-octahydro-24-methoxy-4,6-ethanediylidenepyrimido[4,5-b]pyrrolo[2,1-k][6,1,9,12]benzoxatriazacyclopentadecine-11(8H)-thione, 19-chloro-18-fluoro-9,10,11a,12,13,14,16,21-octahydro-22-methoxy-4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine, 19-chloro-8,9,10,11,12,13,14,15,16,21-decahydro-22-methoxy-15-methyl-4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine, 17-chloro-8,9,10,11,12,13,14,19-octahydro-20-methoxy-13-methyl-4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8H,13H)-dione, 17-chloro-10,11,14,19-tetrahydro-20-methoxy-11,11-dimethyl-9,11-ethanediylidenepyrimido[4,5-b]pyrrolo[1,2-i][6,1,9,12]benzoxatriazacyclopentadecine-14,19(5H,13H)-dione, 3-chloro-16,17,18,18a,20,21-hexahydro-17-hydroxy-22-methoxy-4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,15]benzoxadiazacyclooctadecine-15(16H)-one, 20-chloro-9,12,13,14,17,22-hexahydro-23-methoxy-4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacyclooctadecine, 20-chloro-9,10,11,12,13,14,15,16,17,22-decahydro-23-methoxy-16-methyl-4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-11,14-dione, 18-chloro-9,10,12,13,15,20-hexahydro-21-methoxy-10-(2-methoxyethyl)-4,6-ethanediylidenepyrimido[4,5-b][6,1,12,16]benzoxatriazacyclononadecine-12,16(13H,17H)-dione, 21-chloro-8,9,10,11,14,15,18,23-octahydro-24-methoxy-4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,12,17]benzoxatriazacycloeicosine-12,17(18H)-dione, 22-chloro-9,10,11,13,14,15,16,19,24-nonahydro-25-methoxy-4,6-ethanediylidene-8H-pyrimido[4,5-b]pyrrolo[1,2-i][6,1,12,15]benzoxatriazacyclooctadecine-12,17(18H)-dione, 22-chloro-9,10,11,14,15,16,16a,19,24-nonahydro-25-methoxy-4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-9,12-dione, 18-chloro-10,11,13,14,15,20-hexahydro-21-methoxy-11,11-dimethyl-4,6-ethanediylidene-8H-pyrimido[4,5-b]pyrrolo[1,2-i][6,1,12,15]benzoxatriazacyclooctadecine-12,17(18H)-dione, 22-chloro-9,10,11,14,15,16,16a,19,24-nonahydro-15-hydroxy-25-methoxy-4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacyclooctadecine-

-25-

- 12,15(16H)-dione, 20-chloro-9,10,11,13,14,17,22-heptahydro-23-methoxy-13-methyl-
- 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacyclooctadecine-12,15(16H)-dione, 20-chloro-9,10,11,13,14,17,22-heptahydro-23-methoxy-14-(2-methylpropyl)-
- 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacyclooctadecine-12,15(16H)-dione, 20-chloro-9,10,11,13,14,17,22-heptahydro-23-methoxy-14,14-dimethyl-
- 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacyclooctadecine-12,15(16H)-dione, 20-chloro-9,10,11,13,14,17,22-heptahydro-23-methoxy-14-(phenylmethyl)-
- 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacyclooctadecine-12,15(16H)-dione, 20-chloro-9,10,11,13,14,17,22-heptahydro-23-methoxy-14-methyl-
- 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-9,12-dione, 18-chloro-10,11,13,14,15,20-hexahydro-21-methoxy-11-methyl-
- 1,21-ethanediylidene-5H-pyrimido[4,5-b]pyrrolo[1,2-i][6,1,9,12]benzoxatriazacyclohexadecine-13,18(19H)-dione, 7-chloro-10,11,12,13a,14,15,16-heptahydro-23-methoxy-
- 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-9,12-dione, 18-chloro-10,11,13,14,15,20-hexahydro-21-methoxy-10-methyl-
- 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-9,12-dione, 18-chloro-10,11,13,14,15,20-hexahydro-21-methoxy-11-(2-methylpropyl)-
- 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-9,12-dione, 18-chloro-10,11,13,14,15,20-hexahydro-11-(1-hydroxyethyl)-21-methoxy-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine-11,14(8H,15H)-dione, 19-chloro-9,10,12,13,16,21-hexahydro-22-methoxy-13-(2-methylpropyl)-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine-11,14(8H,15H)-dione, 19-chloro-9,10,12,13,16,21-hexahydro-22-methoxy-13,13-dimethyl-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine-11,14(8H,15H)-dione, 19-chloro-9,10,12,13,16,21-hexahydro-22-methoxy-13-(phenylmethyl)-
- 4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine-

-26-

11,14(8H,15H)-dione, 19-chloro-9,10,12,13,16,21-hexahydro-13-(1-hydroxyethyl)-22-methoxy-4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacyclooctadecine-12,15(16H)-dione, 20-chloro-9,10,11,13,14,17,22-heptahydro-14-(1-hydroxyethyl)-23-methoxy-4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-11,14-dione, 18-chloro-9,10,12,13,15,20-hexahydro-21-methoxy-10-[2-(4-morpholinyl)ethyl]-carbamic acid, (20-chloro-9,10,13,14,15,16,17,22-octahydro-23-methoxy-15-oxo-4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,15]benzoxadiazacyclooctadecin-14-yl)-, 1,1-dimethylethyl ester

carbamic acid, (20-chloro-9,10,13,14,15,16,17,22-octahydro-23-methoxy-15-oxo-4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,15]benzoxadiazacyclooctadecin-14-yl)-, 1,1-dimethylethyl ester

4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-9,12-dione, 18-chloro-10,11,13,14,15,20-hexahydro-11-(hydroxymethyl)-21-methoxy-1,21-ethanediylidene-5H-pyrimido[4,5-b]pyrrolo[1,2-i][6,1,9,12]benzoxatriazacyclohexadecine-13,18(19H)-dione, 7-chloro-10,11,12,13a,14,15,16-heptahydro-15-hydroxy-23-methoxy-carbamic acid, (20-chloro-9,10,11,12,13,14,15,16,17,22-decahydro-23-methoxy-15-oxo-4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,15]benzoxadiazacyclooctadecin-14-yl)-, 1,1-dimethylethyl ester

4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,15]benzoxadiazacyclooctadecin-15(16H)-one, 14-amino-20-chloro-9,10,11,12,13,14,17,22-octahydro-23-methoxy-carbamic acid, (18-chloro-11,12,13,14,15,20-hexahydro-21-methoxy-13-oxo-4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,13]benzoxadiazacyclohexadecin-12-yl)-, 1,1-dimethylethyl ester

4,6-ethanediylidenepyrimido[4,5-b]pyrrolo[2,1-k][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 19-chloro-9,10,11a,12,13,14,16,21-octahydro-13-hydroxy-22-methoxy-4,6-ethanediylidene-13,16-ethano-8H-pyrimido[4,5-b][6,1,9,12,15]benzoxatetraazacyclooctadecin-11(12H)-one, 20-chloro-9,10,14,15,17,22-hexahydro-25-methoxy-8H-4,6-ethanediylidene-12,15-ethanopyrimido[4,5-b][6,1,9,14]benzoxatriazacycloheptadecin-11(12H)-one, 19-chloro-

-27-

9,10,13,14,16,21-hexahydro-24-methoxy-
4,6-ethanediylidene-12,16-methano-6H-pyrimido[4,5-
b][6,1,9,15]benzoxatriazacyclooctadecin-11(8H)-one, 20-chloro-
9,10,12,13,14,15,17,22-octahydro-24-methoxy-
4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-
one, 17-chloro-9,10,12,13,14,19-hexahydro-20-methoxy-12,13-dimethyl-
4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-
one, 17-chloro-13-ethyl-9,10,12,13,14,19-hexahydro-20-methoxy-
4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-
one, 17-chloro-9,10,12,13,14,19-hexahydro-12-(hydroxymethyl)-20-methoxy-
4,6-ethanediylidene-12H-pyrimido[4,5-b]pyrrolo[2,1-
l][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 20-chloro-
8,9,10,11,12a,13,14,15,17,22-decahydro-14-hydroxy-23-methoxy-
4,6-ethanediylidene-14,17-ethanopyrimido[4,5-
b][6,1,10,13,16]benzoxatetraazacyclononadecin-12(13H)-one, 21-chloro-
8,9,10,11,15,16,18,23-octahydro-26-methoxy-
4,6-ethanediylidene-13,16-ethano-6H-pyrimido[4,5-
b][6,1,10,15]benzoxatriazacyclooctadecin-12(13H)-one, 20-chloro-
8,9,10,11,14,15,17,22-octahydro-25-methoxy-
12H-4,6-ethanediylidene-13,17-methanopyrimido[4,5-
b][6,1,10,16]benzoxatriazacyclononadecin-12-one, 21-chloro-
8,9,10,11,13,14,15,16,18,23-decahydro-25-methoxy-
4,6-ethanediylidene-12H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclohexadecin-
12-one, 18-chloro-8,9,10,11,13,14,15,20-octahydro-21-methoxy-13,14-
dimethyl-
4,6-ethanediylidene-12H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclohexadecin-
12-one, 18-chloro-14-ethyl-8,9,10,11,13,14,15,20-octahydro-21-methoxy-
4,6-ethanediylidene-12H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclohexadecin-
12-one, 18-chloro-8,9,10,11,13,14,15,20-octahydro-13-(hydroxymethyl)-21-
methoxy-
4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecin-
13(8H)-one, 19-chloro-15-ethyl-9,10,11,12,14,15,16,21-octahydro-22-methoxy-
4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecin-
13(8H)-one, 19-chloro-9,10,11,12,14,15,16,21-octahydro-22-methoxy-14,15-
dimethyl-
4,6-ethanediylidenepyrimido[4,5-b][6,1,16]benzoxadiazacyclononadecin-16(17H)-
one, 21-chloro-8,9,10,11,12,13,14,15,18,23-decahydro-24-methoxy-

-28-

4,6-ethanediylidene-8*H*-pyrimido[4,5-*b*][6,1,15]benzoxadiazacyclooctadecin-15(16*H*)-one, 20-chloro-14-(dimethylamino)-9,10,11,12,13,14,17,22-octahydro-23-methoxy-

In a particular embodiment of the present invention the compounds of formula (I) are selected from the group consisting of;

- 4,6-ethanediylidenepyrimido[4,5-*b*][6,1,12]benzoxadiazacyclopentadecine, 17-bromo-8,9,10,11,12,13,14,19-octahydro-20-methoxy-13-methyl,
- 4,6-ethanediylidenepyrimido[4,5-*b*][6,1,12]benzoxadiazacyclopentadecine, 17-bromo-8,9,10,11,12,13,14,19-octahydro-20-methoxy-,
- 4,6-ethanediylidenepyrimido[4,5-*b*][6,1,10,13]benzoxatriazacycloheptadecine-12,15-dione, 19-chloro-8,9,10,11,13,14,16,21-octahydro-22-methoxy-13-(2-methylpropyl)-,
- 4,6-ethanediylidenepyrimido[4,5-*b*][6,1,10,13]benzoxatriazacycloheptadecine-12,15-dione, 19-chloro-8,9,10,11,13,14,16,21-octahydro-22-methoxy-,
- 4,6-ethanediylidenepyrimido[4,5-*b*]pyrrolo[2,1-*k*][6,1,9,12]benzoxatriazacyclopentadecin-11(8*H*)-one, 19-chloro-18-fluoro-9,10,11a,12,13,14,16,21-octahydro-22-methoxy-,
- 4,6-ethanediylidene-8*H*-pyrimido[4,5-*b*][6,1,10,13]benzoxatriazacyclohexadecine, 18-chloro-9,10,11,12,13,14,15,20-octahydro-21-methoxy-14-methyl-,
- 4,6-ethanediylidenepyrimido[4,5-*b*][6,1,11,14]benzoxatriazacycloheptadecine, 19-chloro-8,9,10,11,12,13,14,15,16,21-decahydro-22-methoxy-15-methyl-,
- 4,6-ethanediylidenepyrimido[4,5-*b*][6,1,9,12]benzoxatriazacyclopentadecine, 17-chloro-8,9,10,11,12,13,14,19-octahydro-20-methoxy-13-methyl-,
- 12*H*-4,6-ethanediylidene-13,17-methanopyrimido[4,5-*b*][6,1,10,16]benzoxatriazacyclononadecin-12-one, 21-chloro-8,9,10,11,13,14,15,16,18,23-decahydro-25-methoxy-,
- 4,6-ethanediylidene-12*H*-pyrimido[4,5-*b*][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 18-chloro-8,9,10,11,13,14,15,20-octahydro-21-methoxy-13,14-dimethyl-,
- 4,6-ethanediylidenepyrimido[4,5-*b*][6,1,11,14]benzoxatriazacycloheptadecin-13(8*H*)-one, 19-chloro-15-ethyl-9,10,11,12,14,15,16,21-octahydro-22-methoxy-, or
- 4,6-ethanediylidenepyrimido[4,5-*b*][6,1,11,14]benzoxatriazacycloheptadecin-13(8*H*)-one, 19-chloro-9,10,11,12,14,15,16,21-octahydro-22-methoxy-14,15-dimethyl-,

Other special group of compounds are:

- those compounds of formula (I) wherein -X¹- represents -O-;

-29-

- those compounds of formula (I) wherein $-X^1$ represents $-NR^{11}$ -, in particular $-NH$ -;
- those compounds of formula (I) wherein R^1 is fluoro, chloro or bromo;
- those compounds of formula (I) wherein R^2 is fluoro, chloro or bromo;
- 5 - those compounds of formula (I) wherein R^2 is Het^1 , in particular thiazolyl optionally substituted with methyl;
- those compounds of formula (I) wherein R^2 is C_{2-6} alkynyl-, in particular ethynyl;
- those compounds of formula (I) wherein R^2 is Ar^5 , in particular phenyl optionally substituted with cyano;
- 10 - those compounds of formula (I) wherein R^4 represents methoxy and wherein said methoxy is at position 7 of the structure of formula (I).
- those compounds of formula (I) wherein R^4 represents C_{1-4} alkyloxy substituted with one substituent selected from C_{1-4} alkyloxy- or Het^2 -, in particular propyloxy substituted with morpholinyl;
- 15 - those compounds of formula (I) wherein R^{12} is hydrogen or C_{1-4} alkyl-, in particular methyl or wherein R^{12} is C_{1-4} alkyl-oxy-carbonyl-, in particular t-butyl-oxy-carbonyl-
- those compounds of formula (I) wherein Het^2 represent morpholinyl optionally substituted with C_{1-4} alkyl, preferably morpholinyl attached through the nitrogen
- 20 atom to the remainder of the compounds of formula (I);
- those compounds of formula (I) with Het^3 represent morpholinyl optionally substituted with C_{1-4} alkyl, preferably morpholinyl attached through the nitrogen atom to the remainder of the compounds of formula (I);
- those compounds of formula (I) wherein Het^{12} represent morpholinyl optionally
- 25 substituted with C_{1-4} alkyl, preferably morpholinyl attached through the nitrogen atom to the remainder of the compounds of formula (I).


In a further embodiment of the present invention the X^2 substituent is at position 2', the R^1 substituent represents hydrogen or halo and is at position 4', the R^2 substituent

30 represents halo and is at position 5', the R^3 substituent is at position 2 and the R^4 substituent at position 7 of the structure of formula (I). Alternatively, the X^2 substituent is at position 3', the R^1 substituent represents hydrogen or halo and is at position 4', the R^2 substituent represents halo and is at position 5', the R^3 substituent is at position 2 and the R^4 substituent at position 7 of the structure of formula (I).

-30-

The compounds of this invention can be prepared by any of several standard synthetic processes commonly used by those skilled in the art of organic chemistry and described for instance in the following references; "Heterocyclic Compounds" – Vol.24 (part4) p 261-304 Fused pyrimidines, Wiley – Interscience ; Chem. Pharm.

5 Bull., Vol 41(2) 362-368 (1993); J.Chem.Soc., Perkin Trans. 1, 2001, 130-137.



Y_1 and Y_2 each independently represent a C_{1-5} alkyl, C_{1-6} alkyl, $CO-C_{1-6}$ alkyl, $CO-C_{1-5}$ alkyl, $Het^{22}-CH_2-CO$, $CO-CR^{16}R^{17}-NH$, $CR^{18}R^{19}-CO$, $CH_2-CO-NH-C_{1-3}$ alkyl-, $-C_{1-2}$ alkyl- $NR^{21}-CH_2-CO$ - or $CO-C_{1-3}$ alkyl- NH -


X_3 and X_4 represent optionally protected functional groups, such as for example a primary or secondary amine, hydroxy, hydrogen or halo (Cl, Br or I), which upon reaction produce together with the Y_1 respectively Y_2 substituent to which they are attached, the divalent Y radical as defined for formula (I)

As further exemplified in the experimental part of the description, a particular group of compounds are those compounds of formula (I) where $-X^1-$ represents $-O-$,

10 hereinafter referred to as compounds of formula (I') which are generally prepared using the following synthesis scheme. The compounds of this invention may be prepared starting from the known 6-acetoxy-4-chloro-7-methoxy quinazoline (II') or from 6-acetoxy-7-benzyloxy-4-chloroquinazoline (II''), which can be prepared from commercially available veratric acid and 4-hydroxy-3-methoxy benzoic acid, respectively.

Coupling of the latter compounds with suitable substituted anilines (III') under standard conditions, for example stirred in propanol at an elevated temperature ranging from 40-100°C during 3-12 h, anilines which in their turn can be prepared according to reaction schemes 4-8, furnish the intermediate compounds (IV', IV'') (Scheme 1).

-31-
Scheme 1




V = hydrogen or a protective group such as for example, methylcarbonyl, t-butyl, methyl, ethyl, benzyl or trialkylsilyl groups

R²³ represents Ar³, Ar⁴-C₁₋₄alkyl, C₁₋₄alkyl, C₂₋₆alkenyl optionally substituted with Het¹² or R²³ represents C₁₋₄alkyl substituted with one or where possible two or more substituents selected from C₁₋₄alkyloxy, hydroxy, halo, Het², NR⁷R⁸, NR⁹R¹⁰-carbonyl or Het³-carbonyl, wherein X², Ar³, Ar⁴, Het¹², Het², R¹, R², R⁷, R⁸, R⁹, R¹⁰ and Het³ are defined as for the compounds of formula (I)

Deprotection of the intermediates of formula (IV' - IV^a) as described in *Protective Groups in Organic Synthesis* by T.W. Greene and P.G.M. Wuts, 3rd edition, 1998


- 5 followed by ring closure under Mitsunobu conditions give the target compounds (I' - I^a). (Scheme 2 - wherein V and R¹⁶ are defined as hereinbefore)

Scheme 2



-32-

Alternatively, those compounds of formula (I^b) wherein Y represents -C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-, -C₁₋₅alkyl-NR¹⁴-CO-C₁₋₅alkyl-, -C₁₋₅alkyl-CO-NR¹⁵-C₁₋₅alkyl-, -C₁₋₆alkyl-NH-, -C₁₋₆alkyl-CO-, -C₁₋₃alkyl-NH-CO-Het²⁰-,
 -Het²²-CH₂-CO-NH-C₁₋₃alkyl-, -C₁₋₂alkyl-NH-CO-CR¹⁶R¹⁷-NH-, -C₁₋₂alkyl-CO-NH-CR¹⁸R¹⁹-CO-, -C₁₋₂alkyl-CO-NR²⁰-C₁₋₃alkyl-CO-,
 -C₁₋₂alkyl-NR²¹-CH₂-CO-NH-C₁₋₃alkyl- or -NR²²-CO-C₁₋₃alkyl-NH- are prepared using the following synthesis scheme. The intermediates of formula (IV^b) are obtained as described hereinbefore. Deprotection and subsequent formation of the corresponding ether using the appropriate aminated alcohol under standard conditions provides the intermediates of formula (XXVIII). Deprotection followed by ring closure provides the target compounds of formula (I^b).




V = hydrogen or a protective group such as for example, methylcarbonyl, t-butyl, methyl, ethyl, benzyl or trialkylsilyl groups or in case of solid phase chemistry the resin to which the remainder of the molecule is attached
 R²³ represents Ar³, Ar⁴-C₁₋₄alkyl, C₁₋₄alkyl, C₂₋₆alkenyl optionally substituted with Het¹² or R²³ represents C₁₋₄alkyl substituted with one or where possible two or more substituents selected from C₁₋₄alkoxy, hydroxy, halo, Het², NR⁷R⁸, NR⁹R¹⁰-carbonyl or Het³-carbonyl, wherein Ar³, Ar⁴, Het¹², Het², R⁷, R⁸, R⁹, R¹⁰ and Het³ are defined as for the compounds of formula (I)
 Y₁ and Y₂ each independently represent a C₁₋₆alkyl, C₁₋₆alkyl, CO-C₁₋₆alkyl, CO-C₁₋₆alkyl, Het²²-CH₂-CO, CO-CR¹⁶R¹⁷-NH-, CR¹⁸R¹⁹-CO-, CH₂-CO-NH-C₁₋₃alkyl-, -C₁₋₂alkyl-NR²¹-CH₂-CO- or CO-C₁₋₃alkyl-NH-

-33-

More specific examples for the synthesis of compounds of formula (I^b) are provided in reaction schemes 9-12.

- For those compounds where X² represents -O-, the suitable substituted anilines of formula (III^a) are generally prepared from the commercially available nitro-phenols (X) and the α , ω -protected halogenated alcohols (XI) under alkaline conditions in a reaction inert solvent, for example, using dimethylacetamide (DMA) in the presence of K₂CO₃. The resulting nitro-phenyl derivative (XII) is subsequently reduced according to standard conditions, for example, using iron/acetic acid, to yield the substituted anilines of formula (III^a) (Scheme 4).

Scheme 4



X represents a halogen such as for example, Cl, Br, and I
V represents hydrogen or a protective group such as for example methylcarbonyl

- For those compounds where X² represents -NR¹²-C₁₋₂alkyl-, the suitable substituted anilines of formula (III^b) are generally prepared from the commercially available 2-nitro-benzaldehydes (XIII) and the amine substituted alcohols (XIV) by reductive amination under standard conditions, for example using NaBH₄ and titanium(iv)isopropoxide as reducing agents in ethanol as solvent, yielding in a first step the nitro-benzylamines of formula (XV).


Next the primary free alcohol is protected using art known procedures, for example, using an esterification reaction with acetic anhydride in the presence of pyridine.

-34-

The thus obtained intermediate of formula (XVI) is subsequently reduced according to standard conditions, for example, using hydrogenolysis (H_2 , Pt/C, thiophene, MeOH) or tin(II)chloride ($SnCl_2 \cdot H_2O$, EtOH) to yield the substituted anilines of formula (III^b) (Scheme 5).

5

Scheme 5




V represents a protective group such as for example methylcarbonyl

Using the aforementioned method in two alternative forms, 4-chloro-2-nitrobenzaldehyde has been converted in suitable substituted anilines of formula (III^b). In a first method (Scheme 5a) the suitable substituted anilines of formula (III^b) were obtained by reductive amination of 4-chloro-2-nitrobenzaldehyde with primary amino acids.

A methanolic solution of 10 mmol aldehyde 1, 20 mmol amino acid 2, 19 mmol KF, 1 mL 4% thiophene (inDIPE) and 1 g Pt/C (slurry in THF or MeOH) under 1 atm of hydrogen is stirred at 50 °C (scheme 5a). The reaction mixture is filtered after consumption of 4 equivalents of hydrogen (typically after 48 h) and 3 equivalents of Boc anhydride are added. Next, the solution is stirred for 1-3 h at room temperature (LCMS monitoring), then an excess of 6 N ammonia in MeOH is added, and stirring is continued for 1 h to work-up excess Boc anhydride. Finally, the solution is evaporated to dryness (sublimation of tert-butyl carbamate is observed) and the resulting Boc-protected N-benzyl amino acid 4 is purified by HPLC. When R' is not equal to hydrogen, Boc-protected aniline 5 is observed as the major product. In this case method B can be employed to obtain the aniline of type 4 (vide infra).

25



Scheme 5a. a) 1 atm H_2 , Pt/C, KF, thiophene, MeOH, 50 °C, 1-2 days b) Boc_2O , MeOH, RT, 1-3 h.

5

A second method to obtain the suitable substituted anilines of formula (III)^b has been the reductive amination of primary and secondary amino acids, amino acid hydrochlorides, N-methylallylamine and methylaminoacetaldehyde dimethyl acetal with 4-chloro-2-nitrobenzaldehyde 1 and 4-chloro-3-fluoro-2-nitrobenzaldehyde 6 (Scheme 5b). Overall yields vary between 13 and 100%.

To a solution (suspension) of 5 mmol amine 7, 5 mmol aldehyde 1 or 6 in 30 mL dichloromethane is added 5 mmol titanium(IV)tert-butoxide and 5 mmol DIPEA (when 7 is hydrochloride).¹ After stirring for 15 min, 12 mmol sodium triacetoxyborohydride is added and stirring is continued for 1-5 h (LCMS monitoring). Next, the reaction is worked up with 10-20 mL of a saturated NaHCO_3 -solution and stirring is continued until bubbling stops. The resulting emulsion is filtered over a P3 sintered glass filter and washed with dichloromethane. The organic layer is separated and the aqueous phase extracted with dichloromethane. Drying of the combined organic layers with magnesium sulfate (or potassium carbonate), followed by filtration and evaporation of the solvent yields the crude N-benzylamine 8, which is usually pure enough to use for the next reaction step.


When R is hydrogen, the secondary amine can be protected with a Boc or Cbz group by respectively adding three equivalents of Boc anhydride or benzyl chloroformate and three eq of DIPEA to a methylene chloride solution of the amine and stirring for

¹ With Ti(IV)isopropoxide, transesterification was observed once when 7 was an amino acid tBu ester. The reduction was unexpectedly exothermic in this case, and the heat may have caused this side reaction.

-36-

16-24 h at room temperature. When R' is bulky, Boc protection is generally slow and requires prolonged refluxing in methylene chloride. Next, excess protecting agent is worked up by adding 6 N ammonia in methanol and stirring for 1 h at room temperature. After evaporation of the solvent, the product is purified by RP HPLC.

5



Scheme 5b. a) NaBH(OAc)₃, Ti(OtBu)₄, (DIPEA when 7 is hydrochloride), RT, 1-2 h
b) Boc₂O or CbzCl, DIPEA, CH₂Cl₂, RT to reflux, 16-24 h. c) H₂, Pt/C, thiophene, MeOH (or EtOAc or THF), 24-48 h d) SnCl₂.H₂O, EtOH, 50 °C, 1.5 h.

10

The thus obtained benzylamine 8 is subsequently reduced either by hydrogenolysis or in case R' contains a double bond by reduction with tin(II)chloride

15 Nitro reduction by hydrogenolysis

The benzylamine 8 is dissolved in methanol (or ethyl acetate or THF) and, upon addition of 1 g Pt/C (slurry in EtOAc) and thiophene (1 mL 4% in DIPE), stirred under 1 atm hydrogen at 50 °C (scheme 3, step a). After consumption of three equivalents of hydrogen, the mixture is filtered over dicalite. Removal of the solvent yields the crude aniline 9 which, depending on the nature and purity, can be crystallized from heptane, purified by HPLC or used as a crude in the next reaction step.

20

Nitro reduction with tin(II)chloride

This method was used when R' contains a double bond and hence cannot be reduced hydrogenolytically.

25

-37-


- To an ethanolic solution of crude nitro compound 8 is added 5 eq of tin(II)chloride dihydrate (scheme 3, step b). This mixture is stirred for 1.5 h at 50 °C. Next, the solution is cooled to RT, and saturated sodium bicarbonate and methylene chloride are added (bubbling). The resulting emulsion is filtered over a P3 sintered glass filter.
- 5 Separation of the organic layer, followed by drying on anhydrous potassium carbonate, filtering and removal of the solvent yields the crude aniline 9, which is usually pure enough to be used in the next reaction step.

- For those compounds where X^2 represents $-O-N=CH-$, the suitable substituted anilines of formula (III^c) are generally prepared according to reaction scheme 6.

In a first step the known 2-nitro-benzaldehydes (XIII) are converted into the corresponding oxime (XVII) using, for example, the art known condensation reaction with hydroxylamine.

- 15 Next said oxime of formula XVII is allowed to react with an halogenated alkylacetate under alkaline conditions, for example using K_2CO_3 in DMSO, followed by reducing the nitro group, for example, with hydrogenolysis (H_2 , Pt/C, thiophene, MeOH) or tin(II)chloride ($SnCl_2 \cdot H_2O$, EtOH), to provide the suitable substituted aniline of formula (III^c).

20



X represents a halogen such as for example Cl, Br, or I

- For those compounds where X^2 represents a direct bond and Y represents C₁₋₆alkyl-NH-CO-, the suitable substituted anilines of formula (III^d) are generally prepared according to reaction scheme 7.
- 25


In a first step the known 2-nitro-benzoic acids (XX) are amidated to the intermediates of formula (XXII) under art known conditions, for example, using a hydroxylated

-38-

amine of formula (XXI) that is added dropwise to a mixture of (XX) in CH_2Cl_2 in the presence of 1,1'-carbonylbis-1H-imidazole.

Next the primary free alcohol is protected using art known procedures, for example, using an esterification reaction with acetic anhydride in the presence of pyridine.

- 5 The thus obtained intermediate of formula (XXIII) is subsequently reduced according to standard conditions, for example, using hydrogenolysis (H_2 , Pt/C, thiophene, MeOH) or tin(II)chloride ($\text{SnCl}_2 \cdot \text{H}_2\text{O}$, EtOH) to yield the substituted anilines of formula (III^d).



10

V represents a protective group such as for example methylcarbonyl


For those compounds where X² represents a direct bond the suitable substituted anilines of formula (III^e) are generally prepared according to reaction scheme 8.

- 15 In a first step the known 2-nitro-benzaldehydes (XIII) are alkenated to the intermediates of formula (XXV) under art known conditions, for example, using the Wittig Reaction with the appropriate phosphonium salt of formula (XXIV).

Following esterification of the free carboxylic acid under standard conditions for example, using ethanol under acidic conditions, the intermediate of formula (XXVI) are reduced to yield the desired substituted anilines of formula (III^e).


20

-39-
Scheme 8



Y₃ represent C₁₋₇alkyl


Those compounds of formula (I^b) wherein -X₁-Y-X₂- comprises an amine-amide linker, i.e. -X₁-Y-X₂- represents -O-C₁₋₅alkyl-NR¹⁴-CO-C₁₋₅alkyl-NR¹²-C₁₋₂alkyl-, -O-C₁₋₃alkyl-NH-CO-Het²⁰-C₁₋₂alkyl- or -C₁₋₆alkyl-NH-CO-CH₂-Het²²-, -O-C₁₋₂alkyl-NH-CO-CR¹⁶R¹⁷-NH-C₁₋₂alkyl- were either prepared according to reaction scheme 9 in case m is 1, 2 or 4, or by reaction scheme 10 in case m is 3.



10 Scheme 9a. a) iPrOH, 80 °C, 2-24 h b) 6 N NH₃/ MeOH, RT, 1 h c) Br(CH₂)_mCH₂NHBoc (m = 1, 2, 4), Cs₂CO₃, DMF, RT, overnight.

-40-

To a solution of N-benzyl amino acid **4** or **9** (See Scheme 5a and 5b) in isopropanol is added one equivalent of chloroquinazoline **10** (Scheme 9a). The resulting solution is stirred for 2-24 h at 80 °C to give **11** (LCMS monitoring). Next, the mixture is cooled to RT and a 6 N methanolic solution of ammonia is added. After stirring for one hour, the solution is evaporated to dryness. The crude phenol **12** is then redissolved in dry DMF and, upon addition of 5 equivalents of cesium carbonate, stirred for one h at RT. To the resulting phenolate is then added 1-1.2 eq (to prevent overalkylation) of Boc-aminoalkyl bromide, and the mixture is stirred overnight at RT, evaporated to dryness, redissolved in dichloromethane and filtered over dicalite to remove cesium salts. This yields the crude Boc-aminoalkylated phenols **13**.



Scheme 9b. a) 6 N HCl, dioxane, 60 °C, overnight (R = Me, Et), or TFA/CH₂Cl₂/TIS (90:8:2) (R = *t*Bu) b) HBTU (or PyBop), DIPEA, DMF, RT, 1 h c) 48 % HBr, RT, 1-2 h.

As shown in scheme 9b, the ester function is then hydrolyzed, and Boc group(s) removed, by stirring a dioxane solution of **13** overnight at 60 °C in the presence of 6 N HCl (R = Me, Et) or at RT in the presence of TFA/CH₂Cl₂/TIS (90:8:2) (R = *t*Bu). After evaporation to dryness, the resulting amino acid **14** is redissolved in dry DMF and, upon addition of 6 eq of DIPEA, added dropwise to a solution of 3 eq HBTU (or


-41-

PyBOP)² in dry DMF. Stirring this solution for 1 h at RT, followed by evaporation of the solvent yields the crude macrocycle **15a**, which is purified by RP HPLC. A pre-purification can be done by dissolving the residue in CH₂Cl₂ and washing it with saturated sodium bicarbonate in water, followed by drying on potassium carbonate and removal of the solvent. The yield for this sequence is 10-65% starting from the aniline **4** or **9**.

A Cbz group, if present, can be removed prior purification by dissolving the crude macrocycle **15a** (R^{'''} = Cbz) in 48 % aqueous HBr and stirring for 1-2 h at room temperature (RT). After concentration of the reaction mixture and quenching with solid potassium carbonate, the deprotected macrocycle **15b** is obtained by extraction with CH₂Cl₂ (with added methanol in case of solubility problems), and is similarly purified by RP HPLC. The yield for Cbz deprotection is quantitative by LCMS.

To a solution of N-benzyl amino acid **4** or **9** in isopropanol is added one equivalent of chloroquinazoline **10** (scheme 10a). The resulting solution is stirred for 2-24 h at 80 °C to give **11**. Next, the mixture is cooled to RT and a 6 N methanolic solution of ammonia is added. After stirring for one hour, the solution is evaporated to dryness. The crude phenol **12** is then redissolved in dry DMF and, upon addition of 5 equivalents of cesium carbonate, stirred for one hour at RT. To the resulting phenolate is then added 0.8 eq (to prevent overalkylation) of 4-bromobutyronitrile, and the mixture is stirred overnight at RT, evaporated to dryness, redissolved in dichloromethane and filtered over dicalite to remove cesium salts.


² PyBop generates tris(pyrrolidino)phosphin oxide, which is often difficult to separate from the macrocycle. HBTU generates tetramethylurea, which is more easy to remove.



Scheme 10a. a) iPrOH, 80 °C, 2-24 h b) 6 N NH_3/MeOH , RT, 1 h c) $\text{Br}(\text{CH}_2)_3\text{CN}$, Cs_2CO_3 , DMF, RT, overnight


Crude **16** is then thoroughly dried, redissolved in 6 N ammonia/MeOH (to prevent dimerisation) and, after addition of some thiophene solution in DIPE (to prevent dechlorination), treated with wet Raney nickel under 1 atm hydrogen pressure at 14 °C (scheme 10b). After consumption of 2 equivalents of hydrogen gas (typically after 16-24 h), the mixture is filtered over dicalite and concentrated to provide the crude amine **17**. The ester function is then hydrolyzed to **18** (and, if $R'' = \text{Boc}$, the Boc group removed) by stirring a dioxane solution of **17** overnight in the presence of 6 N HCl ($R = \text{Me, Et}$). After evaporation to dryness, the resulting amino acid **18** is redissolved in dry DMF and, upon addition of 6 eq of DIPEA, added dropwise to a solution of 3 eq HBTU (or PyBOP) in dry DMF. Stirring this solution for 1 h at RT, followed by evaporation of the solvent yields the crude macrocycle **19**, which is purified by RP HPLC. A pre-purification can be done by dissolving the residue in CH_2Cl_2 and washing it with saturated sodium bicarbonate in water. The yield for this sequence is 4-30% starting from the aniline **4** or **9**.

-43-



Scheme 10b. a) H_2 , RaNi , 6 N NH_3/MeOH , 14 °C, 16-24 h b) 6 N HCl , dioxane, 60 °C, overnight c)
 HBTU (or PyBop), DIPEA , DMF , RT, 1 h.

- 5 Those compounds of formula (I^b) wherein $-\text{X}_1-\text{Y}-\text{X}_2-$ comprises an amine-amine linker, i.e. $-\text{X}_1-\text{Y}-\text{X}_2-$ represents $-\text{O}-\text{C}_{1-5}\text{alkyl}-\text{NR}^{13}-\text{C}_{1-5}\text{alkyl}-\text{NR}^{12}-\text{C}_{1-2}\text{alkyl}-$ were generally prepared according to reaction scheme 11.



Scheme 11. a) *i*PrOH, 80 °C, 7-8 h b) NH₃/ MeOH, 1 h c) BrCH₂(CH₂)_nNHBoc ($n = 1, 2, 4$), * Cs₂CO₃, DMF, overnight. *Br(CH₂)₃CN used instead of $n = 3$ d) 6 N HCl, dioxane, 60 °C, 24 h e) NaBH(OAc)₃, CH₂Cl₂, RT, 1h.

5

To a solution of 5-chloro-2-([(2,2-dimethoxyethyl)(methyl)amino]methyl)aniline **20** or the corresponding dioxolane (prepared via reductive amination Scheme 5b) in isopropanol is added one equivalent of chloroquinazoline **10**. The resulting solution is stirred for 7-8 h at 80 °C to give **21**. Next, the mixture is cooled to RT and a 6 N methanolic solution of ammonia is added to remove the acetyl group. After stirring for one hour, the solution is evaporated to dryness. To the crude phenol **22** is then added 5 eq of cesium carbonate and, after stirring for 1 h, Boc-aminoalkyl bromide (1.0-1.2 eq) or 5-bromobutyronitrile (0.8 eq), and the mixture is subsequently stirred overnight at RT evaporated to dryness, redissolved in dichloromethane and filtered

10


-45-

over dicalite to remove cesium salts. This yields the protected aminoalkyl-substituted phenols **23**. The butyronitrile-substituted phenol is first hydrogenated to the corresponding amine under the abovementioned conditions.

- 5 Next the compounds are dissolved in 6 N HCl and dioxane, and the resulting mixture is stirred for about 24 h at 60 °C (LCMS monitoring, formation of imine **24** is observed) (scheme 11). After completion of the reaction, the mixture is poured carefully into an ice-cooled sodium bicarbonate solution or evaporated to dryness (slow, leads to decomposition). In the first case, the imine is extracted with methylene chloride and, after drying on potassium carbonate, immediately reduced to the corresponding amine by addition of sodium triacetoxyborohydride. In the latter case, the oily residue is redissolved in methylene chloride and excess sodium triacetoxyborohydride is added to yield the corresponding amine. The crude macrocycle **25** is obtained after addition of saturated sodium carbonate, extraction with dichloromethane followed by drying on potassium carbonate and removal of the solvent, and can be purified RP HPLC.

Those compounds of formula (I^b) wherein -X₁-Y-X₂- comprises an amide-amide linker, i.e. -X₁-Y-X₂- represents -O-C₁₋₄alkyl-CO-NH-CR¹⁸R¹⁹-CO-NR¹²-C₁₋₂alkyl-,
 20 -O-C₁₋₄alkyl-CO-NR²⁰-C₁₋₃alkyl-CO-NR¹²-C₁₋₂alkyl-, or
 -O-C₁₋₄alkyl-CO-Het²⁰-CO-NR¹²-C₁₋₂alkyl- were generally prepared according to reaction scheme 12.

Scheme 12



-46-

Reagents and conditions: a) i) PL-DCC Resin, HOBT, DCM/DMF, RT, 5h; ii) Polystyrene-methylisocyanate, (polystyrylmethyl)trimethylammonium bicarbonate, RT, 12 h; b) i) chloro quinazoline, iPrOH, 55°C, 3h; ii) 7N NH₃ in MeOH, rt, 2h; c) ClCH₂(CH₂)_mCO₂CH₃, K₂CO₃; d) conc. HCl, H₂O, dioxane, 60°C, 12h; e) PyBOP, DIPEA, DMF, RT, 3h.

In this procedure the aniline 26 is coupled with the appropriate amino protected amino acid, to form the amide of formula 27 using art known conditions, see for example A42 d) hereinbelow. Subsequent coupling with the chloroquinazoline under standard conditions, for example stirred in propanol at an elevated temperature ranging from 40-100°C during 3-12 h, provides the intermediates of formula 28. Alkylation with the appropriate haloacetate followed by deprotection and ring closure, i.e. amide formation using art known conditions, provides the compounds of formula 30. Deprotection of the intermediates of formula 28 can be done as described in *Protective Groups in Organic Synthesis* by T.W. Greene and P.G.M. Wuts, 3rd edition, 1998.


Alternatively, the compounds of formula (I) are prepared by ring closing metathesis of the corresponding olefins (Scheme 13). This reaction is particular useful for those compounds of formula (I) wherein Y represents -C₃₋₉alkenyl-, -C₃₋₉alkyl-, -C₃₋₇alkyl-CO-NH- optionally substituted with amino, mono-or di(C₁₋₄alkyl)amino or C₁₋₄alkyloxycarbonylamino-, or Y represents -C₃₋₇alkenyl-CO-NH- optionally substituted with amino, mono-or di(C₁₋₄alkyl)amino or C₁₋₄alkyloxycarbonylamino- hereinafter referred to as the compounds of formula (I^c).

In a first step the aniline of formula (III^c) is coupled to the 4-chloroquinazoline (II^b) under standard conditions, for example stirred in propanol at an elevated temperature ranging from 40-100°C during 3-12 h. Deprotection of the intermediate of formula (IV^c) as described in *Protective Groups in Organic Synthesis* by T.W. Greene and P.G.M. Wuts, 3rd edition, 1998, followed by alkylation with the appropriate alkyl bromides (XXXII) under art known conditions, such as for example stirring overnight at room temperature in the presence of Cs₂CO₃ in a reaction inert solvent such as for example N,N-dimethylformamide (DMF), provides the olefins of formula (XXXIII). Ring closing metathesis as described in *Advanced Organic Chemistry* by J. March, 3rd edition, 1985, p1036-1039 provides the compounds of formula (I^c) that can optionally be reduced using art known procedures, for example, stirring for 3-10 h at room temperature under H₂-atmosphere in the presence of Pt/C in tetrahydrofuran

-47-

(THF)/ methanol as solvent. The aniline of formula (III^e) is synthesized similar to the synthesis of the amide of formula 27 (Scheme 12 above) by acylation of the corresponding amine.

Scheme 13



R²³ represents Ar³, Ar⁴-C₁₋₄alkyl, C₁₋₄alkyl, C₂₋₆alkenyl optionally substituted with Het¹² or R²³ represents C₁₋₄alkyl substituted with one or where possible two or more substituents selected from C₁₋₄alkoxy, hydroxy, halo, Het², NR⁷R⁸, NR⁸R¹⁰-carbonyl or Het³-carbonyl, wherein Ar³, Ar⁴, Het¹², Het², R⁷, R⁸, R⁹, X², R¹, R², R³ and Het³ are defined as for the compounds of formula (I)
 Y₃ represent a C₁₋₅alkyl, CO-C₁₋₅alkyl or CO-CR¹⁶R¹⁷-NH- or C₁₋₅alkyl-CO- optionally substituted with amino, mono- or di(C₁₋₄alkyl)amino or C₁₋₄alkyloxycarbonylamino
 m represents 1, 2, 3 or 4

5

Where necessary or desired, any one or more of the following further steps in any order may be performed :

- 10 (i) removing any remaining protecting group(s);
- (ii) converting a compound of formula (I) or a protected form thereof into a further compound of formula (I) or a protected form thereof;
- (iii) converting a compound of formula (I) or a protected form thereof into a *N*-oxide, a salt, a quaternary amine or a solvate of a compound of formula (I) or a protected
- 15 form thereof;

-48-

(iv) converting a *N*-oxide, a salt, a quaternary amine or a solvate of a compound of formula (I) or a protected form thereof into a compound of formula (I) or a protected form thereof;

(v) converting a *N*-oxide, a salt, a quaternary amine or a solvate of a compound of formula (I) or a protected form thereof into another *N*-oxide, a pharmaceutically acceptable addition salt a quaternary amine or a solvate of a compound of formula (I) or a protected form thereof;

(vi) where the compound of formula (I) is obtained as a mixture of (R) and (S) enantiomers resolving the mixture to obtain the desired enantiomer.

Compounds of formula (I), *N*-oxides, addition salts, quaternary amines and stereochemical isomeric forms thereof can be converted into further compounds according to the invention using procedures known in the art.

It will be appreciated by those skilled in the art that in the processes described above the functional groups of intermediate compounds may need to be blocked by protecting groups.

Functional groups, which it is desirable to protect, include hydroxy, amino and carboxylic acid. Suitable protecting groups for hydroxy include trialkylsilyl groups (e.g. *tert*-butyldimethylsilyl, *tert*-butyldiphenylsilyl or trimethylsilyl), benzyl and tetrahydropyranyl. Suitable protecting groups for amino include *tert*-butyloxycarbonyl or benzyloxycarbonyl. Suitable protecting groups for carboxylic acid include C₍₁₋₆₎alkyl or benzyl esters.

The protection and deprotection of functional groups may take place before or after a reaction step.

Additionally, the N-atoms in compounds of formula (I) can be methylated by art-known methods using CH₃-I in a suitable solvent such as, for example 2-propanone, tetrahydrofuran or dimethylformamide.

The compounds of formula (I) can also be converted into each other following art-known procedures of functional group transformation of which some examples are mentioned hereinafter.

-49-


- The compounds of formula (I) may also be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with 3-phenyl-2-(phenylsulfonyl)oxaziridine or with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboperoxoic acid or halo substituted benzenecarboperoxoic acid, e.g. 3-chlorobenzenecarboperoxoic acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g. *t*-butyl hydroperoxide. Suitable solvents are, for example, water, lower alkanols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.
- Pure stereochemically isomeric forms of the compounds of formula (I) may be obtained by the application of art-known procedures. Diastereomers may be separated by physical methods such as selective crystallization and chromatographic techniques, e.g. counter-current distribution, liquid chromatography and the like.
- Some of the compounds of formula (I) and some of the intermediates in the present invention may contain an asymmetric carbon atom. Pure stereochemically isomeric forms of said compounds and said intermediates can be obtained by the application of art-known procedures. For example, diastereoisomers can be separated by physical methods such as selective crystallization or chromatographic techniques, e.g. counter current distribution, liquid chromatography and the like methods. Enantiomers can be obtained from racemic mixtures by first converting said racemic mixtures with suitable resolving agents such as, for example, chiral acids, to mixtures of diastereomeric salts or compounds; then physically separating said mixtures of diastereomeric salts or compounds by, for example, selective crystallization or chromatographic techniques, e.g. liquid chromatography and the like methods; and finally converting said separated diastereomeric salts or compounds into the corresponding enantiomers. Pure stereochemically isomeric forms may also be obtained from the pure stereochemically isomeric forms of the appropriate intermediates and starting materials, provided that the intervening reactions occur stereospecifically.

-50-

An alternative manner of separating the enantiomeric forms of the compounds of formula (I) and intermediates involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase.

- 5 Some of the intermediates and starting materials as used in the reaction procedures mentioned hereinabove are known compounds and may be commercially available or may be prepared according to art-known procedures. However, in the synthesis of the compounds of formula (I), the present invention further provides;

- 10 a) the intermediates of formula (III)



the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

- 15 V represents hydrogen or a protective group preferably selected from the group consisting of methylcarbonyl, t-butyl, methyl, ethyl, benzyl or trialkylsilyl;
Y represents -C₃₋₉alkyl-, -C₃₋₉alkenyl-, -C₃₋₇alkyl-CO-NH- optionally substituted with amino, mono- or di(C₁₋₄alkyl)amino or C₁₋₄alkyloxycarbonylamino-, -C₃₋₇alkenyl-CO-NH- optionally substituted with amino, mono- or di(C₁₋₄alkyl)amino or C₁₋₄alkyloxycarbonylamino-,
20 -C₁₋₅alkyl-oxy-C₁₋₅alkyl-, -C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-,
-C₁₋₅alkyl-NR¹⁴-CO-C₁₋₅alkyl-,
-C₁₋₅alkyl-CO-NR¹⁵-C₁₋₅alkyl-, -C₁₋₆alkyl-CO-NH-, -C₁₋₆alkyl-NH-CO-,
-C₁₋₃alkyl-NH-CS-Het²⁰-, -C₁₋₃alkyl-NH-CO-Het²⁰-, C₁₋₂alkyl-CO-Het²¹-CO-,
25 -Het²²-CH₂-CO-NH-C₁₋₃alkyl-, -CO-NH-C₁₋₆alkyl-, -NH-CO-C₁₋₆alkyl-,
-CO-C₁₋₇alkyl-, -C₁₋₇alkyl-CO-, -C₁₋₆alkyl-CO-C₁₋₆alkyl-,
-CO-Het²⁰-, -C₁₋₂alkyl-NH-CO-CR¹⁶R¹⁷-NH-, -C₁₋₂alkyl-CO-NH-CR¹⁸R¹⁹-CO-,
-C₁₋₂alkyl-CO-NR²⁰-C₁₋₃alkyl-CO-, -C₁₋₂alkyl-NR²¹-CH₂-CO-NH-C₁₋₃alkyl-, or
-NR²²-CO-C₁₋₃alkyl-NH-;
30 X² represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹²,
-NR¹²-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or -C₁₋₂alkyl-;
R¹ represents hydrogen, cyano, halo, hydroxy, formyl, C₁₋₆alkoxy-, C₁₋₆alkyl-,
halo-phenyl-carbonylamino-,
C₁₋₆alkoxy- substituted with halo,

-51-

- C₁₋₄alkyl substituted with one or where possible two or more substituents selected from hydroxy or halo;
- R² represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-, Het¹⁶-carbonyl-, C₁₋₄alkyloxy-carbonyl-, C₁₋₄alkylcarbonyl-, aminocarbonyl-, mono-or
 5 di(C₁₋₄alkyl)aminocarbonyl-, Het¹, formyl, C₁₋₄alkyl-, C₂₋₆alkynyl-, C₃₋₆cycloalkyl-, C₃₋₆cycloalkyloxy-, C₁₋₆alkoxy-, Ar⁵, Ar¹-oxy-, dihydroxyborane, C₁₋₆alkoxy- substituted with halo,
 C₁₋₄alkyl substituted with one or where possible two or more substituents selected from halo, hydroxy or NR⁵R⁶,
- 10 C₁₋₄alkylcarbonyl- wherein said C₁₋₄alkyl is optionally substituted with one or where possible two or more substituents selected from hydroxy or C₁₋₄alkyl-oxy-;
- R⁵ and R⁶ are each independently selected from hydrogen or C₁₋₄alkyl;
- R¹² represents hydrogen, C₁₋₄alkyl, C₁₋₄alkyl-oxy-carbonyl-, Het¹⁸-C₁₋₄alkyl-,
 15 phenyl-C₁₋₄alkyl-oxy-carbonyl-, Het¹⁷, C₂₋₄alkenylcarbonyl- optionally substituted with Het¹⁹-C₁₋₄alkylaminocarbonyl-, C₂₋₄alkenylsulfonyl-, C₁₋₄alkyloxyC₁₋₄alkyl- or R¹² represents phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;
- 20 R¹³ represents hydrogen, C₁₋₄alkyl, Het¹³, Het¹⁴-C₁₋₄alkyl- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;
- R¹⁴ and R¹⁵ are each independently selected from hydrogen, C₁₋₄alkyl, Het¹⁵-C₁₋₄alkyl- or C₁₋₄alkyloxyC₁₋₄alkyl-;
- 25 R¹⁶ and R¹⁷ each independently represents hydrogen or C₁₋₄alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl, aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;
- R¹⁸ and R¹⁹ each independently represents hydrogen or C₁₋₄alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl,
 30 aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;
- R²⁰ and R²² each independently represents hydrogen or C₁₋₄alkyl optionally substituted with hydroxy or C₁₋₄alkyloxy;
- R²¹ represents hydrogen, C₁₋₄alkyl, Het²³-C₁₋₄alkylcarbonyl- or
 R²¹ represents mono-or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-carbonyl- optionally
 35 substituted with hydroxy, pyrimidinyl, dimethylamine or C₁₋₄alkyloxy;
- Het¹ represents a heterocycle selected from piperidinyl, morpholinyl, piperazinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl,

-52-

- oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het¹ is optionally substituted with amino, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl-, phenyl, phenyl-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl- mono- or di(C₁₋₄alkyl)amino- or amino-carbonyl-;
- 5 Het¹³ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;
- 10 Het¹⁴ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;
- 15 Het¹⁵ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;
- 20 Het¹⁶ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl, 1,3,2-dioxaborolane or piperidinyl wherein said heterocycle is optionally substituted with one or more substituents selected from C₁₋₄alkyl; and
- 25 Het¹⁷ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;
- 30 Het¹⁸ and Het¹⁹ each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;
- 35 Het²⁰, Het²¹ and Het²² each independently represent a heterocycle selected from pyrrolidinyl, 2-pyrrolidinonyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from hydroxy, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl- or polyhydroxy-C₁₋₄alkyl-;
- Ar¹, Ar², Ar³, Ar⁴ and Ar⁵ each independently represent phenyl optionally substituted with cyano, C₁₋₄alkylsulfonyl-, C₁₋₄alkylsulfonylamino-, aminosulfonylamino-, hydroxy-C₁₋₄alkyl, aminosulfonyl-, hydroxy-, C₁₋₄alkyloxy- or C₁₋₄alkyl.

In one embodiment the intermediates of formula (III) consists of the intermediates of formula (III) wherein one or more of the following restrictions apply;

-53-

V represents hydrogen or a protective group preferably selected from the group consisting of methylcarbonyl, t-butyl, methyl, ethyl, benzyl or trialkylsilyl;

Y represents -C₃₋₉alkyl-, -C₃₋₉alkenyl-, -C₁₋₅alkyl-oxy-C₁₋₅alkyl-,
 -C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-, -C₁₋₅alkyl-NR¹⁴-CO-C₁₋₅alkyl-,
 -C₁₋₅alkyl-CO-NR¹⁵-C₁₋₅alkyl-, -C₁₋₆alkyl-CO-NH-, -C₁₋₆alkyl-NH-CO-,
 -C₁₋₇alkyl-CO-, C₁₋₆alkyl-CO-C₁₋₆alkyl;

X² represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹²,
 -NR¹²-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or C₁₋₂alkyl;

R¹ represents hydrogen, cyano, halo, hydroxy, formyl, C₁₋₆alkoxy-, C₁₋₆alkyl-,

C₁₋₆alkoxy- substituted with halo,

C₁₋₄alkyl substituted with one or where possible two or more substituents selected from hydroxy or halo; and

R² represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-, Het¹⁶-carbonyl-,

C₁₋₄alkyloxycarbonyl-, C₁₋₄alkylcarbonyl-, aminocarbonyl-, mono-or

di(C₁₋₄alkyl)aminocarbonyl-, Het¹, formyl, C₁₋₄alkyl-, C₂₋₆alkynyl-,

C₃₋₆cycloalkyl-, C₃₋₆cycloalkyloxy-, C₁₋₆alkoxy-, Ar⁵, Ar¹-oxy-, dihydroxyborane,
 C₁₋₆alkoxy- substituted with halo,

C₁₋₄alkyl substituted with one or where possible two or more substituents selected from halo, hydroxy or NR⁵R⁶,

C₁₋₄alkylcarbonyl- wherein said C₁₋₄alkyl is optionally substituted with one or where possible two or more substituents selected from hydroxy or C₁₋₄alkyl-oxy-;

R⁵ and R⁶ are each independently selected from hydrogen or C₁₋₄alkyl;

R¹² represents hydrogen, C₁₋₄alkyl, C₁₋₄alkyl-oxy-carbonyl-,

phenyl-C₁₋₄alkyl-oxy-carbonyl-, Het¹⁷, Het¹⁸-C₁₋₄alkyl-,

C₂₋₄alkenylcarbonyl- optionally substituted with Het¹⁹-C₁₋₄alkylaminocarbonyl-,

C₂₋₄alkenylsulfonyl-, C₁₋₄alkyloxyC₁₋₄alkyl- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;

R¹³ represents hydrogen, C₁₋₄alkyl, Het¹³, Het¹⁴-C₁₋₄alkyl- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;

R¹⁴ and R¹⁵ are each independently selected from hydrogen, C₁₋₄alkyl,

Het¹⁵-C₁₋₄alkyl- or C₁₋₄alkyloxyC₁₋₄alkyl-;

Het¹ represents a heterocycle selected from piperidinyl, morpholinyl, piperazinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het¹ is optionally substituted

-54-

with amino, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl-, phenyl, phenyl-C₁₋₄alkyl-,
C₁₋₄alkyl-oxy-C₁₋₄alkyl- mono- or di(C₁₋₄alkyl)amino- or amino-carbonyl-;

Het¹³ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally
substituted with one or where possible two or more substituents selected from
5 C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or
polyhydroxy-C₁₋₄alkyl-;

Het¹⁴ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or
piperidinyl optionally substituted with one or where possible two or more
substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-,
10 C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het¹⁵ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or
piperidinyl optionally substituted with one or where possible two or more
substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-,
C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

15 Het¹⁶ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl,
1,3,2-dioxaborolane or piperidinyl wherein said heterocycle is optionally
substituted with one or more substituents selected from C₁₋₄alkyl; and

Het¹⁷ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally
substituted with one or where possible two or more substituents selected from
20 C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or
polyhydroxy-C₁₋₄alkyl-;

Het¹⁸ and Het¹⁹ each independently represent a heterocycle selected from
morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with
one or where possible two or more substituents selected from C₁₋₄alkyl,
25 C₃₋₆cycloalkyl,

hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Ar¹, Ar², Ar³, Ar⁴ and Ar⁵ each independently represent phenyl optionally substituted
with cyano, C₁₋₄alkylsulfonyl-, C₁₋₄alkylsulfonylamino-, aminosulfonylamino-,
hydroxy-C₁₋₄alkyl, aminosulfonyl-, hydroxy-, C₁₋₄alkyloxy- or C₁₋₄alkyl.

30

In particular the intermediates of formula (III) wherein one or more of the following
restrictions apply;


i) V represents hydrogen, methyl or ethyl;

ii) Y represents -C₃₋₉alkyl-, -C₁₋₅alkyl-oxy-C₁₋₅alkyl-, -C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-,
35 -C₁₋₆alkyl-NH-CO-;

-55-

- iii) Y represents $-C_{1-5}\text{alkyl-oxy-}C_{1-5}\text{alkyl}$, $C_{1-2}\text{alkyl-CO-Het}^{21}\text{-CO-}$, $\text{-CO-}C_{1-7}\text{alkyl-}$,
or
 -CO-Het^{20} ;
- iv) X^2 represents a direct bond, O, $\text{-O-}C_{1-2}\text{alkyl-}$, NR^{12} , $\text{-NR}^{12}\text{-}C_{1-2}\text{alkyl-}$, $\text{-CH}_2\text{-}$,
5 -O-N=CH- or $C_{1-2}\text{alkyl}$;
- v) X^2 represents $\text{-NR}^{12}\text{-}C_{1-2}\text{alkyl-}$ or $C_{1-2}\text{alkyl}$;
- vi) R^1 represents hydrogen, cyano, halo or hydroxy, preferably halo;
- vii) R^2 represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-,
 $C_{1-4}\text{alkyloxycarbonyl-}$, $\text{Het}^{16}\text{-carbonyl-}$, $C_{1-4}\text{alkyl-}$, $C_{2-6}\text{alkynyl-}$, Ar^5 or Het^1 ;
10 preferably halo;
- viii) R^2 represents hydrogen, cyano, halo, hydroxy, $C_{2-6}\text{alkynyl-}$ or Het^1 ; in particular
 R^2 represents hydrogen, cyano, halo, hydroxy, or Ar^5 ;
- ix) R^{12} represents hydrogen, $C_{1-4}\text{alkyl}$, $C_{1-4}\text{alkyloxycarbonyl}$ or
phenyl- $C_{1-4}\text{alkyl-oxy-carbonyl-}$;
- x) R^{13} represents $\text{Het}^{14}\text{-}C_{1-4}\text{alkyl}$, in particular morpholinyl- $C_{1-4}\text{alkyl}$;
- xi) Het^1 represents thiazolyl optionally substituted with amino, $C_{1-4}\text{alkyl}$,
hydroxy- $C_{1-4}\text{alkyl-}$, phenyl, phenyl- $C_{1-4}\text{alkyl-}$, $C_{1-4}\text{alkyl-oxy-}C_{1-4}\text{alkyl-}$ mono- or
di($C_{1-4}\text{alkyl}$)amino- or amino-carbonyl-;
- xii) Het^{16} represents a heterocycle selected from piperidinyl or pyrrolidinyl;
- 20 xiii) Het^{20} represents piperidine, piperazine, pyrrolidinyl or 2-pyrrolidinonyl wherein
said Het^{20} is optionally substituted with hydroxy.

b) the intermediates of formula (XXX)



- 25 the pharmaceutically acceptable addition salts and the stereochemically isomeric
forms thereof, wherein
- Y1 and Y2 each independently represent a $C_{1-5}\text{alkyl}$, $C_{1-6}\text{alkyl}$, $\text{CO-}C_{1-6}\text{alkyl}$,
 $\text{CO-}C_{1-5}\text{alkyl}$, $\text{Het}^{22}\text{-CH}_2\text{-CO}$, $\text{CO-CR}^{16}\text{R}^{17}\text{-NH-}$, Het^{20} , $\text{CR}^{18}\text{R}^{19}\text{-CO-}$,
 $\text{CH}_2\text{-CO-NH-}C_{1-3}\text{alkyl-}$, $\text{-}C_{1-2}\text{alkyl-NR}^{21}\text{-CH}_2\text{-CO-}$ or $\text{CO-}C_{1-3}\text{alkyl-NH-}$;
- 30 X^1 represents a direct bond, O, $\text{-O-}C_{1-2}\text{alkyl-}$, CO, $\text{-CO-}C_{1-2}\text{alkyl-}$, NR^{11} ,
 $\text{-NR}^{11}\text{-}C_{1-2}\text{alkyl-}$, $\text{-CH}_2\text{-}$, -O-N=CH- or $\text{-}C_{1-2}\text{alkyl-}$;

-56-

X^2 represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹²,
-NR¹²-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or -C₁₋₂alkyl-;

R¹ represents hydrogen, cyano, halo, hydroxy, formyl, C₁₋₆alkoxy-, C₁₋₆alkyl-,
halo-phenyl-carbonylamino-,

5 C₁₋₆alkoxy- substituted with halo,

C₁₋₄alkyl substituted with one or where possible two or more substituents selected
from hydroxy or halo;

R² represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-, Het¹⁶-carbonyl-,
C₁₋₄alkyloxy-carbonyl-, C₁₋₄alkylcarbonyl-, aminocarbonyl-, mono-or

10 di(C₁₋₄alkyl)aminocarbonyl-, Het¹, formyl, C₁₋₄alkyl-, C₂₋₆alkynyl-,

C₃₋₆cycloalkyl-, C₃₋₆cycloalkyloxy-, C₁₋₆alkoxy-, Ar⁵, Ar¹-oxy-, dihydroxyborane,
C₁₋₆alkoxy- substituted with halo,

C₁₋₄alkyl substituted with one or where possible two or more substituents selected
from halo, hydroxy or NR⁵R⁶,

15 C₁₋₄alkylcarbonyl- wherein said C₁₋₄alkyl is optionally substituted with one or
where possible two or more substituents selected from hydroxy or
C₁₋₄alkyl-oxy-;

R³ represents hydrogen, C₁₋₄alkyl, or C₁₋₄alkyl substituted with one or more
substituents selected from halo, C₁₋₄alkyloxy-, amino-, mono-or

20 di(C₁₋₄alkyl)amino-,

C₁₋₄alkyl-sulfonyl- or phenyl;

R⁵ and R⁶ are each independently selected from hydrogen or C₁₋₄alkyl;

R⁷ and R⁸ are each independently selected from hydrogen, C₁₋₄alkyl, Het⁸,

aminosulfonyl-, mono- or di (C₁₋₄alkyl)-aminosulfonyl, hydroxy-C₁₋₄alkyl-,

25 C₁₋₄alkyl-oxy-C₁₋₄alkyl-, hydroxycarbonyl-C₁₋₄alkyl-, C₃₋₆cycloalkyl,

Het⁹-carbonyl-C₁₋₄alkyl-, Het¹⁰-carbonyl-, polyhydroxy-C₁₋₄alkyl-,

Het¹¹-C₁₋₄alkyl- or Ar²-C₁₋₄alkyl-;

R⁹ and R¹⁰ are each independently selected from hydrogen, C₁₋₄alkyl, C₃₋₆cycloalkyl,
Het⁴, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl- or polyhydroxy-C₁₋₄alkyl-;

30 R¹¹ represents hydrogen, C₁₋₄alkyl, Het⁵, Het⁶-C₁₋₄alkyl-, C₂₋₄alkenylcarbonyl-

optionally substituted with Het⁷-C₁₋₄alkylaminocarbonyl-, C₂₋₄alkenylsulfonyl-,

C₁₋₄alkyloxyC₁₋₄alkyl- or phenyl optionally substituted with one or where
possible two or more substituents selected from hydrogen, hydroxy, amino or
C₁₋₄alkyloxy-;

35 R¹² represents hydrogen, C₁₋₄alkyl, C₁₋₄alkyl-oxy-carbonyl-, Het¹⁷, Het¹⁸-C₁₋₄alkyl-,

C₂₋₄alkenylcarbonyl- optionally substituted with Het¹⁹-C₁₋₄alkylaminocarbonyl-,

C₂₋₄alkenylsulfonyl-, C₁₋₄alkyloxyC₁₋₄alkyl- or phenyl optionally substituted with

-57-

one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;

R¹⁶ and R¹⁷ each independently represents hydrogen or C₁₋₄alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl, aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;

R¹⁸ and R¹⁹ each independently represents hydrogen or C₁₋₄alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl, aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;

R²¹ represents hydrogen, C₁₋₄alkyl, Het²³-C₁₋₄alkylcarbonyl- or

R²¹ represents mono- or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-carbonyl- optionally substituted with hydroxy, pyrimidinyl, dimethylamine or C₁₋₄alkyloxy;

R²³ represents Ar³, Ar⁴-C₁₋₄alkyl, C₁₋₄alkyl, C₂₋₆alkenyl optionally substituted with Het¹² or R¹⁷ represents C₁₋₄alkyl substituted with one or where possible two or more substituents selected from C₁₋₄alkyloxy, hydroxy, halo, Het², NR⁷R⁸,

NR⁹R¹⁰-carbonyl or Het³-carbonyl;

Het¹ represents a heterocycle selected from piperidinyl, morpholinyl, piperazinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het¹ is optionally substituted with amino, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl-, phenyl, phenyl-C₁₋₄alkyl-,

C₁₋₄alkyl-oxy-C₁₋₄alkyl- mono- or di(C₁₋₄alkyl)amino- or amino-carbonyl-;

Het² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl, thiomorpholinyl or dithianyl wherein said Het² is optionally substituted with one or where possible two or more substituents selected from hydroxy, halo, amino, C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-oxy-C₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino-, mono- or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-, aminoC₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino-sulfonyl-, aminosulfonyl-;

Het³, Het⁴ and Het⁸ each independently represent a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het³, Het⁴ or Het⁸ is optionally substituted with one or where possible two or more substituents selected from hydroxy-, amino-, C₁₋₄alkyl-, C₃₋₆cycloalkyl-C₁₋₄alkyl-, aminosulfonyl-, mono- or di(C₁₋₄alkyl)aminosulfonyl or amino-C₁₋₄alkyl-;


Het⁵ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from

-58-

C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het⁶ and Het⁷ each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het⁹ and Het¹⁰ each independently represent a heterocycle selected from furanyl, piperidinyl, morpholinyl, piperazinyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het⁹ or Het¹⁰ is optionally substituted C₁₋₄alkyl, C₃₋₆cycloalkyl-C₁₋₄alkyl- or amino-C₁₋₄alkyl-;



Het¹¹ represents a heterocycle selected from indolyl or

Het¹² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl, thiomorpholinyl or dithianyl wherein said Het¹² is optionally

substituted with one or where possible two or more substituents selected from hydroxy, halo, amino, C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-oxy-C₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino- or mono- or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-;

Het¹⁶ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl, 1,3,2-dioxaborolane or piperidinyl wherein said heterocycle is optionally substituted with one or more substituents selected from C₁₋₄alkyl; and

Het¹⁷ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het¹⁸ and Het¹⁹ each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het²⁰, Het²¹ and Het²² each independently represent a heterocycle selected from pyrrolidinyl, 2-pyrrolidinonyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from hydroxy, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl- or polyhydroxy-C₁₋₄alkyl-;

Het²³ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more

-59-

substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-,
 C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;
 Ar¹, Ar³, Ar⁴ and Ar⁵ each independently represent phenyl optionally substituted with
 cyano, C₁₋₄alkylsulfonyl-, C₁₋₄alkylsulfonylamino-, aminosulfonylamino-,
 5 hydroxy-C₁₋₄alkyl, aminosulfonyl-, hydroxy-, C₁₋₄alkyloxy- or C₁₋₄alkyl.

In one embodiment the intermediates of formula (XXX) consists of the intermediates
 of formula (XXX) wherein one or more of the following restrictions apply;

Y₁ and Y₂ each independently represent C₁₋₅alkyl, CO-C₁₋₅alkyl or CO-CHR¹⁶-NH-;
 10 X¹ represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹¹,
 -NR¹¹-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or -C₁₋₂alkyl-;
 X² represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹²,
 -NR¹²-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or C₁₋₂alkyl-;
 R¹ represents hydrogen, cyano, halo, hydroxy, formyl, C₁₋₆alkoxy-, C₁₋₆alkyl-,
 15 C₁₋₆alkoxy- substituted with halo,
 C₁₋₄alkyl substituted with one or where possible two or more substituents selected
 from hydroxy or halo; and
 R² represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-, Het¹⁶-carbonyl-,
 C₁₋₄alkyloxycarbonyl-, C₁₋₄alkylcarbonyl-, aminocarbonyl-, mono- or
 20 di(C₁₋₄alkyl)aminocarbonyl-, Het¹, formyl, C₁₋₄alkyl-, C₂₋₆alkynyl-,
 C₃₋₆cycloalkyl-,
 C₃₋₆cycloalkyloxy-, C₁₋₆alkoxy-, Ar⁵, Ar¹-oxy-, dihydroxyborane ,
 C₁₋₆alkoxy- substituted with halo,
 C₁₋₄alkyl substituted with one or where possible two or more substituents selected
 25 from halo, hydroxy or NR⁵R⁶,
 C₁₋₄alkylcarbonyl- wherein said C₁₋₄alkyl is optionally substituted with one or
 where possible two or more substituents selected from hydroxy or
 C₁₋₄alkyl-oxy-;

R³ represents hydrogen;
 30 R⁵ and R⁶ are each independently selected from hydrogen or C₁₋₄alkyl;
 R⁷ and R⁸ are each independently selected from hydrogen, C₁₋₄alkyl, Het⁸,
 aminosulfonyl-, mono- or di (C₁₋₄alkyl)-aminosulfonyl, hydroxy-C₁₋₄alkyl-,
 C₁₋₄alkyl-oxy-C₁₋₄alkyl-, hydroxycarbonyl-C₁₋₄alkyl-, C₃₋₆cycloalkyl,
 Het⁹-carbonyl-C₁₋₄alkyl-, Het¹⁰-carbonyl-, polyhydroxy-C₁₋₄alkyl-,
 35 Het¹¹-C₁₋₄alkyl- or Ar²-C₁₋₄alkyl-;
 R⁹ and R¹⁰ are each independently selected from hydrogen, C₁₋₄alkyl, C₃₋₆cycloalkyl,
 Het⁴, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl- or polyhydroxy-C₁₋₄alkyl-;

-60-

- R^{11} represents hydrogen, C_{1-4} alkyl, Het^5 , Het^6-C_{1-4} alkyl-, C_{2-4} alkenylcarbonyl- optionally substituted with Het^7-C_{1-4} alkylaminocarbonyl-, C_{2-4} alkenylsulfonyl-, C_{1-4} alkyloxy C_{1-4} alkyl- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C_{1-4} alkyloxy-;
- R^{12} represents hydrogen, C_{1-4} alkyl, C_{1-4} alkyl-oxy-carbonyl-, Het^{17} , $Het^{18}-C_{1-4}$ alkyl-, C_{2-4} alkenylcarbonyl- optionally substituted with $Het^{19}-C_{1-4}$ alkylaminocarbonyl-, C_{2-4} alkenylsulfonyl-, C_{1-4} alkyloxy C_{1-4} alkyl- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C_{1-4} alkyloxy-;
- R^{16} represents hydrogen or C_{1-4} alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl, aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;
- R^{23} represents Ar^3 , Ar^4-C_{1-4} alkyl, C_{1-4} alkyl, C_{2-6} alkenyl optionally substituted with Het^{12} or R^{17} represents C_{1-4} alkyl substituted with one or where possible two or more substituents selected from C_{1-4} alkyloxy, hydroxy, halo, Het^2 , NR^7R^8 , NR^9R^{10} -carbonyl or Het^3 -carbonyl
- Het^1 represents a heterocycle selected from piperidinyl, morpholinyl, piperazinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het^1 is optionally substituted with amino, C_{1-4} alkyl, hydroxy- C_{1-4} alkyl-, phenyl, phenyl- C_{1-4} alkyl-, C_{1-4} alkyl-oxy- C_{1-4} alkyl- mono- or di(C_{1-4} alkyl)amino- or amino-carbonyl-;
- Het^2 represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl, thiomorpholinyl or dithianyl wherein said Het^2 is optionally substituted with one or where possible two or more substituents selected from hydroxy, halo, amino, C_{1-4} alkyl-, hydroxy- C_{1-4} alkyl-, C_{1-4} alkyl-oxy- C_{1-4} alkyl-, hydroxy- C_{1-4} alkyl-oxy- C_{1-4} alkyl-, mono- or di(C_{1-4} alkyl)amino-, mono- or di(C_{1-4} alkyl)amino- C_{1-4} alkyl-, amino C_{1-4} alkyl-, mono- or di(C_{1-4} alkyl)amino-sulfonyl-, aminosulfonyl-;
- Het^3 , Het^4 and Het^8 each independently represent a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het^3 , Het^4 or Het^8 is optionally substituted with one or where possible two or more substituents selected from hydroxy-, amino-, C_{1-4} alkyl-, C_{3-6} cycloalkyl- C_{1-4} alkyl-, aminosulfonyl-, mono- or di(C_{1-4} alkyl)aminosulfonyl or amino- C_{1-4} alkyl-;


-61-

Het⁵ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

5 Het⁶ and Het⁷ each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

10 Het⁹ and Het¹⁰ each independently represent a heterocycle selected from furanyl, piperidinyl, morpholinyl, piperazinyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het⁹ or Het¹⁰ is optionally substituted C₁₋₄alkyl, C₃₋₆cycloalkyl-C₁₋₄alkyl- or amino-C₁₋₄alkyl-;

Het¹¹ represents a heterocycle selected from indolyl or



15 Het¹² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl, thiomorpholinyl or dithianyl wherein said Het¹² is optionally substituted with one or where possible two or more substituents selected from hydroxy, halo, amino, C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-oxy-C₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino- or mono- or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-;

20 Het¹⁶ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl, 1,3,2-dioxaborolane or piperidinyl wherein said heterocycle is optionally substituted with one or more substituents selected from C₁₋₄alkyl; and

25 Het¹⁷ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

30 Het¹⁸ and Het¹⁹ each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;


35 Ar¹, Ar³, Ar⁴ and Ar⁵ each independently represent phenyl optionally substituted with cyano, C₁₋₄alkylsulfonyl-, C₁₋₄alkylsulfonylamino-, aminosulfonylamino-, hydroxy-C₁₋₄alkyl, aminosulfonyl-, hydroxy-, C₁₋₄alkyloxy- or C₁₋₄alkyl.

-62-

In particular the intermediates of formula (XXX) wherein one or more of the following restrictions apply;

- i) X^1 represents O, $-O-C_{1-2}alkyl-$, $-O-N=CH-$, NR^{11} or $-NR^{11}-C_{1-2}alkyl-$; in a particular embodiment X^1 represents $-NR^{11}-$, $-O-$ or $-O-CH_2-$;
- 5 ii) X^2 represents a direct bond, O, $-O-C_{1-2}alkyl-$, $-O-N=CH-$, NR^{12} or $NR^{12}-C_{1-2}alkyl-$; in a particular embodiment X^2 represents a direct bond, $-C_{1-2}alkyl-$, $-O-C_{1-2}alkyl-$, $-O-$ or $-O-CH_2-$;
- iii) R^1 represents hydrogen, cyano, halo or hydroxy, preferably halo;
- 10 iv) R^2 represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-, $C_{1-4}alkyloxycarbonyl-$, Het^{16} -carbonyl-, $C_{1-4}alkyl-$, $C_{2-6}alkynyl-$, Ar^5 or Het^1 ; In a further embodiment R^2 represents hydrogen, cyano, halo, hydroxy, $C_{2-6}alkynyl-$ or Het^1 ; in particular R^2 represents hydrogen, cyano, halo, hydroxy, or Ar^5 ;
- 15 v) R^{23} represents hydrogen, $C_{1-4}alkyl$ or R^{17} represents $C_{1-4}alkyl$ substituted with one or where possible two or more substituents selected from $C_{1-4}alkyloxy-$ or Het^2- ;
- vi) R^{12} represents hydrogen, $C_{1-4}alkyl-$ or $C_{1-4}alkyl-oxy-carbonyl-$;
- vii) Het^1 represents thiazolyl optionally substituted with amino, $C_{1-4}alkyl$, hydroxy- $C_{1-4}alkyl-$, phenyl, phenyl- $C_{1-4}alkyl-$, $C_{1-4}alkyl-oxy-C_{1-4}alkyl-$ mono- or
- 20 di($C_{1-4}alkyl$)amino- or amino-carbonyl-;
- viii) Het^2 represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or pyrrolidinyl wherein said Het^2 is optionally substituted with one or where possible two or more substituents selected from hydroxy, amino or $C_{1-4}alkyl-$; In a further embodiment Het^2 represents a heterocycle selected from morpholinyl or piperidinyl optionally substituted with $C_{1-4}alkyl-$, preferably methyl;
- 25 ix) Het^3 represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or pyrrolidinyl wherein said Het^3 is optionally substituted with one or where possible two or more substituents selected from hydroxy, amino or $C_{1-4}alkyl-$;
- x) Het^{12} represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or pyrrolidinyl wherein said Het^{12} is optionally substituted with one or where
- 30 possible two or more substituents selected from hydroxy, amino or $C_{1-4}alkyl-$;
- xi) Het^{16} represents a heterocycle selected from piperidinyl or pyrrolidinyl.

35 c) the intermediates of formula (XXXIII)



the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

m represents 1, 2, 3 or 4;

- 5 X^2 represents a direct bond, O, $-O-C_{1-2}\text{alkyl-}$, CO, $-CO-C_{1-2}\text{alkyl-}$, NR^{12} , $-NR^{12}-C_{1-2}\text{alkyl-}$, $-CH_2-$, $-O-N=CH-$ or $C_{1-2}\text{alkyl-}$;
- Y_3 represents a $C_{1-5}\text{alkyl}$, $CO-C_{1-5}\text{alkyl}$ or $CO-CR^{16}R^{17}-NH-$ or $C_{1-5}\text{alkyl-CO-}$ optionally substituted with amino, mono- or di($C_{1-4}\text{alkyl}$)amino or $C_{1-4}\text{alkyloxycarbonylamino}$;
- 10 R^1 represents hydrogen, cyano, halo, hydroxy, formyl, $C_{1-6}\text{alkoxy-}$, $C_{1-6}\text{alkyl-}$, halo-phenyl-carbonylamino-, $C_{1-6}\text{alkoxy-}$ substituted with halo, $C_{1-4}\text{alkyl}$ substituted with one or where possible two or more substituents selected from hydroxy or halo;
- 15 R^2 represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-, Het^{16} -carbonyl-, $C_{1-4}\text{alkyloxycarbonyl-}$, $C_{1-4}\text{alkylcarbonyl-}$, aminocarbonyl-, mono- or di($C_{1-4}\text{alkyl}$)aminocarbonyl-, Het^1 , formyl, $C_{1-4}\text{alkyl-}$, $C_{2-6}\text{alkynyl-}$, $C_{3-6}\text{cycloalkyl-}$, $C_{3-6}\text{cycloalkyloxy-}$, $C_{1-6}\text{alkoxy-}$, Ar^5 , Ar^1 -oxy-, dihydroxyborane, $C_{1-6}\text{alkoxy-}$ substituted with halo,
- 20 $C_{1-4}\text{alkyl}$ substituted with one or where possible two or more substituents selected from halo, hydroxy or NR^5R^6 , $C_{1-4}\text{alkylcarbonyl-}$ wherein said $C_{1-4}\text{alkyl}$ is optionally substituted with one or where possible two or more substituents selected from hydroxy or $C_{1-4}\text{alkyl-oxy-}$;
- 25 R^3 represents hydrogen, $C_{1-4}\text{alkyl}$, or $C_{1-4}\text{alkyl}$ substituted with one or more substituents selected from halo, $C_{1-4}\text{alkyloxy-}$, amino-, mono- or di($C_{1-4}\text{alkyl}$)amino-, $C_{1-4}\text{alkyl-sulfonyl-}$ or phenyl;
- R^5 and R^6 are each independently selected from hydrogen or $C_{1-4}\text{alkyl}$;
- 30 R^{12} represents hydrogen, $C_{1-4}\text{alkyl}$, $C_{1-4}\text{alkyl-oxy-carbonyl-}$, $Het^{18}-C_{1-4}\text{alkyl-}$, phenyl- $C_{1-4}\text{alkyl-oxy-carbonyl-}$ Het^{17} , $C_{2-4}\text{alkenylcarbonyl-}$ optionally substituted with $Het^{19}-C_{1-4}\text{alkylaminocarbonyl-}$, $C_{2-4}\text{alkenylsulfonyl-}$, $C_{1-4}\text{alkyloxy}$ $C_{1-4}\text{alkyl-}$

-64-

- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;
- R¹⁶ and R¹⁷ each independently represents hydrogen or C₁₋₄alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl, aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;
- Het¹ represents a heterocycle selected from piperidinyl, morpholinyl, piperazinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het¹ is optionally substituted with amino, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl-, phenyl, phenyl-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl- mono- or di(C₁₋₄alkyl)amino- or amino-carbonyl-;
- Het¹⁶ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl, 1,3,2-dioxaborolane or piperidinyl wherein said heterocycle is optionally substituted with one or more substituents selected from C₁₋₄alkyl;
- Het¹⁸ and Het¹⁹ each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-; and
- Ar¹ and Ar⁵ each independently represent phenyl optionally substituted with cyano, C₁₋₄alkylsulfonyl-, C₁₋₄alkylsulfonylamino-, aminosulfonylamino-, hydroxy-C₁₋₄alkyl, aminosulfonyl-, hydroxy-, C₁₋₄alkyloxy- or C₁₋₄alkyl.

In particular the intermediates of formula (XXXIII) wherein one or more of the following restrictions apply;

- i) X² represents a direct bond, C₁₋₂alkyl, NR¹² or -NR¹²-C₁₋₂alkyl-; in a particular embodiment X² represents -NR¹²-C₁₋₂alkyl- or C₁₋₂alkyl;
- ii) Y₃ represents a C₁₋₅alkyl, CO-CR¹⁶R¹⁷-NH- or -C₁₋₅alkyl-CO-; in a particular embodiment Y₃ represents -C₁₋₅alkyl-CO-;
- iii) R¹ represents hydrogen, cyano, halo or hydroxy, preferably halo;
- iv) R² represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-, C₁₋₄alkyloxycarbonyl-, Het¹⁶-carbonyl-, C₁₋₄alkyl-, C₂₋₆alkynyl-, Ar⁵ or Het¹; In a further embodiment R² represents hydrogen, cyano, halo or hydroxy; in particular R² represents hydrogen, cyano, halo, hydroxy, or Ar⁵;
- v) R³ represents hydrogen;
- vi) R¹² represents hydrogen or C₁₋₄alkyl;
- vii) R¹⁶ represents hydrogen or C₁₋₄alkyl substituted with hydroxy;
- viii) R¹⁷ represents hydrogen or C₁₋₄alkyl, in particular hydrogen or methyl;

-65-

- ix) Het¹ represents thiazolyl optionally substituted with amino, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl-, phenyl, phenyl-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl- mono- or di(C₁₋₄alkyl)amino- or amino-carbonyl-;
- x) Het¹⁶ represents a heterocycle selected from piperidiny1 or pyrrolidinyl;
- 5 xi) Ar⁵ represents phenyl.

It is also an object of the present invention to provide the use of the intermediates of formula (III), (XXX) or (XXXIII) in the synthesis of a compound of formula (I).

- 10 The compounds of the present invention, including the compounds of formula (I) and the intermediates of formula (III), (XXX) and (XXXIII) are useful because they possess pharmacological properties. They can therefore be used as medicines.

- As described in the experimental part hereinafter, the growth inhibitory effect and anti-tumour activity of the present compounds has been demonstrated in vitro, in enzymatic assays on the receptor tyrosine kinases EGFR, ErbB2, ErbB4, FIT3, BLK or the Sar kinase family such as for example Lyn, Yes cSRC. In an alternative assay, the growth inhibitory effect of the compounds was tested on a number of carcinoma cell lines, in particular in the ovarian carcinoma cell line SKOV3 and the squamous carcinoma cell line A431 using art known cytotoxicity assays such as MTT.
- 15
- 20

- Accordingly, the present invention provides the compounds of formula (I) and their pharmaceutically acceptable *N*-oxides, addition salts, quaternary amines and stereochemically isomeric forms for use in therapy. More particular in the treatment or prevention of cell proliferation mediated diseases. The compounds of formula (I) and their pharmaceutically acceptable *N*-oxides, addition salts, quaternary amines and the stereochemically isomeric forms may hereinafter be referred to as compounds according to the invention.
- 25

- 30 Disorders for which the compounds according to the invention are particularly useful are atherosclerosis, restenosis, cancer and diabetic complications e.g. retinopathy.

- In view of the utility of the compounds according to the invention, there is provided a method of treating a cell proliferative disorder such as atherosclerosis, restenosis and cancer, the method comprising administering to an animal in need of such treatment, for example, a mammal including humans, suffering from a cell proliferative disorder, a therapeutically effective amount of a compound according to the present invention.
- 35

-66-

Said method comprising the systemic or topical administration of an effective amount of a compound according to the invention, to animals, including humans. One skilled in the art will recognize that a therapeutically effective amount of the EGFR inhibitors of the present invention is the amount sufficient to induce the growth inhibitory effect and that this amount varies *inter alia*, depending on the size, the type of the neoplasia, the concentration of the compound in the therapeutic formulation, and the condition of the patient. Generally, an amount of EGFR inhibitor to be administered as a therapeutic agent for treating cell proliferative disorder such as atherosclerosis, restenosis and cancer, will be determined on a case by case by an attending physician.

Generally, a suitable dose is one that results in a concentration of the EGFR inhibitor at the treatment site in the range of 0.5 nM to 200 μ M, and more usually 5 nM to 10 μ M. To obtain these treatment concentrations, a patient in need of treatment likely will be administered between 0.01 mg/kg to 300 mg/kg body weight, in particular from 10 mg/kg to 100 mg/kg body weight. As noted above, the above amounts may vary on a case-by-case basis. In these methods of treatment the compounds according to the invention are preferably formulated prior to admission. As described herein below, suitable pharmaceutical formulations are prepared by known procedures using well known and readily available ingredients.

Due to their high degree of selectivity as EGFR inhibitors, the compounds of formula (I) as defined above, are also useful to mark or identify the kinase domain within the receptor tyrosine kinase receptors. To this purpose, the compounds of the present invention can be labelled, in particular by replacing, partially or completely, one or more atoms in the molecule by their radioactive isotopes. Examples of interesting labelled compounds are those compounds having at least one halo which is a radioactive isotope of iodine, bromine or fluorine; or those compounds having at least one ^{11}C -atom or tritium atom.

One particular group consists of those compounds of formula (I) wherein R^1 is a radioactive halogen atom. In principle, any compound of formula (I) containing a halogen atom is prone for radiolabelling by replacing the halogen atom by a suitable isotope. Suitable halogen radioisotopes to this purpose are radioactive iodides, e.g. ^{122}I , ^{123}I , ^{125}I , ^{131}I ; radioactive bromides, e.g. ^{75}Br , ^{76}Br , ^{77}Br and ^{82}Br , and radioactive fluorides, e.g. ^{18}F . The introduction of a radioactive halogen atom can be performed by a suitable exchange reaction or by using any one of the procedures as described

-67-

hereinabove to prepare halogen derivatives of formula (I).

Another interesting form of radiolabelling is by substituting a carbon atom by a ^{11}C -atom or the substitution of a hydrogen atom by a tritium atom.

Hence, said radiolabelled compounds of formula (I) can be used in a process of specifically marking receptor sites in biological material. Said process comprises the steps of (a) radiolabelling a compound of formula (I), (b) administering this radiolabelled compound to biological material and subsequently (c) detecting the emissions from the radiolabelled compound.

The term biological material is meant to comprise every kind of material which has a biological origin. More in particular this term refers to tissue samples, plasma or body fluids but also to animals, specially warm-blooded animals, or parts of animals such as organs.

When used in *in vivo* assays, the radiolabelled compounds are administered in an appropriate composition to an animal and the location of said radiolabelled compounds is detected using imaging techniques, such as, for instance, Single Photon Emission Computerized Tomography (SPECT) or Positron Emission Tomography (PET) and the like. In this manner the distribution of the particular receptor sites throughout the body can be detected and organs containing said receptor sites can be visualized by the imaging techniques mentioned hereinabove. This process of imaging an organ by administering a radiolabelled compound of formula (I) and detecting the emissions from the radioactive compound also constitutes a part of the present invention.

In yet a further aspect, the present invention provides the use of the compounds according to the invention in the manufacture of a medicament for treating any of the aforementioned cell proliferative disorders or indications.

The amount of a compound according to the present invention, also referred to here as the active ingredient, which is required to achieve a therapeutical effect will be, of course, vary with the particular compound, the route of administration, the age and condition of the recipient, and the particular disorder or disease being treated. A suitable daily dose would be from 0.01 mg/kg to 300 mg/kg body weight, in particular from 10 mg/kg to 100 mg/kg body weight. A method of treatment may also include administering the active ingredient on a regimen of between one and four intakes per day.

While it is possible for the active ingredient to be administered alone, it is preferable

-68-

to present it as a pharmaceutical composition. Accordingly, the present invention further provides a pharmaceutical composition comprising a compound according to the present invention, together with a pharmaceutically acceptable carrier or diluent. The carrier or diluent must be "acceptable" in the sense of being compatible with the other ingredients of the composition and not deleterious to the recipients thereof.

The pharmaceutical compositions of this invention may be prepared by any methods well known in the art of pharmacy, for example, using methods such as those described in Gennaro et al. Remington's Pharmaceutical Sciences (18th ed., Mack Publishing Company, 1990, see especially Part 8 : Pharmaceutical preparations and their Manufacture). A therapeutically effective amount of the particular compound, in base form or addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable, preferably, for systemic administration such as oral, percutaneous or parenteral administration; or topical administration such as via inhalation, a nose spray, eye drops or via a cream, gel, shampoo or the like. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wettable agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not cause any significant deleterious effects on the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on or as an ointment.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

Experimental part

Hereinafter, the term 'RT' means room temperature, 'ADDP' means 1,1'-(azodicarbonyl)dipiperidine, 'DCM' dichloromethane, 'DMA' means dimethylacetamide, 'DME' means dimethyl ether, 'DMF' means *N,N*-dimethylformamide, 'DMSO' means dimethylsulfoxide, 'DIPE' means diisopropyl ether, 'DIPEA' means *N*-ethyl-*N*-(1-methylethyl)-2-propanamine, 'EtOH' means ethanol, 'EtOAc' means ethyl acetate, 'HBTU' means 1-[bis(dimethylamino)methylene]-*H*-benzotriazolium, hexafluorophosphate(1-), 3-oxide) 'LAH' means lithiumaluminiumhydride i.e. LiAlH_4 , 'TFA' means trifluoroacetic acid and 'THF' means tetrahydrofuran, 'PyBOP' means (1-hydroxy-1*H*-benzotriazolato-*O*)tri-1-pyrrolidinyl-, (T-4)-phosphorus(1+), hexafluorophosphate(1-), 'NaBH(OAc)₃' means sodium triacetoxyborohydride, RP means reversed-phase.

A. Preparation of the intermediates

Example A1

a) Preparation of hexanoic acid, 6-(2-methyl-6-nitrophenoxy)-, methyl ester (intermediate 1)

A mixture of 2-methyl-6-nitro-phenol (0.0065 mol) and K_2CO_3 (0.026 mol) in *N,N*-dimethyl-formamide (DMF) (80 ml) was stirred at 50 °C for 15 min., then 6-bromo-,methyl ester hexanoic acid (0.0195 mol) was added dropwise and the reaction mixture was stirred for 18 hours at 50°C. After completion, the reaction was quenched with ice-water and the mixture was extracted 3 times with toluene. The organic layer

-70-

was separated, dried (MgSO_4), filtered and concentrated. The residue was used as such in the next step, yielding 100% of intermediate 1.

b) Preparation of hexanoic acid, 6-(2-amino-6-methylphenoxy)-, methyl ester (intermediate 2)

A mixture of intermediate 1 (0.013 mol) and ethylamine (0.5 g) in THF (100 ml) was hydrogenated with Pt/C 5% (2 g) as a catalyst. After uptake of H_2 (3 equiv.), the reaction mixture was filtered over a small plug of Dicalite and the filtrate was concentrated, yielding 1.4 g of intermediate 2 which was used as such in the next step.

c) Preparation of hexanoic acid, 6-[2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-6-methylphenoxy]-, methyl ester (intermediate 3)

A mixture of 4-chloro-6-methylcarbonyloxy-7-methoxyquinazoline (0.0045 mol) and intermediate 2 (0.0056 mol) in 2-propanol (40 ml) was stirred and refluxed for 1 day. The reaction mixture was concentrated and the residue, treated with DIPE and this mixture was stirred overnight. The solid was collected by filtration, washed and dried, yielding intermediate 3.

d) Preparation of hexanoic acid, 6-[2-[(6-hydroxy-7-methoxy-4-quinazolinyl)amino]-6-methylphenoxy]-, methyl ester (intermediate 4)

A solution of intermediate 3 (0.0045 mol) and NH_4OH (1.5 ml) in CH_3OH (50 ml) was stirred for 18 hours at RT and the solvent was evaporated, yielding intermediate 4 (impure, used as such in the next reaction step).

e) Preparation of hexanoic acid, 6-[2-[(6-hydroxy-7-methoxy-4-quinazolinyl)amino]-6-methylphenoxy] (intermediate 5)

A mixture of intermediate 4 (0.00024 mol), LiOH (0.00047 mol), THF (3 ml), CH_3OH (1 ml) and H_2O (1 ml) was stirred and heated at 70 °C for 30 min. and then the reaction mixture was allowed to reach RT. The organic solvent (THF/ CH_3OH) was evaporated and the aqueous concentrate was neutralised with HCl (1N) filtered and the solid retained was washed and dried (vac.) at 65 °C, yielding 0.040 g of intermediate 5.

Example A2

a) Preparation of hexanoic acid, 6-(2-chloro-6-nitrophenoxy)-, methyl ester (intermediate 6)

A solution of 2-chloro-6-nitro-phenol (0.046 mol) in N,N-dimethylformamide (150 ml) was heated to 50 °C, then K_2CO_3 (0.069 mol) was added and the reaction mixture was stirred for 15 min. 6-Bromo-,methyl ester hexanoic acid (0.069 mol) was added and the mixture was stirred overnight. The reaction mixture was filtered and the

-71-

filtrate was concentrated and the residue was used as such in the next step, yielding 13.88 g of intermediate 6.

b) Preparation of hexanoic acid, 6-(2-amino-6-chlorophenoxy)-, methyl ester (intermediate 7)

A mixture of intermediate 6 (0.046 mol) and ethanime (2 g) in THF (ml) was hydrogenated with Pt/C 5% (3 g) as a catalyst in the presence of DIPE (2 ml). After uptake of H₂ (3 equiv.), the reaction mixture was filtered over small plug of Dicalite the filtrate was concentrated, yielding intermediate 7.

c) Preparation of hexanoic acid, 6-[2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-6-chlorophenoxy]-, methyl ester (intermediate 8)

A mixture of 4-chloro-6-methylcarbonyloxy-7-methoxyquinazoline (0.022 mol) and intermediate 7 (0.022 mol) in 2-propanol (170 ml) was stirred and heated at 80 °C for 2 hours, concentrated and the residue was chromatographed over silica gel (eluent: DCM/CH₃OH 97/3). The product fractions were collected and the solvent was evaporated, yielding 5.1 g intermediate 8 (used as such in the next reaction step).

d) Preparation of 6-quinazolinol, 4-[[3-chloro-2-[(6-hydroxyhexyl)oxy]phenyl]amino]-7-methoxy-(intermediate 9)

A mixture of LAH (0.0246 mol) in THF (40 ml) was stirred at RT. A solution of intermediate 8 (0.006 mol) in THF (60 ml) was added dropwise. The reaction mixture was stirred for 1 day then, extra LAH (0.0123 mol) was added portionwise. The mixture was stirred further over the weekend then, H₂O (2 ml) was added dropwise, followed by the dropwise addition of a 15 % NaOH soln. (2 ml) and H₂O (6 ml). This mixture was stirred for 15 min filtered and the filtrate was concentrated The residue was stirred in boiling CH₃CN, filtered and the solid retained was dried (vac.) at 60 °C. The solids were re-dissolved in CH₃OH/DCM (10/90) and this mixture was neutralised with HCl (1N). The organic layer was separated, dried (MgSO₄), filtered and concentrated, yielding 1 g of intermediate 9.

Example A3

a) Preparation of hexanoic acid, 6-(4-chloro-2-nitrophenoxy)-, methyl ester (intermediate 10)

A mixture of 4-chloro-6-nitro-phenol (0.029 mol) and K₂CO₃ (0.035 mol) in DMA (80 ml) was stirred at 50 °C for 30 min., then 6-bromo-,methyl ester hexanoic acid (0.035 mol) was added dropwise and the reaction mixture was stirred for another 18 hours at 50 °C. After completion, the mixture was filtered and the filtrate was neutralised with HCl (1N), then poured onto ice water and stirred for 30 min. The resulting precipitate was collected by filtration, washed, dissolved in DCM, dried

-72-

(MgSO₄), filtered and concentrated, yielding intermediate 10 (used as such in the next reaction step).

b) Preparation of hexanoic acid, 6-(2-amino-4-chlorophenoxy)-, methyl ester (intermediate 11)

A mixture of intermediate 10 (0.026 mol) and ethylamine (1 g) in THF (100 ml) was hydrogenated with Pt/C 5% (5 g) as a catalyst in the presence of DIPE (1 ml). After uptake of H₂ (3 equiv.), and the reaction mixture was filtered over a small plug of Dicalite, the filtrate was concentrated, yielding intermediate 11.

c) Preparation of hexanoic acid, 6-[2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-4-chlorophenoxy]-, methyl ester (intermediate 12)

A mixture of 4-chloro-6-methylcarbonyloxy-7-methoxyquinazoline (0.014 mol) and intermediate 11 (0.014 mol) in 2-propanol (120 ml) was heated at 80 °C and stirred for 3 hours. The reaction mixture was filtered and the filtrate was concentrated. The residue was purified by column chromatography over silica gel (eluent: DCM/CH₃OH 96.5/3.5). The product fractions were collected and the solvent was evaporated to dryness, yielding 1.8 g of intermediate 12 (used as such in the next reaction step).

d) Preparation of 6-quinazolinol, 4-[[5-chloro-2-[(6-hydroxyhexyl)oxy]phenyl]amino]-7-methoxy- (intermediate 13)

LAH (0.015 mol) was stirred in THF (40 ml) and then a solution of intermediate 12 (0.0037 mol) in THF (80 ml) was added dropwise under N₂ at RT. The reaction mixture was stirred over the weekend, then treated with H₂O (0.9 ml), aq. NaOH soln. (15 %, 0.9 ml) and H₂O (2.7 ml). The reaction mixture was filtered, the residue washed and the filtrate was concentrated under reduced pressure. The residue was stirred in DIPE and the solid was collected by filtration, yielding 0.8 g (53 %) of intermediate 13.

Example A4

a) Preparation of pentanoic acid, 5-(4-chloro-2-nitrophenoxy)-, methyl ester (intermediate 14)

A mixture of 4-chloro-6-nitrophenol (0.023 mol), K₂CO₃ (0.027 mol) and N,N-dimethylformamide (80 ml) was stirred at 50 °C for 30 min., then 5-bromo-, methylester pentanoic acid (0.027 mol) was added dropwise and the reaction mixture was stirred for 18 hours at 50 °C. The mixture was filtered and the filtrate was neutralised with HCl (1N). This mixture was poured in ice water and stirred for 30 min. The resulting precipitate was collected by filtration, washed, re-dissolved in

-73-

DCM/CH₃OH(95/5) dried (MgSO₄), filtered and concentrated, yielding 6.6 g of intermediate 14 (used as such in the next reaction step).

b) Preparation of pentanoic acid, 5-(2-amino-4-chlorophenoxy)-, methyl ester (intermediate 15)

A mixture of intermediate 14 (0.023 mol) and ethylamine (1 g) in THF (100 ml) was hydrogenated with Pt/C 5% (2 g) as a catalyst in the presence of DIPE (1 ml). After uptake of H₂ (3 equiv.), the reaction mixture was over a small plug of Dicalite and the filtrate was concentrated, yielding intermediate 15.

c) Preparation of pentanoic acid, 5-[2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-4-chlorophenoxy]-, methyl ester (intermediate 16)

A mixture of 4-chloro-6-methylcarbonyloxy-7-methoxyquinazoline (0.0067 mol) and intermediate 15 (0.0048 mol) in 2-propanol (60 ml) was stirred and heated at 80 °C for 4 hours. The reaction mixture was filtered, the solid retained was washed and dried 0.7 g. The filtrate was concentrated and the residue (oil) was chromatographed over silica gel (eluent: DCM/CH₃OH 96.5/3.5). The pure fraction were collected and concentrated, yielding 1.5 g of intermediate 16.

d) Preparation of 6-quinazolinol, 4-[[5-chloro-2-[(5-hydroxypentyl)oxy]phenyl]amino]-7-methoxy- (intermediate 17)

A mixture of LAH (0.013 mol) in THF (25 ml) was stirred at RT under N₂, then a solution of intermediate 16 (0.0032 mol) in THF (45 ml) was added dropwise and the reaction mixture was stirred overnight. The reaction mixture was treated with water (0.8 ml), NaOH (0.8 ml, 15 %) and again water (2.4 ml) and the mixture was stirred for 15 min. The mixture was filtered and the filtrate was concentrated, re-dissolved in DCM/CH₃OH (95/5), neutralised with HCl (1N), concentrated and then purified over silica gel (eluent: DCM/CH₃OH 91.5/8.5). The product fractions were collected and concentrated, yielding 0.400 g of intermediate 17.

Example A5

a) Preparation of hexanoic acid, 6-(2-nitrophenoxy)-, methyl ester (intermediate 18)

A mixture of 2-nitro-phenol (0.014 mol) and K₂CO₃ (0.017 mol) in N, N-dimethylformamide (50 ml) was stirred for 15 min. at 50 °C, then 6-bromo-hexanoic acid methyl ester (0.017 mol) was added dropwise and the reaction mixture was stirred for 18 hours at 50 °C. The reaction mixture was filtered and poured onto ice water. The resulting precipitate was filtered, washed and dried, yielding 3.0 g of intermediate 18 (used as such in the next reaction step).

-74-

b) Preparation of hexanoic acid, 6-(2-aminophenoxy)-, methyl ester (intermediate 19)

A mixture of intermediate 18 (0.011 mol) in THF (100 ml) was hydrogenated with Pt/C 5% (0.5 g) as a catalyst in the presence of DIPE (0.5 ml). After uptake of H₂ (3 equiv.), the reaction mixture was filtered over a small plug of Dicalite and concentrated, yielding intermediate 19.

c) Preparation of hexanoic acid, 6-[2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]phenoxy]-, methyl ester (intermediate 20)

5 A mixture of 4-chloro-6-methylcarbonyloxy-7-methoxyquinazoline (0.011 mol) and intermediate 19 (0.011 mol) in 2-propanol (100 ml) was heated and stirred at 80 °C for 5 hours. The reaction mixture was concentrated and the residue (oil) was purified over silica gel (eluent: DCM/CH₃OH 96.5/3.5). The product fractions were collected and and concentrated, yielding 2.3 g of intermediate 20 (used as such in the next reaction step).

d) Preparation of 6-quinazolinol, 4-[[2-[(6-hydroxyhexyl)oxy]phenyl]amino]-7-methoxy- (intermediate 21)

15 A mixture of LAH (0.020 mol) in THF (30 ml) was stirred at RT and then a solution of intermediate 20 (0.005 mol) in THF (50 ml) was added dropwise. The reaction mixture was stirred overnight, treated with H₂O (1 ml), aqueous NaOH soln. (1 ml, 15 %) and again with H₂O (3 ml). This mixture was filtered, the residue was washed and the filtrate was neutralised with HCl (1N). The filtrate was then concentrated and the residue was dried (vac.) at 55 °C, yielding 0.5 g of intermediate 21.

Example A6

a) Preparation of hexanoic acid, 6-(4-bromo-2-nitrophenoxy)-, methyl ester (intermediate 22)

20 A mixture of 4-bromo-2-nitrophenol (0.046 mol) in DMA (100 ml) was heated to 40°C and then K₂CO₃ (0.046 mol) was added. The reaction mixture was stirred for 15 min. 6-Bromo-hexanoic acid methyl ester (0.046 mol) was added and the mixture was stirred overnight at 40 °C. Extra 6-bromo-hexanoic acid methyl ester (2 g) was added and the reaction mixture was stirred for another 2 hours. The mixture was cooled to RT and poured into ice water (400 ml). The resulting precipitate was filtered, dissolved in DCM, dried (MgSO₄) and filtered again. Finally, the filtrate was evaporated, yielding 14.24 g (90 %) of intermediate 22.

25 b) Preparation of hexanoic acid, 6-(2-amino-4-bromophenoxy)-, methyl ester (intermediate 23)

-75-

A mixture of intermediate 22 (0.04 mol) and ethylamine (0.044 mol) in THF (250 ml) was hydrogenated with Pt/C 5% (2 g) as a catalyst in the presence of DIPE (2 ml). After uptake of H₂ (3 equiv.), the reaction mixture was filtered over a small plug of Dicalite and concentrated, yielding 12.8 g of intermediate 23 (98 %).

c) Preparation of hexanoic acid, 6-[2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-4-bromophenoxy]-, methyl ester (intermediate 24)

- 5 A mixture of 4-chloro-6-acetoxy-7-methoxyquinazoline hydrochloride (0.00554 mol) and intermediate 23 (0.00554 mol) in 2-propanol (10 ml) was heated to 80 °C. After 1 hour, the reaction mixture was homogeneous and was coloured black. The mixture was stirred overnight at RT, concentrated and then purified by column chromatography over silica gel. Two product fractions were collected and
- 10 concentrated. Fraction 2 was stirred in 2-propanol/DIPE (1/24), filtered and the solid retained was dried, yielding intermediate 24.

d) Preparation of 6-quinazolinol, 4-[[5-bromo-2-[(6-hydroxyhexyl)oxy]phenyl]amino]-7-methoxy- (intermediate 25)

- A mixture of intermediate 24 (0.00188 mol) in THF (40 ml) was added dropwise to a stirred suspension of LAH (0.0075 mol) in THF (20 ml) at RT under N₂-atm., for 16h. The reaction mixture was subsequently treated with H₂O (0.4 ml), and after 15 min
- 15 with aqueous NaOH soln. (0.4 ml, 15 %), and finally with H₂O (1.2 ml) (colour change from grey/green to yellow). The reaction mixture was filtered and the filtrate was neutralised with HCl (1N) and concentrated. The residue was stirred in CH₃CN/DIPE (24/1) and the solid was collected by filtration and dried, yielding
- 20 intermediate 25.

Example A7

a) Preparation of heptanoic acid, 7-(4-chloro-2-nitrophenoxy)-, ethyl ester (intermediate 26)

- A mixture of 4-chloro-6-nitro-phenol (0.017 mol) and K₂CO₃ (0.019 mol) in DMA (70 ml) was stirred at 50 °C for 15 min., then 6-bromo-,ethyl ester hexanoic acid (0.019 mol) was added and the reaction mixture was stirred overnight at 50 °C. The
- 25 resulting precipitate was filtered and the filtrate was concentrated under reduced pressure, yielding intermediate 26 (used as such in the next reaction step).

b) Preparation of heptanoic acid, 7-(2-amino-4-chlorophenoxy)-, ethyl ester (intermediate 27)

A mixture of intermediate 26 (0.017mol) and dimethyl-amine (1 g) in THF (100 ml) was hydrogenated with Pt/C 5% (2 g) as a catalyst in the presence of DIPE (1 ml).

-76-

After uptake of H₂ (3 equiv.), The reaction mixture was filtered over a small plug of Dicalite and concentrated, yielding intermediate 27.

c) Preparation of heptanoic acid, 7-[2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-4-chlorophenoxy]-, ethyl ester (intermediate 28)

A mixture of 4-chloro-6-methylcarbonyloxy-7-methoxyquinazoline (0.0051 mol) and intermediate 27 (0.006 mol) in 2-propanol (50 ml) was stirred and heated at 80 °C for 6 hours, then the solvent was evaporated under reduced pressure. The crude residue was purified by column chromatography over silica gel (eluent: DCM/CH₃OH 97.5/2.5). The product fractions were collected and concentrated, yielding 2.4 g (92 %) of intermediate 28.

d) Preparation of 6-quinazolinol, 4-[[5-chloro-2-[(7-hydroxyheptyl)oxy]phenyl]amino]-7-methoxy- (intermediate 29)

A mixture of LAH (0.0186 mol) in THF (40 ml) was stirred at RT and then a solution of intermediate 28 (0.0047 mol) in THF (40 ml) was added dropwise. The reaction mixture was stirred for 18 hours and extra LAH (0.0092 mol) was added, then the resulting mixture was stirred for another 1 day. The reaction mixture was subsequently treated with H₂O (1.5 ml), aq. NaOH soln. (15 %, 1.5 ml) and then H₂O (4.5 ml) and the mixture was stirred for 10 min. This mixture was filtered and the filtrate was neutralised with HCl (1N), concentrated and the residue was purified by column chromatography (silica gel, eluent: DCM/CH₃OH 95/5). The product fractions were collected and concentrated, yielding 0.5 g (25 %) of intermediate 29.

Example A8

a) Preparation of 1-octanol, 8-(4-chloro-2-nitrophenoxy)-, acetate (ester) (intermediate 30)

A mixture of 4-chloro-6-nitro-phenol (0.0205 mol), fine molecular sieves (3.5 g), DMA, p.a. (50 ml) and K₂CO₃ (0.0238 mol) was stirred for 1 hour, then 8-bromo-1-octanol-acetate (0.0235 mol) was added and the reaction mixture was heated at 50 °C for 16 hours. The mixture was cooled and poured out into ice water, then extracted with toluene (2 times 150 ml). The organic layers were combined, dried (MgSO₄), filtered off and the solvent was evaporated (vac.). The residue was purified by column chromatography over silica gel (eluent: DCM/Hexane 80/20). The product fractions were collected and the solvent was evaporated, yielding 6.2 g (87.9 %) of intermediate 30.

b) Preparation of 1-octanol, 8-(2-amino-4-chlorophenoxy)-, acetate (ester) (intermediate 31)

-77-

A mixture of intermediate 30 (0.018 mol) in THF (100 ml) was hydrogenated with Pt/C 5% (1 g) as a catalyst in the presence of thiophene (1 ml) [H178-005]. After uptake of H₂ (3 equiv.), the reaction mixture was filtered over a small plug of Dicalite and concentrated, yielding 5.6 g of intermediate 31 (used as such in the next reaction step).

c) Preparation of 6-quinazolinol, 4-[[2-[[8-(acetyloxy)octyl]oxy]-5-chlorophenyl]amino]-7-methoxy-, acetate (ester) (intermediate 32)

A mixture of 4-chloro-6-methylcarbonyloxy-7-methoxyquinazoline (0.01 mol) and intermediate 31 (0.01 mol) in 2-propanol (60 ml) was heated at 80 °C for 2 hours and the reaction mixture was cooled, then concentrated. DIPE was added and the mixture was stirred for 2 hours. The solids were collected and then dried, yielding 5.0 g of intermediate 32.

d) Preparation of 6-quinazolinol, 4-[[5-chloro-2-[(8-hydroxyoctyl)oxy]phenyl]amino]-7-methoxy- (intermediate 33)

A mixture of intermediate 32 (0.0094 mol) in methanol (100 ml) was heated at 60 °C. Then a solution of K₂CO₃ (0.019 mol) in H₂O (10 ml) was added dropwise. The organic solvent was evaporated and the aqueous concentrate was treated with acetic acid. The resulting precipitate was filtered off, washed with H₂O and dried (vac.) at 60°C, yielding 3.7 g (88 %) of intermediate 33.

Example A9

a) Preparation of 1-nonanol, 9-(4-chloro-2-nitrophenoxy)-, acetate (ester) (intermediate 34)

A mixture of 4-chloro-6-nitro-phenol (0.02 mol), DMA, p.a. (70 ml) and K₂CO₃ (0.0246 mol) was heated at 50 °C for 1 hour and then 9-bromo,1-nonanol acetate (0.024 mol) was added. The reaction mixture was heated over the weekend and poured into ice (250 ml). The solids were collected by filtration, dissolved in DCM, dried (MgSO₄), filtered and concentrated, yielding 8.6 g of intermediate 34.

b) Preparation of 1-nonanol, 9-(2-amino-4-chlorophenoxy)-, acetate (ester) (intermediate 35)

A mixture of intermediate 34 (0.023 mol) in THF (200 ml) was hydrogenated at 50°C with Pt/C 5% (2 g) as a catalyst in the presence of thiophene (2 ml). After uptake of H₂ (3 equiv.), the catalyst was filtered off and the filtrate was evaporated (vac.), yielding intermediate 35.

c) Preparation of 6-quinazolinol, 4-[[2-[[9-(acetyloxy)nonyl]oxy]-5-chlorophenyl]amino]-7-methoxy-, acetate (ester) (intermediate 36)

-78-

A mixture of 4-chloro-6-methylcarbonyloxy-7-methoxyquinazoline (0.00099 mol) and intermediate 35 (0.0010 mol) in 2-propanol (15 ml) was heated at 80 °C for 1.5 hours and then the reaction mixture was concentrated under a dry N₂-flow. DIPE was added; the solids were collected and then dried. Yielding intermediate 36 (off-white solid) Alternatively a mixture of 4-chloro-6-methylcarbonyloxy-7-methoxyquinazoline (0.051 mol) and intermediate 35 (0.0051 mol) in 2-propanol (40 ml) was heated at 80 °C for 4 hours and then the reaction mixture was concentrated under a dry N₂-flow. DIPE was added; the solids were collected and then dried, yielding 2.38 g (84.3%) of intermediate 36.

d) Preparation of 6-quinazolinol, 4-[[5-chloro-2-[(9-hydroxynonyl)oxy]phenyl]amino]-7-methoxy- (intermediate 37)

K₂CO₃ (0.34 g) was added to a solution of intermediate 36 (0.00437 mol) in methanol (40 ml) and H₂O (8 ml) and after 2 hours the resulting precipitate was filtered, giving solids (I) and filtrate (I). Filtrate (I) was evaporated and then H₂O was added to the residue (pH: 10). Acetic acid was added until pH: 5-6 and the mixture was stirred for 10 min., then the solids were filtered off. These solids and solids (I) were combined in H₂O/CH₃OH (20ml/100ml) and then K₂CO₃ (0.380 g) was added. The reaction mixture was heated at 60 °C for 30 min. and extra K₂CO₃ (0.400 g) was added, then the resulting mixture was stirred overnight. The solvent was evaporated and the residue was treated with H₂O and acetic acid. The resulting solids were filtered off, washed with CH₃OH and dried in a vacuum oven at 60 °C, yielding 1.7 g of intermediate 37.

Example A10

a) Preparation of benzoic acid, 5-hydroxy-2-nitro-4-(phenylmethoxy)- (intermediate 38) KOH (75 g) was added to H₂O (175 ml), stirred at RT. 4-benzyloxy-5-methoxy-2-nitro-benzoic acid (0.031 mol;) was added portionwise and the suspension was heated for 12 hours at 75 °C. The reaction mixture was filtered and the filtrate was acidified with HCl (conc.). The resulting precipitate was filtered off, stirred in DIPE, filtered off and dried, yielding 5.75 g (65%) of intermediate 38.

b) Preparation of benzoic acid, 5-hydroxy-2-nitro-4-(phenylmethoxy)-, methyl ester (intermediate 39)

A mixture of intermediate 38 (0.020 mol) in thionylchloride (50 ml) was stirred and refluxed for 2 hours and the solvent was evaporated under reduced pressure. The residue was quenched with methanol (50 ml) and the mixture was stirred over the

-79-

weekend. The solvent was evaporated and then co-evaporated with toluene, yielding intermediate 39.

c) Preparation of benzoic acid, 5-(acetyloxy)-2-nitro-4-(phenylmethoxy)-, methyl ester (intermediate 40)

A mixture of intermediate 39 (0.020 mol) in acetic anhydride (40 ml) and pyridine (6 ml) was heated to 90 °C and the reaction mixture was stirred for 2 hours. The solvent was evaporated under reduced pressure and the residue was filtered over silica gel (eluent: DCM). The product fractions were collected and the solvent was evaporated, yielding 5.4 g (78 %) of intermediate 40.

d) Preparation of benzoic acid, 5-(acetyloxy)-2-amino-4-(phenylmethoxy)-, methyl ester (intermediate 41)

A mixture of intermediate 40 (0.015 mol) in THF (100 ml) was hydrogenated with Pt/C 5% (2 g) as a catalyst in the presence of thiophene solution (1 ml) [H178-032].

After uptake of H₂ (3 equiv.), the catalyst was filtered off and the filtrate was evaporated, yielding 4.7 g of intermediate 41.

e) Preparation of 4(3*H*)-quinazolinone, 6-hydroxy-7-(phenylmethoxy)- (intermediate 42)

A mixture of intermediate 41 (0.015 mol) and ammonium and formic acid (0.0225 mol) in formamide (50 ml) was heated to 150 °C and the reaction mixture was stirred for 4 hours, then the mixture was allowed to reach RT and poured out into ice-water.

The resulting precipitate was filtered off, washed with H₂O and dried (vac.) at 60°C, yielding 2.9 g (72.5 %) of intermediate 42.

f) Preparation of 4(3*H*)-quinazolinone, 6-(acetyloxy)-7-(phenylmethoxy)- (intermediate 43)

A mixture of intermediate 42 (0.011 mol) in monoacetate 1,1,1-ethenediol (12 ml) and pyridine (2 ml) was heated to 90 °C and the reaction mixture was stirred for 3 hours, then the mixture was poured out into ice-water. The resulting precipitate was

filtered off, washed and dried, yielding 3.3 g of (97 %) of intermediate 43.

g) Preparation of 6-quinazolinol, 4-chloro-7-(phenylmethoxy)-, acetate (ester) (intermediate 44)

A solution of intermediate 43 (0.0032 mol) and N,N-dimethylformamide (cat. quant.) in thionyl chloride (30 ml) was stirred and refluxed for 6 hours and then the solvent was evaporated under reduced pressure and co-evaporated with toluene. The residue was dissolved in DCM and washed with NaHCO₃. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated. The residue was used as such in the next reaction step, yielding 0.6 g (60 %) of intermediate 44.

-80-

h) Preparation of 1-hexanol, 6-(4-bromo-2-nitrophenoxy)-, acetate (ester) (intermediate 44a)

A stirring solution of 4-bromo-2-nitrophenol (0.115 mol) in DMA (250 ml) was heated at 40°C, then K₂CO₃ (0.115 mol) was added and the reaction mixture was stirred for 15 min. 6-bromo- 1-hexanol, acetate (0.115 mol) was added and the mixture was stirred overnight at 40 °C. Extra 6-bromo- 1-hexanol, acetate (4 g) was added, then the resulting mixture was stirred for 2 hours, cooled to RT and stirred overnight. The mixture was filtered off and the filtrate was poured out into ice-water (2000 ml) and then extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated under reduced pressure, yielding 39.6 g of intermediate 44a.

i) Preparation of 1-hexanol, 6-(2-amino-4-bromophenoxy)-, acetate (ester) (intermediate 45)

A mixture of intermediate 44a (0.105 mol) in THF (250 ml) was hydrogenated with Pt/C, 5% (3 g) as a catalyst in the presence of thiophene solution (3 ml). After uptake of H₂ (3 equiv.), the catalyst was filtered off and the filtrate was evaporated under reduced pressure. The residue was dissolved in DIPE and converted into the hydrochloric acid salt (1:1) with HCl (20 ml, 6N in 2-propanol). The resulting precipitate was filtered off, washed and dried, yielding 36.91 g (96 %) of intermediate 45.

j) Preparation of 6-quinazolinol, 4-[[2-[[6-(acetyloxy)hexyl]oxy]-5-bromophenyl]amino]-7-(phenylmethoxy)-, acetate (ester) (intermediate 46)

A mixture of intermediate 44 (0.0031 mol) and intermediate 45 (0.0031 mol) in 2-propanol (50 ml) was heated to 80 °C and the reaction mixture was stirred for 6 hours, then the mixture was allowed to reach RT and stirred overnight. Finally, the solvent was evaporated under reduced pressure, yielding 1.9 g of intermediate 46.

k) Preparation of 6-quinazolinol, 4-[[5-bromo-2-[(6-hydroxyhexyl)oxy]phenyl]amino]-7-(phenylmethoxy)- (intermediate 47)

A solution of intermediate 46 (0.0031 mol) in methanol (25 ml) was heated to 60 °C and a solution of K₂CO₃ (0.0062 mol) in H₂O (2.5 ml) was added, then the reaction mixture was stirred for 18 hours. Extra K₂CO₃ (0.0031 mol) was added and the mixture was stirred for 3 hours at 60 °C. The organic solvent (CH₃OH) was evaporated under reduced pressure and the aqueous concentrate was treated with acetic acid. The resulting precipitate was filtered off, washed and dried (vac.) at 60 °C, yielding 1.4 g (84 %) of intermediate 47.

Example A11

a) Preparation of 6-quinazolinol, 4-[[2-[[6-(acetyloxy)hexyl]oxy]-5-bromophenyl]amino]-, acetate (ester) (intermediate 48)

A mixture of intermediate 45 (0.0045 mol) and 4-chloro-6-quinazolinol acetate (ester) (0.0045 mol) in 2-propanol (50 ml) was heated to 80 °C and the reaction mixture was stirred for 2.5 hours. The solvent was evaporated under reduced pressure and the residue was used as such in the next reaction step, yielding intermediate 48.

b) Preparation of 6-quinazolinol, 4-[[5-bromo-2-[(6-hydroxyhexyl)oxy]phenyl]amino]- (intermediate 49)

A mixture of intermediate 48 (0.0045 mol) and K₂CO₃ (0.0135 mol) in H₂O (2.5 ml) and methanol (25 ml) was stirred at 60 °C for 16 hours and the organic solvent was evaporated, then the aqueous concentrate was extracted with CH₃OH/DCM. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated. The residue was dried (vac.) at 60 °C and used as such in the next reaction step, yielding intermediate 49.

Example A12

a) Preparation of 1-pentanol, 5-[[[(4-bromo-2-nitrophenyl)methyl]amino]- (intermediate 50)

A solution of 4-bromo-2-nitro-benzaldehyde, (0.013 mol), 5-amino-1-pentanol (0.013 mol) and titanium, tetrakis (2-propanolato) (0.014 mol) in EtOH (15 ml) was stirred at RT for 1 hour, then the reaction mixture was heated to 50 °C and stirred for 30 min. The mixture was cooled to RT and NaBH₄ (0.013 mol) was added portionwise. The reaction mixture was stirred overnight and then poured out into ice water (50 ml). The resulting mixture was stirred for 20 min., the formed precipitate was filtered off (giving Filtrate (I)), washed with H₂O and stirred in DCM (to dissolve the product and to remove it from the Ti-salt). The mixture was filtered and then the filtrate was dried (MgSO₄) and filtered, finally the solvent was evaporated. Filtrate (I) was evaporated until EtOH was removed and the aqueous concentrate was extracted 2 times with DCM. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated, yielding 3.8 g (93 %) of intermediate 50.

b) Preparation of carbamic acid, [(4-bromo-2-nitrophenyl)methyl](5-hydroxypentyl)-, 1,1-dimethylethyl ester (intermediate 51)

-82-

A solution of intermediate 50 (0.0032 mol) in DMC (20 ml) was stirred at RT and a solution of dicarbonic acid, bis(1,1-dimethylethyl) ester (0.0032 mol) in DMC (5 ml) was added dropwise. The reaction mixture was stirred for 1 hour at RT and washed 2 times with H₂O. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated, yielding intermediate 51.

c) Preparation of carbamic acid, [5-(acetyloxy)pentyl][(4-bromo-2-nitrophenyl)methyl]-, 1,1-dimethylethyl ester (intermediate 52)

A solution of intermediate 51 (0.0032 mol) and pyridine (0.032 mol) in acetic anhydride (15 ml) was stirred at RT for 16 hours, then the solvent was evaporated under reduced pressure and co-evaporated with toluene. The residue was used as such in the next reaction step, yielding 1.47 g (100 %) of intermediate 52.

d) Preparation of carbamic acid, [5-(acetyloxy)pentyl][(2-amino-4-bromophenyl)methyl]-, 1,1-dimethylethyl ester (intermediate 53)

A mixture of intermediate 52 (0.0033 mol) in THF (50 ml) was hydrogenated with Pt/C 5% (0.5 g) as a catalyst in the presence of thiophene solution (0.5 ml) [H179-007]. After uptake of H₂ (3 equiv.), the catalyst was filtered off and the filtrate was evaporated, yielding intermediate 53.

e) Preparation of carbamic acid, [[2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-4-bromophenyl)methyl][5-(acetyloxy)pentyl]-, 1,1-dimethylethyl ester (intermediate 54)

A mixture of intermediate 53 (0.0028 mol) and 4-chloro-7-methoxy-, acetate 6-quinazolinol (ester) (0.0028 mol) in 2-propanol (50 ml) was heated to 60 °C and the reaction mixture was stirred for 1 hour. The solvent was evaporated under reduced pressure and the residue was used as such in the next reaction step, yielding intermediate 54.

f) Preparation of carbamic acid, [[4-bromo-2-[(6-hydroxy-7-methoxy-4-quinazolinyl)amino]phenyl)methyl](5-hydroxypentyl)-, 1,1-dimethylethyl ester (intermediate 55)

A solution of intermediate 54 (0.0028 mol) in methanol (50 ml) was stirred at RT and a solution of K₂CO₃ (0.0056 mol) in H₂O (5 ml) was added, then the reaction mixture was heated to 60 °C and stirred for 18 hours. The organic solvent was removed and the aqueous concentrate was acidified with acetic acid. The resulting precipitate was filtered off, dissolved in DCM, dried (MgSO₄), filtered and the solvent was evaporated, yielding 1.2 g of intermediate 55.

Example A13

-83-

a) Preparation of 6-heptenoic acid, 7-(4-bromo-2-nitrophenyl)-, (6E)- (intermediate 56)

A mixture of NaH, 60% (0.026 mol, free from mineral oil) in DMSO, dry (15 ml) was heated to 65 °C and the mixture was stirred for 1.5 hour (until the generation of H₂ stopped), then the suspension (dark green) was cooled to 15 °C and a solution of 5-carboxypentyltriphenylphosphonium bromide (0.013 mol) in DMSO (10 ml) was added dropwise. The resulting solution (red) was stirred at RT for 10 min. and a solution of 4-bromo-2-nitro-benzaldehyde (0.013 mol; ; 4-Bromo-2-nitrobenzaldehyde) in DMSO, dry (8 ml) was added rapidly. The solution (dark brown) was stirred for 105 min. and quenched with H₂O/Et₂O (25/75, 100 ml). The Et₂O-layer was removed and the aqueous layer was extracted 2 times with ethylacetate, then acidified (pH: 1-2) with HCl (37 %) and extracted again with ethylacetate. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated, yielding intermediate 56.

b) Preparation of 6-heptenoic acid, 7-(4-bromo-2-nitrophenyl)-, methyl ester, (6E)- (intermediate 57)

A solution of intermediate 56 (0.013 mol) in concentrated HCl (0.20 ml) and methanol (10 ml) was stirred overnight at RT and then the solvent was evaporated under reduced pressure. The residue was dissolved in DCM and washed with a NaHCO₃ soln. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated. The residue was filtered over silica gel (eluent: DCM); the product fractions were collected and the solvent was evaporated, yielding 0.800 g of intermediate 57.

c) Preparation of benzeneheptanoic acid, 2-amino-4-bromo-, methyl ester (intermediate 58)

A mixture of intermediate 57 (0.0023 mol) in THF (50 ml) was hydrogenated with 5% concentrated Pt/C 5% (0.5 g) as a catalyst in the presence of thiophene solution (0.5 ml) [H179-035]. After uptake of H₂ (4 equiv.), the catalyst was filtered off and the filtrate was evaporated, yielding 0.72 g of intermediate 58.

d) Preparation of benzeneheptanoic acid, 2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-4-bromo-, methyl ester (intermediate 59)

A mixture of intermediate 58 (0.0023 mol) and 4-chloro-7-methoxy-6-quinazolinol acetate (ester) (0.0023 mol) in 2-propanol (40 ml) was heated to 80 °C and stirred for 4 hours, then the solvent was evaporated under reduced pressure. The residue was stirred in CH₃OH/DIPE (1/9); the resulting precipitate was filtered off, washed with DIPE and dried (vac.) at 60 °C, yielding 0.55 g of intermediate 59.

-84-

e) Preparation of 6-quinazolinol, 4-[[5-bromo-2-(7-hydroxyheptyl)phenyl]amino]-7-methoxy- (intermediate 60)

A mixture of LAH (0.005 mol) in 2-propanol (20 ml) was stirred at RT. A solution of intermediate 59 (0.001 mol) in 2-propanol (30 ml) was added dropwise, then the reaction mixture was stirred overnight. Ethylacetate (20 ml) was added and the excess of 4-chloro-7-methoxy-6-quinazolinol acetate (ester) was decomposed with a 10 % HCl soln. (5 ml). The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated. The residue was used as such in the next reaction step, yielding intermediate 60.

Example A14

a) Preparation of 1-pentanol, 5-[[[(4-bromo-2-nitrophenyl)methyl]methylamino]- (intermediate 61)

10 A solution of intermediate 50 (0.0047 mol), formaldehyde (0.025 mol) and titanium, tetrakis (2-propanolato) (0.0051 mol) in EtOH (150 ml) was heated to 50 °C and stirred for 1 hour, then NaBH₄ (0.026 mol) was added portionwise at RT. The reaction mixture was stirred overnight and then quenched with water (100 ml). The resulting mixture was stirred for 1 hour; the formed precipitate was filtered off and washed.
15 The organic filtrate was concentrated, then the aqueous concentrate was extracted with DCM and dried. The solvent was evaporated and the residue was filtered over silica gel (eluent: DCM/CH₃OH from 98/2 to 95/5). The product fractions were collected and the solvent was evaporated, yielding 0.5 g of intermediate 61.

b) Preparation of 1-pentanol, 5-[[[(4-bromo-2-nitrophenyl)methyl]methylamino]-, acetate (ester) (intermediate 62)

A solution of intermediate 61 (0.0015 mol) and pyridine (0.015 mol) in acetic anhydride (8 ml) was stirred overnight at RT, then the solvent was evaporated and co-evaporated with toluene, yielding intermediate 62.

c) Preparation of 1-pentanol, 5-[[[(2-amino-4-bromophenyl)methyl]methylamino]-, acetate (ester) (intermediate 63)

A mixture of intermediate 62 (0.0015 mol) in THF (50 ml) was hydrogenated with Pt/C 5% (0.5 g) as a catalyst in the presence of thiophene solution (0.5 ml) [H179-034]. After uptake of H₂ (3 equiv.), the catalyst was filtered off and the filtrate was
25 evaporated, yielding 0.5 g of intermediate 63.

d) Preparation of 6-quinazolinol, 4-[[2-[[[5-(acetyloxy)pentyl]methylamino]methyl]-5-bromophenyl]amino]-7-methoxy-, acetate (ester) (intermediate 64)

-85-

A mixture of intermediate 63 (0.0015 mol) and 4-chloro-7-methoxy-6-quinazolinol acetate (ester) (0.0015 mol) in 2-propanol (30 ml) was heated to 80 °C and the reaction mixture was stirred for 1 day. The solvent was evaporated under reduced pressure and the residue was used as such in the next reaction step, yielding 0.83 g of intermediate 64.

e) Preparation of 6-quinazolinol, 4-[[5-bromo-2-[(5-hydroxypentyl)methylamino]methyl]phenyl]amino]-7-methoxy- (intermediate 65)
A solution of intermediate 64 (0.0015 mol) in methanol (25 ml) was stirred at RT and a solution of K₂CO₃ (0.003 mol) in H₂O (2.5 ml) was added, then the reaction mixture was heated to 60 °C and stirred for 18 hours. The solvent was evaporated and H₂O (20 ml) was added, then the mixture was neutralised with acetic acid and the formed precipitate was filtered off. The filtrate was concentrated under reduced pressure and the concentrate was extracted with DCM, filtered, then dried (MgSO₄) and the mixture was concentrated under reduced pressure, yielding 0.5 g (70 %) of intermediate 65.

15 Example A15

a) Preparation of methanesulfonic acid, trifluoro-, 2-(4-chloro-2-nitrophenyl)ethyl ester (intermediate 66)
A mixture of 2-(4-chloro-2-nitrophenyl)-ethanol (0.01 mol) and 2,6-di-tert-butylpyridine (0.012 mol) in nitromethane (30 ml) was stirred under N₂ at 0°C and a mixture of trifluoromethylsulfonic anhydride (0.011 mol) in nitromethane (10 ml) was added dropwise at 0°C, then the reaction mixture was allowed to reach RT and stirred for 1 hour, yielding intermediate 66.

b) Preparation of 1-butanol, 4-[2-(4-chloro-2-nitrophenyl)ethoxy]-, acetate (ester) (intermediate 67)
A mixture of 1-acetoxy-4-hydroxybutane (0.01 mol) in nitromethane (10 ml) was added dropwise to intermediate 66 and then the reaction mixture was stirred for 1 hour at 65°C. The mixture was cooled and water was added. The layers were separated and the aqueous layer was extracted 2 times with DCM. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated. The residue was purified 2 times by column chromatography over silica gel (eluent 1: DCM; eluent 2: hexane/EtOAc 90/10). The product fractions were collected and the solvent evaporated, yielding 0.800 g (25%) of intermediate 67.

c) Preparation of 1-butanol, 4-[2-(2-amino-4-chlorophenyl)ethoxy]-, acetate (ester) (intermediate 68)

-86-

A mixture intermediate 67 in dioxane (40 ml) was hydrogenated at 40°C with Pt/C (0.300 g) as a catalyst in the presence of thiophene solution (0.3 ml). After uptake of H₂ (3 equiv.), the catalyst was filtered off and the filtrate was evaporated, yielding intermediate 68.

d) Preparation of 6-quinazolinol, 4-[[2-[2-[4-(acetyloxy)butoxy]ethyl]-5-chlorophenyl]amino]-7-methoxy-, acetate (ester) (intermediate 69)

- 5 A mixture 4-chloro-6-acetoxy-7-methoxyquinazoline (0.00040 mol) and intermediate 68 (0.00035 mol) in dioxane (q.s.) was stirred for 3 hours at 80°C and then the solvent was evaporated, yielding intermediate 69.

e) Preparation of 6-quinazolinol, 4-[[5-chloro-2-[2-(4-hydroxybutoxy)ethyl]phenyl]amino]-7-methoxy- (intermediate 70)

- A mixture of intermediate 69 (residue) and K₂CO₃ (0.0144 mol) in H₂O (25 ml) and EtOH (25 ml) was stirred over the weekend at RT, then H₂O (150 ml) was added and the reaction mixture was extracted 3 times with DCM. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated. The residue (0.900 g) was purified by HPLC. The product fractions were collected and the solvent was evaporated, yielding 0.300 g of intermediate 70.
- 10

15 Example A16

a) Preparation of benzaldehyde, 4-chloro-2-nitro-, oxime (intermediate 71)

- A mixture of 4-chloro-2-nitrobenzaldehyde (0.01077 mol) and hydroxylamine hydrochloride (1:1) (0.01184 mol) in pyridine (20ml) was heated on an oil bath for 2 hours at 80 °C and then the solvent was evaporated under reduced pressure. The residue was taken up in CH₃OH/DCM (10/90) and the resulting mixture was extracted with 1N HCl and then washed with a NaHCO₃ solution and water. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated under reduced pressure. The residue was dried (vacuum) at 50°C, yielding 1.75 g (81%) of intermediate 71.
- 20

b) Preparation of benzaldehyde, 4-chloro-2-nitro-, *O*-[8-(acetyloxy)octyl]oxime (intermediate 72)

- K₂CO₃ (0.00887 mol) was added under heavy stirring to a solution of intermediate 71 (0.00887 mol) in DMSO (25ml), then 8-bromooctyl acetate (0.00887 mol) was added and the reaction mixture was stirred for 4 hours at RT. The mixture was heated for 1 hour on an oil bath at 50-60 °C and then extra 8-bromooctyl acetate (0.669 g) and K₂CO₃ (0.369 g) were added. The reaction mixture was stirred for 4 hours at 50-60 °C and cooled. The mixture was poured out into H₂O/NH₄Cl and extracted with EtOAc. The EtOAc-layer was dried (MgSO₄), filtered off and the solvent was evaporated
- 25
- 30

-87-

(vac.). The residual oil (4 g) was purified by column chromatography over silica gel (eluent: DCM/Hexane 70/30, 80/20, 100/0). The pure product fractions were collected and the solvent was evaporated under reduced pressure, yielding 1.34 g of intermediate 72.

c) Preparation of benzaldehyde, 2-amino-4-chloro-, *O*-[8-(acetyloxy)octyl]oxime (intermediate 73)

5 A mixture of intermediate 72 (0.0036 mol) in THF (100 ml) was hydrogenated overnight under H₂ at RT with Pt/C 5% (0.5 g) as a catalyst in the presence of thiophene solution. (1 ml) and then the reaction mixture was heated overnight at 50°C. After uptake of H₂, the mixture was filtered and the filtrate was evaporated. The residue was taken up in THF (100 ml) and the reaction mixture was hydrogenated
10 overnight at 50°C with Pt/C 5% (0.3 g) as a catalyst in the presence of thiophene solution. (0.1 ml). This mixture in THF (100 ml) was hydrogenated further overnight at RT with Pt/C 5% (0.5 g) as a catalyst. After uptake of H₂ (3 equiv.), the catalyst was filtered off and the filtrate was evaporated, yielding intermediate 73.

d) Preparation of benzaldehyde, 2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-4-chloro-, *O*-[8-(acetyloxy)octyl]oxime (intermediate 74)

15 A mixture of 4-chloro-6-acetoxy-7-methoxyquinazoline (0.0021 mol) and intermediate 73 (0.0022 mol) in 2-propanol, p.a. (30 ml) was heated for 1 hour on an oil bath at 80 °C and then the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: DCM/CH₃OH 99.5/0.5 to gradient with CH₃OH). The pure fractions were collected and the solvent was evaporated, yielding 0.300 g of intermediate 74.

e) Preparation of benzaldehyde, 4-chloro-2-[(6-hydroxy-7-methoxy-4-quinazolinyl)amino]-, *O*-(8-hydroxyoctyl)oxime (intermediate 75)

20 A mixture of intermediate 74 (0.00026 mol) in CH₃OH (10 ml) was treated with a mixture of K₂CO₃ (0.0011 mol) in H₂O (1 ml) and then the reaction mixture was stirred overnight at RT. The organic solvent (CH₃OH) was evaporated and the aqueous concentrate was diluted with H₂O (30 ml). The resulting mixture was acidified with acetic acid until pH: 4-5, then the mixture was stirred for 1 hour and
25 filtered. The obtained solid was washed with CH₃OH (5 ml) and dried in a vacuum oven at 60 °C, yielding 0.199 g (78%) of intermediate 75.

Example A17

a) Preparation of 1-octanol, 8-(3-nitrophenoxy)-, acetate (ester) (intermediate 76)

30 A mixture of 3-nitrophenol (0.0144 mol) and K₂CO₃ (0.0144 mol) in 2-propanone (20 ml) was stirred for 2.5 hours at RT and then 8-bromooctyl acetate (0.0144 mol) was

-88-

added The reaction mixture was stirred for 3 hours at RT and was then stirred and refluxed for 18 hours. Extra 8-bromooctyl acetate (0.004 mol) was added and then the resulting mixture was stirred and refluxed for 18 hours. The mixture was cooled to RT, filtered and the filter residue was washed with 2-propanone. The filtrates were combined and concentrated under reduced pressure. The concentrate was then purified by column chromatography (eluent: Hexane/EtOAc 85/15). The product fractions were collected and the solvent was evaporated, yielding intermediate 76.

b) Preparation of 1-octanol, 8-(3-aminophenoxy)-, acetate (ester) (intermediate 77)

A mixture of intermediate 76 (0.0123 mol) in THF (50 ml) was hydrogenated at 50°C with Pt/C 5% (2 g) as a catalyst in the presence of thiophene solution. (1 ml). After uptake of H₂ (3 equiv.), the catalyst was filtered off and the filtrate was evaporated, yielding 3.6 g of intermediate 77.

c) Preparation of 6-quinazolinol, 4-[[3-[[8-(acetyloxy)octyl]oxy]phenyl]amino]-7-methoxy-, acetate (ester) (intermediate 78)

A mixture of intermediate 77 (0.0123 mol) and 4-chloro-6-acetoxy-7-methoxyquinazoline (0.0123 mol) in 2-propanol (50 ml) was heated at 85°C until complete dissolution and then the reaction mixture was cooled to RT. The resulting precipitate was filtered off, washed with DIPE and dried (vacuum), yielding 5.33 g (88%) of intermediate 78.

d) Preparation of 6-quinazolinol, 4-[[3-[(8-hydroxyoctyl)oxy]phenyl]amino]-7-methoxy- (intermediate 79)

A mixture of intermediate 78 (0.00404 mol) and K₂CO₃ (0.00807 mol) in H₂O (8 ml) and CH₃OH (80ml) was heated at 65°C for 18 hours and then the organic solvent was evaporated under reduced pressure. The residue was diluted with H₂O and the resulting mixture was acidified with 1N HCl to pH: 4. The precipitate was filtered off and dried (vacuum), yielding 1.5 g (90%) of intermediate 79.

Example A18

a) Preparation of 2-propenamide, 3-(4-chloro-2-nitrophenyl)-N-(3-hydroxypropyl)-, (intermediate 80)

1,1'-carbonylbis- 1H-imidazole (0.009 mol) was added to a mixture of 4-chloro-2-nitro- cinnamic acid (0.006 mol) in THF (100 ml) at RT and the resulting mixture was stirred for 2 hours at RT, giving Mixture (I). Mixture (I) was added portionwise to a mixture of 3-amino- 1-propanol (0.06 mol) in THF (100 ml) and the reaction mixture was stirred for 2 hours at RT. The solvent was evaporated and the residue was taken up in water (100 ml). The aqueous layer was extracted with DCM (3 times 100 ml), then the organic layers were combined, dried and filtered. The solvent was evaporated

-89-

under reduced pressure (several co-evaporations with toluene were required). The residue was stirred overnight in toluene at RT and the resulting precipitate was filtered off, then purified by column chromatography over silica gel (eluent: DCM/CH₃OH 100/0 to 97/3). The product fractions were collected and the solvent was evaporated, yielding 1g (59 %) of intermediate 80.

b) Preparation of 2-propenamide, *N*-[3-(acetyloxy)propyl]-3-(4-chloro-2-nitrophenyl)- (intermediate 81)

Pyridine (0.035 mol) was added dropwise to a mixture of intermediate 80 (0.0035 mol) in acetic anhydride (20 ml) at RT and then the reaction mixture was stirred for 1 hour at RT. Finally, the solvent was evaporated under reduced pressure, yielding 1.1 g (100 %) of intermediate 81.

c) Preparation of benzenepropanamide, *N*-[3-(acetyloxy)propyl]-2-amino-4-chloro- (intermediate 82)

10 A mixture of intermediate 81 (0.0033 mol) in THF (50 ml) was hydrogenated at RT for 10 days with Pt/C 5% (0.5 g) as a catalyst in the presence of thiophene solution (0.5 ml). After uptake of H₂ (4 equiv.), the catalyst was filtered off and the filtrate was evaporated under reduced pressure. The residue was purified by column chromatography over silica gel (eluent: DCM/CH₃OH 99.5/0.5 to 95/5). The product
15 fractions were collected and the solvent was evaporated under reduced pressure, yielding 0.8 g (81 %) of intermediate 82.

d) Preparation of benzenepropanamide, 2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-*N*-[3-(acetyloxy)propyl]-4-chloro- (intermediate 83)

A mixture of intermediate 82 (0.0027 mol) and 4-chloro-6-acetoxy-7-methoxyquinazoline (0.0027 mol) in 2-propanol (50 ml) was stirred and refluxed for 2 hours and then the solvent was evaporated under reduced pressure. The residue was
20 purified by column chromatography over silica gel (eluent: DCM/CH₃OH 99/1 to 90/10). The product fractions were collected and the solvent was evaporated under reduced pressure, yielding 0.91 g (65 %) of intermediate 83.

e) Preparation of benzenepropanamide, 4-chloro-2-[(6-hydroxy-7-methoxy-4-quinazolinyl)amino]-*N*-(3-hydroxypropyl)- (intermediate 84)

A mixture of intermediate 83 (0.0017 mol) and potassium carbonate (0.0072 mol) in methanol (20 ml) and water (2 ml) was stirred for 1 hour at RT and then the solvent
25 was evaporated under reduced pressure. The residue was taken up in water and the aqueous layer was acidified with acetic acid. The resulting precipitate was filtered off and dried, yielding 0.46 g (63 %) of intermediate 84.

-90-

Example A19

a) Preparation of 6-quinazolinol, 4-chloro-7-methoxy-, acetate (ester) (intermediate 85)

A mixture of 6-(acetyloxy)-7-methoxy-4(1*H*)-quinazolinone (0.23 mol) and DMF (1 ml) in thionyl chloride (500 ml) was stirred and refluxed for 5 hours and then the reaction mixture was cooled to RT. The solvent was evaporated under reduced pressure and then co-evaporated with toluene. The residue was dissolved in DCM and washed with a saturated aqueous NaHCO₃ solution. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated under reduced pressure. The residue was stirred in DIPE and then the resulting precipitate was filtered off, yielding 55.4 g (95%) of intermediate 85.

b) Preparation of 6-quinazolinol, 4-[(4-chloro-2-hydroxyphenyl)amino]-7-methoxy-, 6-acetate monohydrochloric acid (intermediate 86)

A mixture of intermediate 85 (0.00696 mol) and 2-amino-5-chloro-phenol (0.00696 mol) in 2-propanol (100 ml) was heated under stirring for 4 hours at 85°C and the reaction mixture was cooled to RT, then the resulting precipitate was filtered off, yielding intermediate 86, isolated as a monohydrochloric acid.

c) Preparation of 6-quinazolinol, 4-[[4-chloro-2-[(6-hydroxyhexyl)oxy]phenyl]amino]-7-methoxy-, 6-acetate (intermediate 87)

A solution of intermediate 86 (0.00076 mol) in DMA (20 ml) was stirred at RT and sodium hydride (0.00091 mol) was added portionwise, then the mixture was stirred for 30 min. and a solution of 6-bromo-1-hexanol (0.00091 mol) in DMA (2 ml) was added dropwise. The reaction mixture was stirred overnight at RT and an aqueous NH₄Cl solution (1 ml) was added. The reaction mixture was poured out into ice water and the solvent was evaporated. The residue was purified by HPLC and then the product fractions were collected and the solvent was evaporated, yielding 0.030 g of intermediate 87.

Example A20

Preparation of boronic acid, (8,9,10,11,12,13-hexahydro-20-methoxy-4,6-ethanediylidene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclopentadecin-17-yl)- (intermediate 88)

A mixture of compound 6 (0.0006 mol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi-1,3,2-dioxaborolane (0.00066 mol), dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (0.024 g), potassium acetate (0.00092 mol) and [1,1'-biphenyl]-2-ylidicyclohexyl-phosphine (0.024 g) in DMSO (5 ml) was stirred at 80°C for 2 hours, then the reaction mixture was poured out into ice-water

-91-

and was stirred for 1 hour. The resulting precipitate was filtered off and then purified by column chromatography (eluent: DCM/CH₃OH from 98/2 to 90/10). The pure fractions were collected and the solvent was evaporated, yielding 0.080 g (33 %) of intermediate 88.

5

Example A21

a) Preparation of 1-pentanol, 5-[[[4-chloro-5-fluoro-2-nitrophenyl)methyl]amino]-(intermediate 89)

A solution of 4-chloro-5-fluoro-2-nitro- benzaldehyde (0.0098 mol), pentanolamine (0.0098 mol) and tetrakis (2-propanolato) titanium, (0.011 mol) in EtOH (10 ml) was stirred for 1 hour at RT and sodium hydroborate (0.015 mol) was added portionwise, then the reaction mixture was stirred overnight at RT and H₂O was added. The mixture was stirred for 15 min. and the precipitate was filtered off. The filtrate was evaporated and then the residue was dissolved in DCM and washed with H₂O. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated. The residue was purified by column chromatography over silica gel. The product fractions were collected and the solvent was evaporated, yielding 2.3 g (48%) of intermediate 89.

10

15

b) Preparation of carbamic acid, [5-(acetyloxy)pentyl][(4-chloro-5-fluoro-2-nitrophenyl)methyl]-, 1,1-dimethylethyl ester (intermediate 90)

A solution of intermediate 89 (0.0079 mol) in DCM (20 ml) was treated for 30 min. with a solution of tert-butoxycarbonyl anhydride (0.082 mol) in DCM (20 ml) and then the reaction mixture was washed with H₂O (2 x 20 ml). The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated. The residue was dissolved in acetic anhydride (30 ml) and then the solution was treated with pyridine (5 ml) and stirred over the weekend. The solvent was evaporated and co-evaporated with toluene. The residue was purified by column chromatography over silica gel (eluent: DCM). The product fractions were collected and the solvent was evaporated, yielding 1.4 g (40.9%) of intermediate 90.

20

25

c) Preparation of carbamic acid, [5-(acetyloxy)pentyl][(2-amino-4-chloro-5-fluorophenyl)methyl]-, 1,1-dimethylethyl ester (intermediate 91)

A mixture of intermediate 90 (0.0016 mol), Fe (0.009 mol) and NH₄Cl (0.016 mol) in toluene (40 ml), CH₃OH (40 ml) and H₂O (20 ml) was stirred and refluxed for 2 hours, then the reaction mixture was cooled and filtered over dicalite. The filtrate was evaporated and then the residue was diluted with DCM (50 ml) and washed with H₂O. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated, yielding 0.513 g (80%) of intermediate 91.

30

-92-

d) Preparation of carbamic acid, [[2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-4-chloro-5-fluorophenyl]methyl][5-(acetyloxy)pentyl]-, 1,1-dimethylethyl ester (intermediate 92)

A mixture of intermediate 91 (0.000379 mol) and 4-chloro-6-acetoxy-7-methoxyquinazoline (0.000379 mol) in 2-propanol (10 ml) was heated on an oil bath for 3 hours at 80°C and then the solvent was evaporated. The residue was purified by column chromatography over silica gel (gradient eluent: DCM/CH₃OH 100/0 to 99/1). The product fractions were collected and the solvent was evaporated, yielding 0.148 g (63%) of intermediate 92.

e) Preparation of carbamic acid, [[4-chloro-5-fluoro-2-[(6-hydroxy-7-methoxy-4-quinazolinyl)amino]phenyl]methyl](5-hydroxypentyl)-, 1,1-dimethylethyl ester (intermediate 93)

A solution of intermediate 92 (0.000239 mol) in CH₃OH (10 ml) was treated with a solution of K₂CO₃ (0.00051 mol) in H₂O (1 ml). The reaction mixture was stirred overnight, then the mixture was neutralized with acetic acid and the solvent was evaporated. The residue was diluted with DCM and washed with H₂O. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated, yielding 0.120 g (93,7%) of intermediate 93.

Example A22

a) Preparation of 1-butanol, 4-[[2-(4-chloro-2-nitrophenyl)ethyl]amino]- (intermediate 94)

1-amino-4-butanol (0.0300 mol) was added to a stirred suspension of 2-propanol (30 ml) and molecular sieves (8 g) under N₂ at RT and then a mixture of 4-chloro-2-nitrobenzeneacetaldehyde (0.0100 mol) in 2-propanol (10 ml) was added dropwise. The mixture was stirred for 90 min. and sodium cyanotrihydroborate (0.120 mol) was added portionwise (generation of gas). The reaction mixture was stirred overnight and acidified with 6N HCl to pH < 2. A saturated aqueous K₂CO₃ solution was added to pH: 10 and the resulting mixture was filtered over dicalite. The residual fraction was washed with 2-propanol and stirred in hot EtOAc, then this mixture was filtered over dicalite and the filtrate was evaporated under reduced pressure. The residue was taken up in EtOAc and the mixture was extracted with 1N HCl (250 ml), then the layers were separated to give an aqueous layer(*) and an organic layer (1).

(*) Aqueous layer was separated, neutralized with K₂CO₃ and extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated under reduced pressure, yielding intermediate 94. (fraction 1).

-93-

Organic layer (1) was dried (MgSO_4), filtered off and the solvent was evaporated under reduced pressure. The residue was purified by column chromatography over silica gel. The product fractions were collected and the solvent was evaporated yielding intermediate 94. (fraction 2) The product fractions were collected (Yield 100%).

b) Preparation of carbamic acid, [2-(4-chloro-2-nitrophenyl)ethyl](4-hydroxybutyl)-, 1,1-dimethylethyl ester (intermediate 95)

A mixture of intermediate 94 (0.0015 mol) in DCM (10 ml) was stirred at RT and a solution of bis(1,1-dimethylethyl) dicarbonate (0.0015 mol) in DCM (5 ml) was added dropwise, then the reaction mixture was stirred for 1 hour and an extra solution of bis(1,1-dimethylethyl) dicarbonate (0.0015 mol) in DCM (5 ml) was added. The resulting mixture was stirred for 1 hour and washed 2 times with water. The organic layer was separated, dried (MgSO_4), filtered off and the solvent was evaporated, yielding 0.56g of intermediate 95.

c) Preparation of carbamic acid, [4-(acetyloxy)butyl][2-(4-chloro-2-nitrophenyl)ethyl]-, 1,1-dimethylethyl ester (intermediate 96)

A solution of intermediate 95 (0.0015 mol) and pyridine (0.015 mol) in acetic anhydride (10 ml) was stirred for 18 hours at RT, then the solvent was evaporated under reduced pressure and co-evaporated with toluene. The residue was purified over silica gel (eluent: DCM/ CH_3OH 100/0 to 98/2). The product fractions were collected and the solvent was evaporated, yielding 0.2g (33%) of intermediate 96.

d) Preparation of carbamic acid, [4-(acetyloxy)butyl][2-(2-amino-4-chlorophenyl)ethyl]-, 1,1-dimethylethyl ester (intermediate 97)

A mixture of intermediate 96 (0.0005 mol) in THF (40 ml) was hydrogenated with Pt/C 5% (0.1 g) as a catalyst in the presence of thiophene solution. (0.1 ml). After uptake of H_2 (3 equiv.), the catalyst was filtered off and the filtrate was evaporated, yielding (quantitative yield) intermediate 97.

e) Preparation of carbamic acid, [4-(acetyloxy)butyl][2-[2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-4-chlorophenyl]ethyl]-, 1,1-dimethylethyl ester (intermediate 98)

A solution of intermediate 97 (0.0005 mol) and 4-chloro-6-acetoxy-7-methoxyquinazoline (0.0005 mol) in 2-propanol (15 ml) was stirred for 2 hours at 80°C and then the solvent was evaporated under reduced pressure. The crude residue was purified by Flash column chromatography over silica gel (eluent: DCM/ CH_3OH 99.8/0.2 to 96/4). The product fractions were collected and the solvent was evaporated, yielding 0.150 g of intermediate 98.

-94-

f) Preparation of carbamic acid, [2-[4-chloro-2-[(6-hydroxy-7-methoxy-4-quinazoliny]amino]phenyl]ethyl](4-hydroxybutyl)-, 1,1-dimethylethyl ester (intermediate 99)

A mixture of intermediate 98 (0.00025 mol) and K_2CO_3 (0.0005 mol) in methanol (20 ml) and water (2 ml) was heated and stirred for 18 hours at 50°C and then the solvent was evaporated under reduced pressure. The aqueous layer was neutralised with acetic acid and the product was extracted with DCM. The extract was washed with water, dried ($MgSO_4$) and filtered off and then the filtrate was evaporated, yielding 0.130g of intermediate 99.

Example A23

a) Preparation of benzene, 4-chloro-1-(3-chloropropoxy)-2-nitro- (intermediate 100)
Potassium carbonate (0.15 mol) was added portionwise to a mixture of 4-chloro-2-nitrophenol (0.1 mol) in 2-propanone (500 ml) at RT. The mixture was stirred and refluxed for 30 min. 1-bromo-3-chloro- propane (0.11 mol) was added dropwise and then the reaction mixture was stirred and refluxed for 45 min. Extra 1-bromo-3-chloro- propane (0.44 mol) was added, followed by potassium iodide (1 g) and then the reaction mixture was stirred and refluxed overnight. The obtained mixture was filtered and the filter residue was washed with 2-propanone. The filtrate was evaporated under reduced pressure and the residue was purified by column chromatography over silica gel (eluent: hexane/EtOAc 80/20). The product fractions were collected and the solvent was evaporated, yielding 17.30 g (69 %) of intermediate 100.

b) Preparation of ethanol, 2-[[3-(4-chloro-2-nitrophenoxy)propyl][2-(4-morpholinyl)ethyl]amino]- (intermediate 101)
A mixture of 2-(2-morpholinoethylamino)- ethanol (0.0083 mol), intermediate 100 (0.0085 mol) and sodium carbonate (0.016 mol) in acetonitrile (150 ml) was stirred at reflux temperature for 92 hours and the reaction mixture was filtered off. The filtrate was evaporated under reduced pressure and the residue was purified over silica gel (eluent: DCM/(CH_3OH/NH_3) 99/1 to 95/5). The product fractions were collected and the solvent was evaporated under reduced pressure, yielding 1.4 g (44 %) of intermediate 101.

c) Preparation of ethanol, 2-[[3-(4-chloro-2-nitrophenoxy)propyl][2-(4-morpholinyl)ethyl]amino]-, acetate (ester) (intermediate 102)
Pyridine (0.036 mol) was added to a mixture of intermediate 101 (0.0036 mol) in acetic anhydride (25 ml) at RT and then the reaction mixture was stirred for 1 hour at

-95-

RT. Finally, the solvent was evaporated under reduced pressure, yielding 1.6g (100 %) of intermediate 102.

d) Preparation of ethanol, 2-[[3-(2-amino-4-chlorophenoxy)propyl][2-(4-morpholinyl)ethyl]amino]-, acetate (ester) (intermediate 103)

A mixture of intermediate 102 (0.0037 mol) in THF (50 ml) was hydrogenated at 50°C with Pt/C 5% (0.5 g) as a catalyst in the presence of thiophene solution (0.5 ml).

5 After uptake of H₂ (3 equiv.), the catalyst was filtered off and the filtrate was evaporated under reduced pressure. The residue was purified by column chromatography over silica gel (eluent: DCM/CH₃OH 96/4). The product fractions were collected and the solvent was evaporated under reduced pressure, yielding 0.88 g (60 %) of intermediate 103.

e) Preparation of 6-quinazolinol, 4-[[2-[3-[[2-(acetyloxy)ethyl][2-(4-morpholinyl)ethyl]amino]propoxy]-5-chlorophenyl]amino]-7-methoxy-, acetate (ester) (intermediate 104)

10 A mixture of intermediate 103 (0.0021 mol) and 4-chloro-6-acetoxy-7-methoxyquinazoline (0.0021 mol) in 2-propanol (100 ml) was stirred at reflux temperature for 4 hours and DIPEA (0.3 ml) was added, then the reaction mixture was stirred and refluxed for 2 hours. Finally, the solvent was evaporated under reduced pressure, to give residue (I). A mixture of residue (I) (max. 0.0021 mol), tris[α -(1,2-
15 α :4,5- α)-(1E,4E)-1,5-diphenyl-1,4-pentadien-3-one)]di- palladium (=Pd₂(DBA)₃) (0.00013 mol), [1,1'-binaphthalene]-2,2'-diylbis[diphenyl- phosphine (=BINAP) (0.00026 mol) and calcium oxide (0.021 mol) in dioxane (40 ml) was stirred in a sealed tube at 130°C for 16 hours and then the reaction mixture was filtered over dicalite. The filtrate was evaporated under reduced pressure and the residue was
20 purified by column chromatography over silica gel (eluent: DCM/CH₃OH 98/2 to 90/10). The product fractions were collected and the solvent was evaporated under reduced pressure, yielding 0.480 g (37 %) of intermediate 104.

f) Preparation of 6-quinazolinol, 4-[[5-chloro-2-[3-[(2-hydroxyethyl)[2-(4-morpholinyl)ethyl]amino]propoxy]phenyl]amino]-7-methoxy- (intermediate 105)

A mixture of intermediate 104 (0.00073 mol) and potassium carbonate (0.0057 mol) in methanol (20 ml) and water (2 ml) was stirred for 16 hours at RT, then the solvent
25 was evaporated under reduced pressure and the residue was taken up in water. The aqueous layer was washed with DCM, acidified with acetic acid until pH: 7 and extracted with DCM. The organic layer was dried (MgSO₄), filtered off and the solvent was evaporated, yielding 0.250g (64 %) of intermediate 105 .

-96-

Example A24

a) Preparation of benzamide, 4-chloro-*N*-(6-hydroxyhexyl)-2-nitro- (intermediate 106)

1,1'-carbonylbis-1*H*-imidazole (0.01 mol) was added portionwise to a mixture of 4-chloro-2-nitrobenzoic acid (0.01 mol) in DCM (40 ml) at RT and the mixture was stirred for 1 hour at RT, then 6-hydroxyhexylamine (0.01 mol) was added dropwise at RT and the reaction mixture was stirred for 1 hour at RT. The mixture was washed with water (40 ml) and with HCl (1N, 40 ml). The organic layer was separated, dried, filtered off and the solvent was evaporated under reduced pressure, yielding 1.7 g (57 %) of intermediate 106.

b) Preparation of benzamide, *N*-[6-(acetyloxy)hexyl]-4-chloro-2-nitro- (intermediate 107)

Pyridine (0.057 mol) was added dropwise to a mixture of intermediate 106 (0.0057 mol) in acetic anhydride (26.7 ml) at RT and then the reaction mixture was stirred for 1 hour at RT. The solvent was evaporated under reduced pressure and the residue was taken up in water, then the mixture was extracted with toluene. The organic layer was separated, dried, filtered off and the solvent was evaporated under reduced pressure, yielding 1.8 g (92 %) of intermediate 107.

c) Preparation of benzamide, *N*-[6-(acetyloxy)hexyl]-2-amino-4-chloro- (intermediate 108)

A mixture of intermediate 107 (0.0053 mol) in THF (40 ml) was hydrogenated at 50°C with Pt/C (0.5 g) as a catalyst in the presence of thiophene solution (0.5 ml). Then the reaction mixture was filtered off and the filtrate was evaporated under reduced pressure. The residue was hydrogenated again and after uptake of H₂ (3 equiv.), the catalyst was filtered off and the filtrate was evaporated under reduced pressure, yielding 1.56 g (94 %) of intermediate 108.

d) Preparation of benzamide, *N*-[6-(acetyloxy)hexyl]-2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-4-chloro- (intermediate 109)

A mixture of intermediate 108 (0.0042 mol) and 4-chloro-6-acetoxy-7-methoxyquinazoline (0.0042 mol) in 2-propanol (100 ml) was stirred at 50°C for 16 hours and then the solvent was evaporated under reduced pressure. The residue was purified by column chromatography over silica gel (eluent: DCM/CH₃OH from 99/1 to 90/10). The pure fractions were collected and the solvent was evaporated, yielding 1.3g (59%) of intermediate 109.

e) Preparation of benzamide, 4-chloro-*N*-(6-hydroxyhexyl)-2-[(6-hydroxy-7-methoxy-4-quinazolinyl)amino]- (intermediate 110)

-97-

Intermediate 109 (0.0023 mol) and potassium carbonate (0.0046 mol) were stirred in water (2 ml) and methanol (20 ml) at 50°C for 16 hours, then the reaction mixture was concentrated. The residue was acidified with acetic acid and the aqueous layer was extracted with DCM. The organic layer was separated, dried, filtered off and the solvent was evaporated under reduced pressure, yielding 0.99g (97%) of intermediate 110.

Example A25

a) Preparation of carbamic acid, [(4-bromo-2-nitrophenyl)methyl](5-hydroxypentyl)-, phenylmethyl ester (intermediate 111)

Phenylmethyl chloroformate (0.033 mol) was added dropwise to a mixture of intermediate 50 (max. 0.022 mol) and triethylamine (0.04 mol) in DCM (100 ml) at RT and then the reaction mixture was stirred for 1 hour at RT. Water (100 ml) was added and the mixture was stirred for 30 min. at RT. The organic layer was separated, dried, filtered off and the solvent was evaporated under reduced pressure. The residue was purified by column chromatography (eluent: DCM/CH₃OH 99.5/0.5). The product fractions were collected and the solvent was evaporated, yielding 6.8 g (68 %) of intermediate 111.

b) Preparation of carbamic acid, [(2-amino-4-bromophenyl)methyl](5-hydroxypentyl)-, phenylmethyl ester (intermediate 112)

A mixture of intermediate 111 (0.015 mol) in EtOAc (200 ml) was hydrogenated for 40 hours with Pt/C 5% (2 g) as a catalyst in the presence of thiophene solution (3 ml). After uptake of H₂ (3 equiv.), the catalyst was filtered off and the filtrate was evaporated under reduced pressure, yielding 6.3 g of intermediate 112.

c) Preparation of carbamic acid, [[2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-4-bromophenyl)methyl](5-hydroxypentyl)-, phenylmethyl ester .HCl (1:1) (intermediate 113)

A solution of 6-acetoxy-4-chloro-7-methoxyquinazoline (0.015 mol) in 2-propanol (q.s.) was added to a solution of intermediate 112 (0.015 mol) in 2-propanol (q.s.) at 60°C and then the reaction mixture was stirred for 1 hour at 70°C. The solvent was evaporated under reduced pressure and the obtained residue was stirred in hexane. The resulting precipitate was filtered off and dried, yielding 9.35 g (98 %) of intermediate 113, isolated as a hydrochloric salt (1:1).

d) Preparation of carbamic acid, [[2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-4-bromophenyl)methyl](5-chloropentyl)-, phenylmethyl ester (intermediate 114)

-98-

Methanesulfonyl chloride (0.12 mol) was added to a solution of intermediate 113 (0.0125 mol) in 1-methyl-2-pyrrolidinone (50 ml) at RT and the reaction mixture was stirred for 1 hour at 90°C. The mixture was poured out into water (300 ml) and the aqueous layer was extracted with EtOAc (3 x 100 ml). The organic layer was separated, washed with water (2 x 200 ml), dried, filtered off and the solvent was evaporated under reduced pressure. The obtained residue was purified by column chromatography (eluent: DCM/CH₃OH 99.5/0.5 to 94/6). The pure fractions were collected and the solvent was evaporated under reduced pressure, yielding 3.9 g of intermediate 114.

Example A26

a) Preparation of glycine, N-[(4-chloro-2-nitrophenyl)acetyl]-, ethyl ester (intermediate 115)

A slurry of 4-chloro-2-nitro-benzeneacetic acid (0.0051 mol) and 1-hydroxy-1H-benzotriazole (0.0051 mol) in DCM (50 ml) was treated with 1,1'-carbonylbis-1H-imidazole (0.0051 mol), then after 10 min. DIPEA (0.0051 mol) was added, followed by glycine hydrochloride, ethyl ester (0.0051 mol). The reaction mixture was stirred for 2 hours and was washed with water (50 ml), with a Na₂CO₃ solution (30 ml) and with 1N HCl. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated. DIPE (100 ml) was added to the obtained residue and after stirring the resulting solids were collected, yielding 1.1 g of intermediate 115.

b) Preparation of glycine, N-[(2-amino-4-chlorophenyl)acetyl]-, ethyl ester (intermediate 116)

A mixture of intermediate 115 (0.023 mol) in THF (250 ml) was hydrogenated with Pt/C (2.0 g) as a catalyst in the presence of thiophene solution (1 ml). After uptake of H₂ (3 equiv.), the catalyst was filtered off and the filtrate was evaporated. The obtained residue was suspended in DIPE, then the suspension was stirred at boiling temperature, cooled and the desired product was collected, yield 6.2 g of intermediate 116.

c) Preparation of glycine, N-[[2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-4-chlorophenyl]acetyl]-, ethyl ester (intermediate 117)

A mixture of intermediate 85 (0.00050 mol) and intermediate 116 (0.00050 mol) in 2-propanone (5 ml) was stirred for 16 hours in a pressure tube at 80°C (oil bath temperature), then the reaction mixture was filtered and the filter residue was air-dried, yielding 0.165 g of intermediate 117.

-99-

d) Preparation of glycine, N-[[4-chloro-2-[(6-hydroxy-7-methoxy-4-quinazolinyl)amino]phenyl]acetyl]-, ethyl ester (intermediate 118)

A mixture of intermediate 117 (0.0244 mol) in $\text{NH}_3/\text{CH}_3\text{OH}$ (7N) (50 ml) and CH_3OH (100 ml) was stirred overnight at RT and then the solvent was evaporated under reduced pressure and at RT. Finally, the obtained residue was dried (vac.) overnight at 60°C , yielding 8.2 g (75 %) of intermediate 118.

e) Preparation of glycine, N-[[4-chloro-2-[[6-[3-[(1,1-dimethylethoxy)carbonyl]amino]propoxy]-7-methoxy-4-quinazolinyl]amino]phenyl]acetyl]-, ethyl ester (intermediate 119)

- 5 A mixture of intermediate 118 (0.00308 mol) and cesium carbonate (0.0154 mol) in DMA (20 ml) was stirred for 1 hour at RT, then *N*-(3-bromopropyl)carbamate (0.00308 mol) was added and the reaction mixture was stirred for 1 hour at RT. Extra *N*-(3-bromopropyl)carbamate (0.00308 mol) was added and the resulting mixture was stirred overnight. The mixture was filtered and the filter residue was washed with
- 10 DMA. The filtrates were combined and concentrated under reduced pressure. The crude concentrate was purified by column chromatography (eluent: $\text{DCM}/\text{CH}_3\text{OH}$ 100/0 to 95/5). The pure product fractions were collected and the solvent was evaporated under reduced pressure, yielding intermediate 119.

f) Preparation of glycine, N-[[2-[[6-(3-aminopropoxy)-7-methoxy-4-quinazolinyl]amino]-4-chlorophenyl]acetyl]-, ethyl ester (intermediate 120)

- 15 A mixture of intermediate 119 (0.003 mol) in TFA (50 ml) and DCM (50 ml) was stirred for 1 hour at RT, then the reaction mixture was concentrated under reduced pressure and the concentrate was used as such in the next reaction step without further purification, yielding intermediate 120.

g) Preparation of glycine, N-[[2-[[6-(3-aminopropoxy)-7-methoxy-4-quinazolinyl]amino]-4-chlorophenyl]acetyl]- (intermediate 121)

- 20 A mixture of intermediate 120 (0.003 mol) and $\text{LiOH} \cdot \text{H}_2\text{O}$ (0.018 mol) in ethanol (15 ml) and water (1 ml) was stirred for 2 hours and then this reaction mixture was used as such in the next reaction step without further purification, yielding intermediate 121.

Example A27

a) Preparation of 1-pentanol, 5-[[[4-bromo-5-fluoro-2-nitrophenyl)methyl]amino]- (intermediate 122)

- 25 Reaction under N_2 : a mixture of 4-bromo-5-fluoro-2-nitro- benzaldehyde (0.0379 mol) and 5-amino-1-pentanol (0.0379 mol; 97 %) in 1,2-dichloro- ethane (150 ml) was stirred for 20 min. and $\text{NaBH}(\text{OAc})_3$ (0.0417 mol) was added at RT, then the

-100-

reaction mixture was stirred overnight at RT (after 2 hours, extra $\text{NaBH}(\text{OAc})_3$ (q.s.) was added). The mixture was washed with a saturated NaHCO_3 solution. The organic layer was separated, dried (MgSO_4), filtered off and the solvent was evaporated. The residue was purified by column chromatography (eluent: $\text{DCM}/(\text{CH}_3\text{OH}/\text{NH}_3)$ 90/10).

- 5 The pure product fractions were collected and the solvent was evaporated, yielding 10.5 g of intermediate 122.

b) Preparation of carbamic acid, [(4-bromo-5-fluoro-2-nitrophenyl)methyl](5-hydroxypentyl)-, 1,1-dimethylethyl ester (intermediate 123)

- A solution of dicarbonic acid, bis(1,1-dimethylethyl) ester (0.034 mol) in DCM (20 ml) was added dropwise to a mixture of intermediate 122 (0.031 mol) in DCM (200 ml) and then the resulting mixture was reacted for 4 hours. The reaction mixture was washed with water and the organic layer was separated. The organic layer was then dried (MgSO_4), filtered off and the solvent was evaporated. The residue was purified by column chromatography (eluent: $\text{DCM}/\text{CH}_3\text{OH}$ 95/5). The product fractions were collected and then the solvent was evaporated and co-evaporated with toluene, yielding 9.8 g of intermediate 123.

c) Preparation of carbamic acid, [(2-amino-4-bromo-5-fluorophenyl)methyl](5-hydroxypentyl)-, 1,1-dimethylethyl ester (intermediate 124)

- 15 A mixture of intermediate 123 (0.0022 mol) in EtOAc (100 ml) was hydrogenated with Pt/C 5% (0.5 g) as a catalyst in the presence of thiophene solution (q.s.). After uptake of H_2 (3 equiv.), the reaction mixture was filtered over dicalite and the filtrate was evaporated. The residue was purified by column chromatography (gradient eluent: DCM to $\text{DCM}/\text{CH}_3\text{OH}$). The pure product fractions were collected and the solvent was evaporated, yielding 0.623 g of intermediate 124.

d) Preparation of carbamic acid, [[2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-4-bromo-5-fluorophenyl)methyl](5-hydroxypentyl)-, 1,1-dimethylethyl ester (intermediate 125)

- A mixture of intermediate 85 (0.00154 mol) in acetonitrile (3 ml) was warmed to 60°C and then a solution of intermediate 124 (0.00154 mol) in acetonitrile (3 ml) was added dropwise at 60°C . The reaction mixture was stirred for 30 min. at 60°C and then cooled to RT. The solvent was evaporated and the residue was purified by column chromatography over Biotage (gradient eluent: $\text{DCM}/\text{CH}_3\text{OH}$). The pure product fractions were collected and the solvent was evaporated, yielding 0.800 g of intermediate 125.

e) Preparation of carbamic acid, [[4-bromo-5-fluoro-2-[(6-hydroxy-7-methoxy-4-quinazolinyl)amino]phenyl)methyl](5-hydroxypentyl)-, 1,1-dimethylethyl ester (intermediate 126)

-101-

A solution of potassium carbonate (0.00105 mol) in water (5 ml) was added dropwise to a mixture of intermediate 125 (0.00105 mol) in methanol (50 ml) and then the reaction mixture was stirred for 1 hour at RT. The solvent was evaporated and the residue was extracted with DCM. The organic layer was separated, dried (MgSO₄),
5 filtered off and the solvent was evaporated. The residue was purified by column chromatography over Biotage (gradient eluent: DCM/CH₃OH). The pure product fractions were collected and the solvent was evaporated, yielding 0.590 g of intermediate 126.

f) Preparation of 4,6-ethenopyrimido[4,5-b][6,1,12]benzoxadiazacyclopentadecine-13(8H)-carboxylic acid, 17-bromo-16-fluoro-9,10,11,12,14,19-hexahydro-20-methoxy-, 1,1-dimethylethyl ester (intermediate 127)

A mixture of ADDP (0.00166 mol) in THF dry (70 ml) under N₂ was cooled on an ice
10 bath to 5°C, then tributyl-phosphine (0.00166 mol) was added and the mixture was stirred for 5 min. at 5°C. A solution of intermediate 126. (0.00055 mol) in THF dry (10 ml) was slowly added dropwise and the reaction mixture was stirred for 20 min. at 5°C and was then allowed to reach RT. The mixture was stirred for 3 hours at RT and
15 the solvent was evaporated. The residue was purified by column chromatography (gradient eluent: DCM/CH₃OH). The pure product fractions were collected and the solvent was evaporated, yielding intermediate 127.

Example A28

a) Preparation of glycine, N-[[2-[[6-(2-bromoethoxy)-7-methoxy-4-quinazolinyl]amino]-4-chlorophenyl]acetyl]-, ethyl ester (intermediate 128)

A mixture of intermediate 118 (0.0138 mol) and cesium carbonate (0.0690 mol) in
20 DMF (120 ml) was stirred for 30 min. at RT, then 1,2-dibromo-ethane (0.117 mol) was added and the reaction mixture was stirred overnight at RT. The solvent was evaporated under reduced pressure and the residue was co-evaporated with toluene. The obtained residue was stirred in DIPE and the desired product was filtered off, yielding 6.93 g (91 %) of intermediate 128.

b) Preparation of glycine, N-[[4-chloro-2-[[7-methoxy-6-[2-[[2-(4-morpholinyl)ethyl]amino]ethoxy]-4-quinazolinyl]amino]phenyl]acetyl]-, ethyl ester (intermediate 129)

25 A mixture of intermediate 128 (0.00181 mol) and 4-morpholineethanamine (0.00907 mol) in ethanol (20 ml) was heated in a microwave oven for 90 min. at 100°C and then the reaction mixture was purified by RP high-performance liquid chromatography. The product fractions were collected and the solvent was evaporated, yielding 0.39 g (36 %) of intermediate 129.

-102-

c) Preparation of glycine, N-[[4-chloro-2-[[7-methoxy-6-[2-[[2-(4-morpholinyl)ethyl]amino]ethoxy]-4-quinazolinyl]amino]phenyl]acetyl]- (intermediate 130)

A mixture of intermediate 129 (0.00065 mol) and LiOH .H₂O (0.0032 mol) in ethanol (20 ml) and water (2 ml) was stirred for 2 hours at RT and then the solvent was evaporated under reduced pressure, yielding intermediate 130 (used as such in the next reaction step without further purification).

5

Example A29

a) Preparation of benzoic acid, 4-fluoro-, 5-[(4-fluorobenzoyl)amino]-2-nitrophenyl ester (intermediate 131)

A solution of 5-amino-2-nitro- phenol (0.032 mol) and triethylamine (0.065 mol) in DCM (100 ml) was stirred at RT, then a solution of 4-fluoro- benzoyl chloride (0.065 mol) in DCM (10 ml) was added dropwise and the reaction mixture was stirred for 1 day at RT. The mixture was washed 2 times with 1N HCl and once with water. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated. The obtained residue was stirred in ethanol/hexane (50/50), filtered off and dried, yielding 5.25 g of intermediate 131.

b) Preparation of benzamide, 4-fluoro-N-(3-hydroxy-4-nitrophenyl)- (intermediate 132)

A mixture of intermediate 131 (0.0132 mol) in methanol (80 ml) was stirred at RT and then a solution of NaOCH₃ 30% in methanol (0.0132 mol) in methanol (10 ml) was added. The reaction mixture was stirred for 30 minutes and the organic solvent was evaporated under reduced pressure. The obtained concentrate was stirred in 1N HCl, then the resulting precipitate was filtered off, washed with water and dried (vacuum) at 60°C, yielding 3.3 g of intermediate 132 .

c) Preparation of benzamide, N-[3-[[6-(acetyloxy)hexyl]oxy]-4-nitrophenyl]-4-fluoro- (intermediate 133)

A mixture of intermediate 132 (0.011 mol) and potassium carbonate (0.012 mol) in DMA (100 ml) was stirred for 1 hour at 60°C, then 6-bromohexyl acetate (0.012 mol) was added and the reaction mixture was stirred for 18 hours at 60°C. Extra 6-bromohexyl acetate (0.300 g) was added and the mixture was stirred for 5 hours more. The resulting mixture was poured out into ice water and was extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated, yielding 4.6 g of intermediate 133.

d) Preparation of benzamide, N-[3-[[6-(acetyloxy)hexyl]oxy]-4-aminophenyl]-4-fluoro- (intermediate 134)

-103-

A mixture of intermediate 133 (0.0024 mol) in methanol (75 ml) in a closed vessel was hydrogenated overnight at 50°C with Pd/C 10% (1 g) as a catalyst in the presence of thiophene solution (1 ml). After uptake of H₂ (3 equiv.), the catalyst was filtered off, to give Filtrate (I). A mixture of intermediate 133 (0.0024 mol) in THF (75 ml) in a closed vessel was hydrogenated overnight at 50°C with Pd/C 10% (1 g) as a catalyst in the presence of thiophene solution (1 ml). After uptake of H₂ (3 equiv.), the catalyst was filtered off, to give Filtrate (II). Filtrate (I) and Filtrate (II) were combined and evaporated under reduced pressure, yielding 1.85 g of intermediate 134.

e) Preparation of benzamide, N-[3-[[6-(acetyloxy)hexyl]oxy]-4-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]phenyl]-4-fluoro- (intermediate 135)

10 A mixture of intermediate 85 (0.0048 mol) and intermediate 134 (0.0048 mol) in 2-propanol (80 ml) was stirred for 1 hour at 80°C, then the resulting precipitate was filtered off, washed and dried, yielding 1.15 g of intermediate 135.

f) Preparation of benzamide, 4-fluoro-N-[3-[(6-hydroxyhexyl)oxy]-4-[(6-hydroxy-7-methoxy-4-quinazolinyl)amino]phenyl]- (intermediate 136)

A solution of intermediate 135 (0.0019 mol) and potassium carbonate (0.0038 mol) in water (4 ml) and methanol (40 ml) was stirred for 1 hour at 50°C, then a mixture of potassium carbonate (0.26 g) in water (2 ml) was added and the reaction mixture was stirred for 4 hours. The solvent was evaporated under reduced pressure and the obtained residue was dissolved in acetic acid (100 %). After stirring for 30 min., the solvent was evaporated and the crude residue was stirred in DCM. The formed precipitate was filtered off, washed and then stirred in water. The obtained precipitate was filtered off, washed and dried (vacuum) at 60°C, yielding 0.650 g (66 %) of intermediate 136.

Example A30

a) Preparation of alanine, N-[(4-chloro-2-nitrophenyl)acetyl]-2-methyl-, ethyl ester (intermediate 137)

A mixture of 4-chloro-2-nitro- benzeneacetic acid (0.00456 mol), 1-hydroxy-1H-benzotriazole (0.00456 mol), 1,1'-carbonylbis-1H-imidazole (0.00456 mol) and DIPEA (0.00456 mol) in DCM (20 ml) was stirred at RT for 15 min., then 2-methyl-alanine, ethyl ester (0.00456 mol) was added and the reaction mixture was stirred over the weekend at RT. The mixture was washed 2 times with a saturated potassium carbonate solution, 2 times with 1N HCl and once with water. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated, yielding intermediate 137.

-104-

b) Preparation of alanine, N-[(2-amino-4-chlorophenyl)acetyl]-2-methyl-, ethyl ester (intermediate 138)

A mixture of intermediate 137 (0.00456 mol) in ethanol (25 ml) and THF (25 ml) was hydrogenated at 50°C with Pt/C 5% (0.5 g) as a catalyst in the presence of thiophene solution (0.3 ml). After uptake of H₂ (3 equiv.), the catalyst was filtered off and the filtrate was evaporated, yielding 0.65 g of intermediate 138.

c) Preparation of alanine, N-[[2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-4-chlorophenyl]acetyl]-2-methyl-, ethyl ester (intermediate 139)

- 5 A mixture of intermediate 138 (0.0022 mol) and intermediate 85 (0.0022 mol) in acetonitrile (25 ml) was heated to 80°C and stirred for 2 hours. The solvent was evaporated and the residue was purified by flash column chromatography (eluent: DCM/CH₃OH 100/0, 95/5), then the product fractions were collected and the solvent was evaporated, yielding 0.68 g of intermediate 139.

d) Preparation of alanine, N-[[4-chloro-2-[[6-hydroxy-7-methoxy-4-quinazolinyl]amino]phenyl]acetyl]-2-methyl-, ethyl ester (intermediate 140)

- 10 A mixture of intermediate 139 (0.0013 mol) in CH₃OH/NH₃ (7N) (10 ml) and methanol (10 ml) was stirred for 18 hours at RT and then the solvent was evaporated under reduced pressure, yielding 0.600 g of intermediate 140.

e) Preparation of alanine, N-[[4-chloro-2-[[6-[2-[[1,1-dimethylethoxy)carbonyl]amino]ethoxy]-7-methoxy-4-quinazolinyl]amino]phenyl]acetyl]-2-methyl-, ethyl ester (intermediate 141)

- 15 A mixture of intermediate 140 (0.0013 mol) and cesium carbonate (0.0063 mol) in DMA (20 ml) was stirred for 45 min. at RT, then (2-bromoethyl)- carbamic acid 1,1-dimethylethyl ester (0.0014 mol) was added and the mixture was stirred for 4 hours. Extra (2-bromoethyl)- carbamic acid 1,1-dimethylethyl ester (0.0014 mol) was added and the reaction mixture was stirred overnight at RT. The resulting precipitate was filtered off and the filtrate was evaporated, yielding intermediate 141 (quantitative yield, used as such in the next reaction step).

f) Preparation of alanine, N-[[2-[[6-(2-aminoethoxy)-7-methoxy-4-quinazolinyl]amino]-4-chlorophenyl]acetyl]-2-methyl-, ethyl ester (intermediate 142)

- 20 A solution of intermediate 141 (0.0013 mol) in TFA (15 ml) and DCM (15 ml) was stirred for 1 hour at RT and then the solvent was evaporated, yielding 0.670 g of intermediate 142.

g) Preparation of alanine, N-[[2-[[6-(2-aminoethoxy)-7-methoxy-4-quinazolinyl]amino]-4-chlorophenyl]acetyl]-2-methyl- (intermediate 143)

-105-

A mixture of intermediate 142 (0.0013 mol) and LiOH . H₂O (0.0039 mol) in ethanol (20 ml) and water (1 ml) was heated and stirred for 1 hour at 40°C, then extra LiOH . H₂O (0.01192 mol) was added and the reaction mixture was stirred for 3 hours.

5 Again LiOH . H₂O (0.00477 mol) was added and the resulting mixture was stirred for 1 hour at 40°C. Finally, the solvent was evaporated under reduced pressure, yielding intermediate 143 (Quantitative Yield).

Example A31

a) Preparation of beta-alanine, N-[(2-amino-4-chlorophenyl)methyl]-, methyl ester (intermediate 144)

10 A mixture of β-Alanine, methyl ester, hydrochloride (0.020 mol), 4-chloro-2-nitro-benzaldehyde (0.010 mol) and potassium fluoride (0.019 mol) in methanol (100 ml) was hydrogenated at 50°C for 24 hours with Pt/C (1 g, slurry in THF) as a catalyst in the presence of thiophene solution (1 ml, 4% in 2-propanol). After uptake of H₂ (4 equiv.), the reaction mixture was filtered over dicalite, then the filter residue was washed with DCM and the solvent was evaporated. The obtained residue was
15 dissolved in methanol and was used as such in the next reaction step, yielding intermediate 144.

b) Preparation of beta-alanine, N-[(2-amino-4-chlorophenyl)methyl]-N-[(1,1-dimethylethoxy)carbonyl]-, methyl ester (intermediate 145)

Tert-butyl dicarbonate (0.060 mol) was added to intermediate 144 (0.020 mol) and the reaction mixture was stirred for 1 hour, then NH₃/CH₃OH was added and the mixture was stirred for 1 hour. The solvent was evaporated and the dry residue was filtered
20 over silica gel with DCM as eluent. The filter residue was purified by RP high-performance liquid chromatography, then the product fractions were collected and the solvent was evaporated, yielding 1.206 g (36 %) of intermediate 145.

c) Preparation of beta-alanine, N-[[2-[[6-(acetyloxy)-7-methoxy-4-quinazoliny]amino]-4-chlorophenyl)methyl]-N-[(1,1-dimethylethoxy)carbonyl]-, methyl ester (intermediate 146)

A solution of intermediate 145 (0.0035 mol) in acetonitrile (40 ml) was heated to 80°C, then intermediate 85 (0.0035 mol) was added and the reaction mixture was
25 stirred for 4 hours at 80°C. Finally, the solvent was evaporated to dryness, yielding intermediate 146 (used as such in the next reaction step).

d) Preparation of beta-alanine, N-[[4-chloro-2-[(6-hydroxy-7-methoxy-4-quinazoliny]amino)phenyl)methyl]-N-[(1,1-dimethylethoxy)carbonyl]-, methyl ester (intermediate 147)

-106-

NH₃/CH₃OH (7N) (0.035 mol) was added to a solution of intermediate 146 (0.0035 mol) in methanol (5 ml) and the reaction mixture was stirred for 30 min. at RT. The solvent was evaporated and the obtained residue was dissolved in methanol. DIPE was added and the resulting precipitate was filtered off and dried, yielding 0.9125g. A second crop could be obtained by adding heptane to the filtrate, yielding 0.8794 g (total: 93 %) of intermediate 147.

e) Preparation of beta-alanine, N-[[4-chloro-2-[[6-[3-[(1,1-dimethylethoxy)carbonyl]amino]propoxy]-7-methoxy-4-quinazolinyl]amino]phenyl]methyl]-N-[(1,1-dimethylethoxy)carbonyl]-, methyl ester (intermediate 148)

cesium carbonate (0.00775 mol) was added to a solution of intermediate 147 (0.00155 mol) in DMF dry (15 ml) and the mixture was stirred for 15 min. at RT, then (3-bromopropyl)- carbamic acid 1,1-dimethylethyl ester (0.00155 mol) was added and the reaction mixture was stirred overnight at RT. The solvent was evaporated and the residue was dissolved in DCM. This solution was slowly filtered and the filter residue was washed with DCM, yielding intermediate 148 (used as such in the next reaction step).

f) Preparation of beta-alanine, N-[[2-[[6-(3-aminopropoxy)-7-methoxy-4-quinazolinyl]amino]-4-chlorophenyl]methyl]- hydrochloric acid salt (1:1) (intermediate 149)

Water (4 ml) was added to a solution of intermediate 148 (0.00155 mol) in dioxane (10 ml), then hydrochloric acid (4 ml, 36-38%) was added and the mixture was stirred until the generation of gas stopped. Extra hydrochloric acid (2 ml, 36-38%) was added and the reaction mixture was stirred in a closed vessel for 8 hours. Finally, the solvent was evaporated, yielding intermediate 149 (used as such in the next reaction step).

Example A32

a) Preparation of (R)proline, 1-[(4-chloro-2-nitrophenyl)methyl]-, 1,1-dimethylethyl ester (intermediate 150)

Tetrakis (2-propanolato) titanium (0.010 mol) was added to a solution of D-Proline, 1,1-dimethylethyl ester, hydrochloride (0.010 mol) and 4-chloro-2-nitrobenzaldehyde (0.010 mol) in DCM (30 ml), then the mixture was stirred for 1 hour at RT and NaBH(OAc)₃ (0.011 mol) was added. The reaction mixture was stirred for 2 hours at RT and then water was added. The mixture was filtered over a P2 glass filter and washed with DCM, then the organic layer was separated and the aqueous layer was extracted 2 times with DCM. The organic layers were combined, dried, filtered

-107-

off and the solvent was evaporated, yielding intermediate 150 (used as such in the next reaction step).

b) Preparation of (R) proline, 1-[(2-amino-4-chlorophenyl)methyl]-, 1,1-dimethylethyl ester (intermediate 151)

A mixture of intermediate 150 (0.01 mol) in ethanol (100 ml) and THF (50 ml) was hydrogenated with Pt/C 5% (1 g) as a catalyst in the presence of thiophene solution (1 ml: 4% in DIPE). After uptake of H₂ (3 equiv.), the catalyst was filtered off and the filtrate was evaporated. The residue was purified by RP high-performance liquid chromatography, then the product fractions were collected and the organic solvent was evaporated. The obtained concentrate was filtered and then the filter residue was washed with water and dried in an oven, to give 1.0453 g (34 %) of intermediate 151.

10 The filter was washed with DCM and the aqueous layer was extracted with DCM. The organic layer was then dried and filtered over potassium carbonate, to give 0.0480 g intermediate 151.

c) Preparation of (R)proline, 1-[[2-[[6-(acetyloxy)-7-methoxy-4-quinazolinyl]amino]-4-chlorophenyl]methyl]-, 1,1-dimethylethyl ester (intermediate 152)

Intermediate 85 (0.001 mol) was added to a solution of intermediate 151 (0.001 mol) in 2-propanol (q.s.) and then the reaction mixture was stirred for 2 hours and the solvent was evaporated. Extra Intermediate 85 (0.0185 g) was added to the previous prepared reaction mixture, then the mixture was stirred for one more hour and the solvent was evaporated, yielding intermediate 152.

15

d) Preparation of (R)proline, 1-[[4-chloro-2-[[6-(hydroxy-7-methoxy-4-quinazolinyl)amino]phenyl]methyl]-, 1,1-dimethylethyl ester (intermediate 153)

NH₃/CH₃OH (10 ml) was added to intermediate 152 (0.001 mol) and the reaction mixture was shaken for 1 hour and then the solvent was evaporated, yielding intermediate 153 (used as such in the next reaction step).

20

e) Preparation of (R)proline, 1-[[4-chloro-2-[[6-(3-cyanopropoxy)-7-methoxy-4-quinazolinyl]amino]phenyl]methyl]-, 1,1-dimethylethyl ester (intermediate 154)

A mixture of intermediate 153 (0.0005 mol), 4-bromo-butanenitrile (0.04 ml) and cesium carbonate (0.815 g) was stirred overnight at RT and then the reaction mixture was stirred for 30 min. at 50°C. Extra 4-bromo-butanenitrile (0.009 ml) was added and the mixture was stirred for 4 hours at RT and for another 15 min. at 50°C. The solvent was evaporated and the residue was dissolved in DCM. This solution was filtered over dicalite and the filtrate was evaporated, yielding intermediate 154.

25

-108-

f) Preparation of (R)proline, 1-[[2-[[6-(4-aminobutoxy)-7-methoxy-4-quinazoliny]]amino]-4-chlorophenyl]methyl]-, 1,1-dimethylethyl ester (intermediate 155)

A mixture of intermediate 154 (0.0005 mol) in CH₃OH/NH₃ (40 ml) was hydrogenated at 14°C with raney nickel (cat. quant.) as a catalyst in the presence of thiophene solution (0.1 ml). After uptake of H₂ (2 equiv.), the catalyst was filtered off and the filtrate was evaporated, yielding intermediate 155.

g) Preparation of (R)proline, 1-[[2-[[6-(4-aminobutoxy)-7-methoxy-4-quinazoliny]]amino]-4-chlorophenyl]methyl]-.TFA (1:1) (intermediate 156)

- 5 A solution of intermediate 155 (residue) in TFA/DCM/TIS (90/8/2) (5 ml) was stirred for 7-8 hours, then the solvent was evaporated and the obtained residue was dried overnight in an oven, yielding intermediate 156, isolated as a trifluoro acetic acid salt.

Example A33

a) Preparation of (S)proline, 1-[(4-chloro-5-fluoro-2-nitrophenyl)methyl]-, 1,1-dimethylethyl ester (intermediate 157)

- 10 A solution of L-Proline, 1,1-dimethylethyl ester (0.010 mol) and 4-chloro-5-fluoro-2-nitro- benzaldehyde (0.010 mol) in DCM (30 ml) was cooled to 0°C and tetrakis (2-propanolato) titanium (0.010 mol) was added, then the mixture was stirred for 1 hour at RT and NaBH(OAc)₃ (0.011 mol) was added. The reaction mixture was stirred for 3 hours at RT and extra tetrakis (2-propanolato) titanium (0.001 mol) and
- 15 NaBH(OAc)₃ (0.001 mol) were added. The resulting mixture was stirred for 6 hours at RT. Water was added and the mixture was filtered. The organic layer was separated, dried, filtered off and the solvent was evaporated, yielding intermediate 157 (used as such in the next reaction step).

b) Preparation of (S) proline, 1-[(2-amino-4-chloro-5-fluorophenyl)methyl]-, 1,1-dimethylethyl ester (intermediate 158)

- A mixture of intermediate 157 (0.009 mol) in EtOAc (150 ml) was hydrogenated with
- 20 Pt/C 5% (1 g) as a catalyst in the presence of thiophene solution (1 ml: 4% in DIPE). After uptake of H₂ (3 equiv.), the catalyst was filtered off and the filtrate was evaporated. The residue was purified by RP high-performance liquid chromatography, then the product fractions were collected and the organic solvent was evaporated. The obtained precipitate was filtered off, washed with water and dried to give 1.1286 g (34
- 25 %) of intermediate 158.

c) Preparation of (S)proline, 1-[[2-[[6-(acetyloxy)-7-methoxy-4-quinazoliny]]amino]-4-chloro-5-fluorophenyl]methyl]-, 1,1-dimethylethyl ester (intermediate 159)

-109-

Intermediate 85 (0.001 mol) was added to a solution of intermediate 158 (0.001 mol) in 2-propanol (q.s.) and then the reaction mixture was stirred for 2 hours and the solvent was evaporated, yielding intermediate 159 (used as such in the next reaction step).

d) Preparation of (S)proline, 1-[[4-chloro-5-fluoro-2-[(6-hydroxy-7-methoxy-4-quinazoliny)]amino]phenyl]methyl]-, 1,1-dimethylethyl ester (intermediate 160)

5 $\text{NH}_3/\text{CH}_3\text{OH}$ (10 ml) was added to intermediate 159 (0.001 mol) and the reaction mixture was shaken for 1 hour and then the solvent was evaporated, yielding intermediate 160 (used as such in the next reaction step).

e) Preparation of (S)proline, 1-[[4-chloro-2-[[6-[3-[(1,1-dimethylethoxy)carbonyl]amino]propoxy]-7-methoxy-4-quinazoliny)]amino]-5-fluorophenyl]methyl]-, 1,1-dimethylethyl ester (intermediate 161)

A mixture of intermediate 160 (0.0005 mol), (3-bromopropyl)- carbamic acid, 1,1-dimethylethyl ester (0.12326 ml) and cesium carbonate (0.815 g) was stirred
10 overnight at RT and then the reaction mixture was stirred for 30 min. at 50°C. Extra (3-bromopropyl)- carbamic acid, 1,1-dimethylethyl ester (0.013 g) was added and the mixture was stirred for 4 hours at RT and for another 15 min. at 50°C. The solvent was evaporated and the residue was dissolved in DCM. This solution was filtered over dicalite and the filtrate was evaporated, yielding intermediate 161.

f) Preparation of (S)proline, 1-[[2-[[6-(3-aminopropoxy)-7-methoxy-4-quinazoliny)]amino]-4-chloro-5-fluorophenyl]methyl]-
.TFA salt (intermediate 162)

15 A solution of intermediate 161 (residue) in TFA/DCM/TIS (90/8/2) (25 ml) was stirred overnight, then the solvent was evaporated and the obtained residue was dried overnight in an oven at 80°C, yielding intermediate 162, isolated as trifluoroacetic acid salt.

20 Example A34

a) Preparation of 5-hexenamide, N-[(2-amino-4-chlorophenyl)methyl]- (intermediate 163)

A mixture of 5-hexenoic acid (0.0075 mol) and PL-DCC resin (0.015 mol; Polymer Laboratories: 3417) in DCM (100 ml) was stirred for 15 min. at RT, then 2-amino-4-chloro- benzenemethanamine (0.01125 mol) was added and the resulting mixture was stirred for 3 hours. After addition of methylisocyanate polystyrene (0.01125 mol;

25 Novabiochem: 01-64-0169), the reaction mixture was stirred for 4 hours, filtered and then the solvent was evaporated, yielding 1.43 g (76 %) of intermediate 163.

-110-

b) Preparation of 5-hexenamide, N-[[4-chloro-2-[(6-hydroxy-7-methoxy-4-quinazolinyl)amino]phenyl]methyl]- (intermediate 164)

A solution of intermediate 163 (0.0057 mol) and intermediate 85 (0.0052 mol) in 2-propanol (20 ml) was stirred for 5 hours at 60°C and then the mixture was cooled. 7N NH₃ in methanol (20 ml) was added and the reaction mixture was stirred for 2 hours at RT. Finally, the solvent was evaporated, yielding 1.5 g of intermediate 164.

c) Preparation of 5-hexenamide, N-[[4-chloro-2-[[7-methoxy-6-(4-pentenyl)oxy]-4-quinazolinyl]amino]phenyl]methyl]- (intermediate 165)

- 5 A mixture of intermediate 164 (0.0018 mol) and cesium carbonate (0.0090 mol) in DMF (20 ml) was stirred for 15 min., then 5-bromo- 1-pentene (0.0021 mol) was added and the reaction mixture was stirred overnight at RT. Water and DCM were added and the layers were separated. The organic layer was washed with a 10 % citric acid solution and with brine, then it was dried (MgSO₄), filtered and the solvent was
- 10 evaporated. The residue (0.694 g) was then purified by RP high-performance liquid chromatography. The product fractions were collected and the solvent was evaporated, yielding 0.270 g of intermediate 165.

Example A35

a) Preparation of (S)carbamic acid, [1-[[[(4-chloro-2-nitrophenyl)methyl]amino]carbonyl]-3-butenyl]-, 1,1-dimethylethyl ester (intermediate 166)

- 15 1-[bis(dimethylamino)methylene]- 1H-benzotriazolium, hexafluorophosphate(1-), 3-oxide (0.0056 mol) was slowly added to a solution of 2-[[[(1,1-dimethylethoxy)carbonyl]amino]- 4-pentenoic acid (0.0046 mol), 4-chloro-2-nitro-benzenemethanamine (0.0056 mol), 1-hydroxy- 1H-benzotriazole (0.0056 mol) and DIPEA (0.93 ml) in DMF (25 ml) and then the reaction mixture was stirred for 3
- 20 hours at RT. The mixture was diluted with EtOAc (200 ml) and then washed with a 10 % aq. citric acid solution (50 ml), with water (50 ml), with an aqueous NaHCO₃ solution (50 ml) and with brine (50 ml). The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated, yielding 2.00 g (100 %) of intermediate 166.

b) Preparation of (S)carbamic acid, [1-[[[(2-amino-4-chlorophenyl)methyl]amino]carbonyl]-3-butenyl]-, 1,1-dimethylethyl ester (intermediate 167)

- 25 A mixture of intermediate 166 (0.003 mol) and tin(II) chloride dihydrate (0.015 mol) in ethanol (50 ml) was stirred for 90 min. at 60°C, then the reaction mixture was poured out into water and extracted 3 times with toluene. The organic layer was

-111-

separated, dried (MgSO_4), filtered off and the solvent was evaporated (vacuum), yielding intermediate 167.

c) Preparation of (S) carbamic acid, [1-[[[4-chloro-2-[(6-hydroxy-7-methoxy-4-quinazoliny]amino]phenyl]methyl]amino]carbonyl]-3-butenyl]-, 1,1-dimethylethyl ester (intermediate 168)

A solution of intermediate 167 (0.003 mol) and intermediate 85 (0.0025 mol) in 2-propanol (150 ml) was stirred overnight at 55°C and then the mixture was cooled. 7N NH_3 in methanol (50 ml) was added and the reaction mixture was stirred for 2 hours at RT. Finally, the solvent was evaporated, yielding 2.17 g of intermediate 168.

d) Preparation of (S) carbamic acid, [1-[[[4-chloro-2-[(7-methoxy-6-(4-pentenyl)oxy)-4-quinazoliny]amino]phenyl]methyl]amino]carbonyl]-3-butenyl]-, 1,1-dimethylethyl ester (intermediate 169)

A solution of intermediate 168 (0.0015 mol) and cesium carbonate (0.0075 mol) in DMF (30 ml) was stirred for 15 min. at RT, then 5-bromo-1-pentene (0.0018 mol) was added and the reaction mixture was stirred for 24 hours at RT. Water and DCM were added and the layers were separated. The organic layer was washed with a 10 % citric acid solution and with brine, then it was dried (MgSO_4), filtered and the solvent was evaporated, yielding intermediate 169.

Example A36

a) Preparation of 1-pentanol, 5-[2-(2-amino-4-chlorophenyl)ethoxy]-, acetate (ester) (intermediate 170)

15 Reaction (I): A mixture of 2,6-bis(1,1-dimethylethyl)-pyridine (0.012 mol) and 4-chloro-2-nitro-benzeneethanol (0.01 mol) in 1,2-dichloro-ethane (30 ml) was stirred under N_2 and at 0°C, then a mixture of triflic anhydride (0.011 mol) in 1,2-dichloro-ethane (10 ml) was added dropwise at 0°C and the reaction mixture was stirred for 1 hour at RT, to give Mixture (I). Reaction (II): A solution of 1,5-pentanediol, monoacetate (0.011 mol) in 1,2-dichloro-ethane (10 ml) was added dropwise to
20 Mixture (I) and the resulting mixture was stirred for 1 hour at 65°C. After cooling, water was added and the mixture was partitioned between ethanol/DCM. The organic layer was separated, dried, filtered off and the solvent was evaporated. The obtained residue was purified by flash column chromatography (eluent: DCM/ CH_3OH 100/0, 98/2). The product fractions were collected and the solvent was evaporated, to give
25 Mixture (II). A mixture of Mixture (II) (0.0133 mol) in THF (50 ml) was hydrogenated 2 times at 50°C with Pt/C (1 g) as a catalyst in the presence of thiophene solution (1 ml). After uptake of H_2 (3 equiv.), the catalyst was filtered off and the filtrate was evaporated. The obtained residue was purified over silica gel on a

-112-

glass filter (eluent: Hexane/EtOAc 80/20, 70/30). The product fractions were collected and the solvent was evaporated, yielding 1.5 g of intermediate 170.

b) Preparation of 6-quinazolinol, 4-[[2-[2-[[5-(acetyloxy)pentyl]oxy]ethyl]-5-chlorophenyl]amino]-7-methoxy-, acetate (ester) (intermediate 171)

A mixture of intermediate 170 (0.005 mol) and intermediate 85 (0.005 mol) in dioxane (20 ml) was reacted for 16 hours at 80°C and then the solvent was evaporated, yielding intermediate 171.

c) Preparation of 6-quinazolinol, 4-[[5-chloro-2-[2-[(5-hydroxypentyl)oxy]ethyl]phenyl]amino]-7-methoxy- (intermediate 172)

A mixture of intermediate 171 (residue) and potassium carbonate (5 g) in water (50 ml) and methanol (50 ml) was stirred overnight at RT, then water was added and the mixture was extracted with DCM. The organic layer was washed 2 times with water, then dried, filtered and the solvent was evaporated. Toluene was added and the solvent was evaporated again, yielding 2 g of intermediate 172.

Example A37

a) Preparation of carbamic acid, [3-(4-chloro-2-nitrophenyl)-2-propynyl]-, 1,1-dimethylethyl ester (intermediate 173)

A mixture of 1-bromo-4-chloro-2-nitro-benzene (0.15 mol), dichlorobis(triphenylphosphine)-palladium (0.0075 mol) and copper (I) iodide (0.0075 mol) in triethylamine (300 ml) was stirred at 50°C and 2-propynyl-carbamic acid, 1,1-dimethylethyl ester (0.375 mol) was added portionwise, then the reaction mixture was stirred for 2 hours at 50°C and the solvent was evaporated. The residue was taken up in water and the mixture was extracted with EtOAc. The organic layer was separated, dried, filtered off and the solvent was evaporated. The residue was purified twice by column chromatography (eluent: Hexane/EtOAc 80/20). The product fractions were collected and the solvent was evaporated. The obtained residue (31.8 g) was stirred in hexane and then the resulting precipitate was filtered off and dried, yielding 31.5 g (67.6 %) of intermediate 173.

b) Preparation of carbamic acid, [3-(2-amino-4-chlorophenyl)propyl]-, 1,1-dimethylethyl ester (intermediate 174)

A mixture of intermediate 173 (0.04 mol) in THF (200 ml) was hydrogenated at 50°C for with Pt/C (3 g) as a catalyst in the presence of thiophene solution (1 ml) (in the meantime, the catalyst was changed 2 times). After uptake of H₂ (6 equiv.), the catalyst was filtered off and the filtrate was evaporated, yielding (66 %) of intermediate 174.

-113-

c) Preparation of carbamic acid, [3-[2-[[6-(acetyloxy)-7-methoxy-4-quinazoliny]amino]-4-chlorophenyl]propyl]-, 1,1-dimethylethyl ester (intermediate 175)

A mixture of intermediate 174 (0.04 mol) and intermediate 85 (0.035 mol) in acetonitrile (100 ml) was reacted for 3 hours at 75°C and then the reaction mixture was cooled. The resulting precipitate was filtered off and dried, yielding 12.2 g (69.6 %) of intermediate 175.

d) Preparation of butanoic acid, 4-[[4-[[5-chloro-2-[3-[(1,1-dimethylethoxy)carbonyl]amino]propyl]phenyl]amino]-7-methoxy-6-quinazolinyloxy]-, ethyl ester (intermediate 176)

5 A mixture of intermediate 175 (0.00020 mol) and potassium carbonate (0.00072 mol) in water (1 ml) and methanol (1 ml) was stirred for 16 hours at RT and then the solvent was evaporated. The residue was taken up in water, then the mixture was neutralised with NaOAc and extracted with DCM. The organic layer was separated, dried, filtered off and the solvent was evaporated, yielding 0.850 g of intermediate
10 176.

e) Preparation of butanoic acid, 4-[[4-[[2-(3-aminopropyl)-5-chlorophenyl]amino]-7-methoxy-6-quinazolinyloxy]- (intermediate 177)

A mixture of intermediate 176 (0.00035 mol) in THF (10 ml) / HCl 36% (2 ml) / water (3 ml) was reacted for 16 hours at RT and then the solvent was evaporated. The obtained residue was stirred in acetonitrile, then the resulting precipitate was filtered off and dried, yielding 0.200 g of intermediate 177.

15

Example A38

a) Preparation of carbamic acid, [3-[4-chloro-2-[(6-hydroxy-7-methoxy-4-quinazoliny]amino]phenyl]propyl]-, 1,1-dimethylethyl ester (intermediate 178)

A mixture of hydrochloric acid salt of intermediate 175 (0.056 mol) and potassium carbonate (25 g) in water (250 ml) and methanol (200 ml) was stirred for 6 hours at
20 RT and then the solvent was evaporated. The residue was taken up in a small amount of water, then NaOAc (25 g) was added and the mixture was extracted with DCM/CH₃OH. The organic layer was separated, dried, filtered off and the solvent was evaporated. The obtained residue was stirred in DIPE and after filtration the filter residue was dried, yielding 23.5 g (91.5 %) of intermediate 178.

b) Preparation of 6-quinazolinol, 4-[[2-(3-aminopropyl)-5-chlorophenyl]amino]-7-methoxy-HCl (1:1) (intermediate 179)

A mixture of intermediate 178 (0.015 mol) in methanol (50 ml) and HCl/2-propanol
25 (10 ml) was stirred for 16 hours at RT and then the solvent was evaporated. The

-114-

obtained residue was stirred in DIPE and after filtration the filter residue was dried, yielding 6.1 g (94.6 %) of intermediate 179, isolated as a hydrochloric acid salt.

c) Preparation of acetamide, N-[3-[4-chloro-2-[(6-hydroxy-7-methoxy-4-quinazolinyl)amino]phenyl]propyl]-2-[(2-hydroxyethyl)amino]- (intermediate 180)

Intermediate 179 (0.02 mol), DMF (100 ml) and DIPEA (0.1 mol) were stirred at 0-10°C, then a mixture of bromo- acetyl chloride (0.05 mol) in DCM (10 ml) was added dropwise and the reaction mixture was stirred for 2 hours at RT, to give mixture (I). A mixture of 2-amino- ethanol (0.2 mol) in DMF (20 ml) was added dropwise to mixture (I) and the resulting mixture was stirred for 5 hours at 60°C. The solvent was evaporated and the obtained residue was purified by RP high-performance liquid-chromatography. The product fractions were collected and the solvent was


10 evaporated, yielding 10.7 g of intermediate 180.

d) Preparation of carbamic acid, [2-[[3-[4-chloro-2-[(6-hydroxy-7-methoxy-4-quinazolinyl)amino]phenyl]propyl]amino]-2-oxoethyl](2-hydroxyethyl)-, 1,1-dimethylethyl ester (intermediate 181)

A mixture of intermediate 180 (0.0043 mol) in DCM (50 ml) and THF (50 ml) was stirred and dicarbonic acid, bis(1,1-dimethylethyl) ester (0.0046 mol) was added, then the reaction mixture was stirred for 2 hours and the solvent was evaporated. The residue was taken up in a small amount of water and the mixture was extracted with DCM. The organic layer was separated and the solvent was evaporated. The obtained residue was taken up in methanol and then CH₃OH/NH₃ was added. The mixture was stirred for 2 hours and the solvent was evaporated. The residue was purified on a glass filter (eluent: DCM/CH₃OH 90/10). The product fractions were collected and the solvent was evaporated, yielding 0.500 g of intermediate 181.

20

Example A39



Preparation of

A mixture of intermediate 179 (0.016 mol) in DMF (80 ml) was stirred and DIPEA (0.040 mol) was added, to give Solution (*). 2-(3,5-Dimethoxy-4-

25 formylphenoxy)ethoxymethyl polystyrene (0.00528 mol, Novabiochem: 01-64-0261) previously washed with DCM, was stirred in DCM (120 ml), then tetrakis (2-propanolato) titanium (0.016 mol) was added and the mixture was stirred. Solution (*) was added and the resulting mixture was stirred for 2 hours. After addition of

-115-

NaBH(OAc)₃ (0.016 mol), the reaction mixture was stirred for 16 hours and was filtered off. The filter residue was washed 2 times with DCM (100 ml)/THF (100 ml), 3 times with successively DCM (200 ml) and methanol (200 ml) and finally 3 times with DCM (200 ml). The washed residue was dried for 16 hours at 50°C and the
5 desired product was collected, yielding 9.46 g (77 %) of intermediate 182.

Example A40

a) Preparation of 1,3-dioxolane-2-methanamine, *N*-[(2-amino-4-chlorophenyl)methyl]-*N*-methyl- (intermediate 183)

A solution of *N*-methyl- 1,3-dioxolane-2-methanamine (0.020 mol) and 4-chloro-2-nitro- benzaldehyde (0.010 mol) in methanol (200 ml) was hydrogenated at 50°C over
10 the weekend with Pt/C (cat. quant., slurry in EtOAc) as a catalyst in the presence of thiophene solution (q.s., 4 % in THF). After uptake of H₂ (4 equiv.), the reaction mixture was filtered over dicalite and the solvent was evaporated. The residue was purified by RP high-performance liquid chromatography. The product fractions were collected and the organic component of the eluent was evaporated. The precipitate
15 was filtered off, to give 0.7879 g (31 %) of intermediate 183.

b) Preparation of 6-quinazolinol, 4-[[5-chloro-2-[[[(1,3-dioxolan-2-ylmethyl)methylamino]methyl]phenyl]amino]-7-methoxy-, acetate (ester) (intermediate 184)

A solution of intermediate 85 (0.00156 mol) and intermediate 183 (0.00156 mol) in acetonitrile (15 ml) was stirred for 3 hours at 80°C and then the reaction mixture was allowed to cool overnight. The mixture was stirred for another hour at 80°C and then 3 drops of glacial acetic acid was added. The resulting mixture was stirred at 80°C
20 and again glacial acetic acid (1 ml) were added. After stirring overnight at 80°C, the mixture was cooled to RT and the obtained precipitate was filtered off. The filtrate was evaporated and the residue was dried in an oven, yielding intermediate 184 (used as such in the next reaction step).

c) Preparation of 6-quinazolinol, 4-[[5-chloro-2-[[[(1,3-dioxolan-2-ylmethyl)methylamino]methyl]phenyl]amino]-7-methoxy- (intermediate 185)

A solution of intermediate 184 (0.00156 mol) in NH₃/CH₃OH (q.s.) was stirred for 1
25 hour at RT and then the reaction mixture was filtered, to give filter residue and filtrate. The filtrate was triturated with acetonitrile and then the desired product was collected, yielding 0.1350 g of intermediate 185.

d) Preparation of carbamic acid, [3-[[4-[[5-chloro-2-[[[(1,3-dioxolan-2-ylmethyl)methylamino]methyl]phenyl]amino]-7-methoxy-6-quinazolinyl]oxy]propyl]-, 1,1-dimethylethyl ester (intermediate 186)

-116-

cesium carbonate (0.00464 mol) was added to a solution of intermediate 185 (0.00093 mol) in DMF (9 ml) and the mixture was stirred for 1 hour at RT. (3-bromopropyl)-carbamic acid, 1,1-dimethylethyl ester (0.00093 mol) was added and the reaction mixture was stirred overnight at RT, then the solvent was evaporated and the residue was dissolved in DCM. This solution was filtered over dicalite and the filtrate was evaporated to dryness, yielding intermediate 186 (used as such in the next reaction step).

Example A41

a) Preparation of benzenemethanamine, 4-chloro-*N*-methyl-2-nitro-*N*-2-propenyl- (intermediate 187)

10 A solution of 4-chloro-2-nitro- benzaldehyde (0.010 mol) and *N*-methyl- 2-propen-1-amine (0.010 mol) in DCM (q.s.) was stirred for 15 hours at RT, then NaBH(OAc)₃ (0.011 mol) was added and the reaction mixture was stirred for 3.5 hours at RT. Extra NaBH(OAc)₃ (0.002 mol) was added and the mixture was filtered over silica gel (eluent: DCM). The second fraction was repurified by column chromatography over silica gel and combined with previously obtained 1st fraction and then the solvent was
15 evaporated, yielding 2.0689 g (86 %) of intermediate 187.

b) Preparation of benzenemethanamine, 2-amino-4-chloro-*N*-methyl-*N*-2-propenyl- (intermediate 188)

Tin (II) chloride dihydrate (0.043 mol) was added to a solution of intermediate 187 (0.0086 mol) in ethanol (40 ml) and after stirring the reaction mixture was heated for 90 min. at 50°C. A saturated aqueous NaHCO₃ solution was added, followed by
20 addition of DCM, then the layers were separated and the separated organic layer was filtered. The aqueous layer was extracted 3 times with DCM and the separated organic layer was filtered again. The filter residue was washed 3 times with DCM and the organic layer of the filtrate was separated, then dried, filtered and the solvent was evaporated, yielding 1.3772 g (76 %) of intermediate 188.

c) Preparation of 6-quinazolinol, 4-[[5-chloro-2-[(methyl-2-propenylamino)methyl]phenyl]amino]-7-methoxy-, acetate (ester) (intermediate 189)

25 Intermediate 85 (0.0016 mol) was added to a solution of intermediate 188 (0.0016 mol) in 2-propanol (20 ml), then reaction mixture was stirred for 3 hours at 80°C and the desired product was collected, yielding intermediate 189.

d) Preparation of 6-quinazolinol, 4-[[5-chloro-2-[(methyl-2-propenylamino)methyl]phenyl]amino]-7-methoxy (intermediate 190)

A solution of intermediate 189 (0.0016 mol) in NH₃/CH₃OH (10 ml) was shaken for 1 hour and then the solvent was evaporated to dryness, yielding intermediate 190.

-117-

e) Preparation of 4-quinazolinamine, 6-(3-butenyloxy)-*N*-[5-chloro-2-[(methyl-2-propenylamino)methyl]phenyl]-7-methoxy- (intermediate 191)

A mixture of intermediate 190 (0.00042 mol), 4-bromo-1-butene (0.0005 mol) and cesium carbonate (q.s.) in DMF (q.s.) was stirred overnight at RT and then the solvent was evaporated. The dry residue was dissolved in DCM and the obtained solution was filtered over dicalite, then the desired product was collected, yielding intermediate 191.

Example A42

a) Preparation of 1*H*-isoindole-1,3(2*H*)-dione, 2-[2-(4-chloro-2-nitrophenyl)ethyl]- (intermediate 192)

A mixture of 4-chloro-1-(2-chloroethyl)-2-nitro- benzene (0.37 mol) and 1*H*-isoindole-1,3(2*H*)-dione, potassium salt (0.55 mol) in DMF (1000 ml) was reacted for 2 hours at 90 °C, then the reaction mixture was cooled and poured out into ice-water. The resulting mixture was stirred for 30 min. at RT and the precipitate was filtered off. The filter residue was dissolved in DCM with MgSO₄ and after filtration the filtrate was evaporated. Yield: 118 g (96 %) of intermediate 192.

b) Preparation of benzeneethanamine, 4-chloro-2-nitro- (intermediate 193)

Hydrazine, monohydrate (2.0 mol) was slowly added dropwise to a mixture of intermediate 192 (0.37 mol) in methanol (1000 ml) and then the reaction mixture was reacted for 6 hours at 55°C. After filtration, the filtrate was evaporated and water was added to the obtained residue. The mixture was extracted 3 times with toluene, then the organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated, yielding 61.5 g of intermediate 193.

c) Preparation of benzeneethanamine, 2-amino-4-chloro- (intermediate 194)

A mixture of intermediate 193 (0.225 mol) in THF (500 ml) was hydrogenated with Pt/C 5% (5 g) as a catalyst in the presence of thiophene solution (5 ml). After uptake of H₂ (3 equiv.), the catalyst was filtered off and the filtrate was evaporated. The obtained residue was dissolved in toluene and a 1N HCl solution (600 ml), then this solution was stirred for 1 hour at 60°C and after cooling sodium hydroxide was added until pH: 9. The organic layer was separated and the aqueous layer was extracted 2 times with toluene. The organic layers were combined, dried (MgSO₄), filtered off and the solvent was evaporated, yielding 30 g of intermediate 194.

d) Preparation of (S) carbamic acid, [2-[[2-(2-amino-4-chlorophenyl)ethyl]amino]-1-methyl-2-oxoethyl]-, 1,1-dimethylethyl ester (intermediate 195)

-118-

A mixture of *N*-[(1,1-dimethylethoxy)carbonyl]-L-Alanine (0.0015 mol) and PL-DCC resin (0.0030 mol; Polymer Laboratories, Part No 3417) in DCM (20 ml) was stirred for 30 min at RT. A mixture of 1-[bis(dimethylamino)methylene]-1*H*-benzotriazolium, hexafluorophosphate(1-), 3-oxide (0.0015 mol) in a small amount of DMF (5 ml) was added. A mixture of intermediate 194 (0.00225 mol) in DCM (2 ml) was added and the reaction mixture was stirred for 5 hours, then methylisocyanate polystyrene (0.00225 mol; NovaBiochem, No 01-64-0169) was added [and additionally, (polystyrylmethyl)trimethylammonium bicarbonate (0.00450 mol; NovaBiochem, No 01-64-0419) was added. After 15 hours, the reaction mixture was filtered and the solvent was evaporated, yielding intermediate 195.

e) Preparation of (S) carbamic acid, [2-[[2-[4-chloro-2-[(6-hydroxy-7-methoxy-4-quinazoliny)amino]phenyl]ethyl]amino]-1-methyl-2-oxoethyl]-, 1,1-dimethylethyl ester (intermediate 196)

A solution of intermediate 195 (0.00110 mol) and intermediate 85 (0.00100 mol) in 2-propanol (20 ml) was stirred for 5 hours at 50°C, then the mixture was cooled and NH₃, 7N in methanol (10 ml) was added. The reaction mixture was stirred for 2 hours at RT and the solvent was evaporated, yielding intermediate 196.

f) Preparation of (S) acetic acid, [[4-[[2-[2-[(2-amino-1-oxopropyl)amino]ethyl]-5-chlorophenyl]amino]-7-methoxy-6-quinazolinyloxy]-.HCl (1:1) (intermediate 197)

Step I 'alkylation with chloroacetate': A solution of intermediate 196 (0.001 mol), chloro- acetic acid, methyl ester (0.002 mol) and potassium carbonate (0.003 mol) in acetonitrile dry (20 ml) was stirred for 3 hours at 75°C, then water (2 ml) and DCM (10 ml) were added and the reaction mixture was stirred for 5 min. at RT. The mixture was filtered through Isolute HM-N cartridges, followed by elution with DCM and then the solvent was evaporated, to give Residue (I). Step II 'deprotection': A solution of Residue (I) in concentrated HCl (2.5 ml), water (2.5 ml) and dioxane (5.0 ml) was stirred for 24 hours at 60°C and then the solvent was evaporated, yielding intermediate 197 isolated as a hydrochloric acid salt (1:1).

Example A43

a) Preparation of carbamic acid, [4-[[[(4-chloro-2-nitrophenyl)methyl]amino]-4-oxobutyl]-, 1,1-dimethylethyl ester (intermediate 198)

N-(ethylcarbonimidoyl)-*N,N*-dimethyl-1,3-propanediamine (0.0049 mol) was added portionwise to a mixture of 4-[[[(1,1-dimethylethoxy)carbonyl]amino]-butanoic acid (0.0049 mol), 4-chloro-2-nitro-benzenemethanamine (0.0041 mol) and DIPEA (0.0049 mol) in DMF (30 ml) at RT and then the reaction mixture was stirred 3 hours at RT. The mixture was diluted with EtOAc (150 ml), washed with a 10 % aqueous

-119-

citric acid solution, with water, with an aqueous NaHCO_3 solution and then with brine. The organic layer was separated, dried, filtered off and the solvent was evaporated, yielding 1.225g of intermediate 198.

b) Preparation of carbamic acid, [4-[[[(2-amino-4-chlorophenyl)methyl]amino]-4-oxobutyl]-, 1,1-dimethylethyl ester (intermediate 199)

A mixture of intermediate 198 (0.003 mol) in THF (25 ml) and methanol (25 ml) was hydrogenated at 50°C with Pt/C 5% (0.5 g) as a catalyst in the presence of thiophene solution (0.5 ml). After uptake of H_2 (3 equiv.), the catalyst was filtered off and the filtrate was evaporated, yielding intermediate 199.

c) Preparation of carbamic acid, [4-[[[4-chloro-2-[(6-hydroxy-7-methoxy-4-quinazolinyl)amino]phenyl]methyl]amino]-4-oxobutyl]-, 1,1-dimethylethyl ester (intermediate 120)

A solution of intermediate 199 (0.0033 mol) and intermediate 85 (0.00275 mol) in 2-propanol (100 ml) was stirred for 3 hours at 50°C and after cooling NH_3 , 7N in methanol (50 ml) was added. The reaction mixture was stirred for 2 hours and then the solvent was evaporated, yielding intermediate 120.

d) Preparation of acetic acid, [[4-[[5-chloro-2-[[[4-[(1,1-dimethylethoxy)carbonyl]amino]-1-oxobutyl]amino]methyl]phenyl]amino]-7-methoxy-6-quinazolinyl]oxy]-, methyl ester (intermediate 121)

A mixture of intermediate 120 (0.001 mol), potassium carbonate (0.003 mol) and chloro- acetic acid, methyl ester (0.003 mol) in acetonitrile (10 ml) was stirred for 3 hours at 75°C, then the reaction mixture was filtered over silica gel and the filter residue was washed with 2-propanone. Finally, the filtrate was evaporated overnight under vacuum, yielding intermediate 121.

e) Preparation of acetic acid, [[4-[[2-[[[(4-amino-1-oxobutyl)amino]methyl]-5-chlorophenyl]amino]-7-methoxy-6-quinazolinyl]oxy]- (intermediate 122)

A solution of intermediate 121 (0.001 mol) in HCl concentrated (3 ml), THF (6 ml) and water (3 ml) was stirred for 24 hours at 60°C and then the solvent was evaporated, yielding intermediate 122.

B. Preparation of the compounds

Example B1

Preparation of 4,6-ethanediylidene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclopentadecin-8(9*H*)-one, 10,11,12,13-tetrahydro-20-methoxy-15-methyl- (compound 1)

A solution of intermediate 5 (0.00008 mol), *N'*-(ethylcarbonimidoyl)-*N,N*-dimethyl-1,3-propanediamine (0.00024 mol) and DMC (5 ml) was stirred at RT and then *N'*-

-120-

(ethylcarbonimidoyl)-*N,N*-dimethyl-1,3-propanediamine, monohydrochloride (0.00008 mol) was added. The reaction mixture was stirred over the weekend at RT. The reaction was completed and the mixture was washed 2 times with H₂O. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated. The residue was purified by high-performance liquid chromatography over RP-18 (Normal Phase). The product fractions were collected, the solvent was evaporated and the residue was dried (vac.) at 65 °C, yielding 0.009g of compound 1.

Example B2

Preparation of 4,6-ethanediylidene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclopentadecine, 15-chloro-8,9,10,11,12,13-hexahydro-20-methoxy- (compound 2)

10 A solution of intermediate 9 (0.0024 mol) and triphenylphosphine (0.0036 mol) in THF, dry (100 ml) was stirred at RT and then a solution of bis(1-methylethyl)diazenedicarboxylate (0.0036 mol) in THF (10 ml) was added dropwise. The reaction mixture was stirred for 6 hours and extra bis(1-methylethyl)diazenedicarboxylate (0.35 ml) in THF (10 ml) was added. The mixture

15 was stirred overnight and concentrated. The residue was purified by column chromatography over silica gel (eluent: DCM/CH₃OH/THF 90/5/5). The product fractions were collected and further purified by RP high-performance liquid chromatography. The product fractions were collected and concentrated. The aqueous concentrate was filtered, and the solid retained washed and dried (vac.) at 65 °C,

20 yielding 0.065 g of compound 2, melting point 255.5-260.2°C.

Example B3

Preparation of 4,6-ethanediylidene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclopentadecine, 17-chloro-8,9,10,11,12,13-hexahydro-20-methoxy- (compound 3)

A solution of intermediate 13 (0.0012 mol) and tributylphosphine (0.0018 mol) in THF (dry)(50 ml) was stirred under N₂ conditions at RT and then a mixture of 1,1'-(azodicarbonyl)bis-piperidine (0.0018 mol) in THF (dry) (10 ml) was added

25 dropwise. The reaction mixture was stirred overnight and extra tributylphosphine (0.30 ml) was added. The mixture was stirred for another 4 hours and the solvent was evaporated. The residue was purified by RP high-performance liquid chromatography and the product fractions were collected and concentrated, the aqueous concentrate

30 was filtered, and the solid retained washed and dried (vac.) at 65 °C, yielding 0.040 g of compound 3, melting point 241.5-242.7°C.

Example B4

Preparation of 4,6-ethanediylidene-8*H*,18*H*-pyrimido[4,5-*b*][6,12,1]benzodioxazacyclotetradecine, 16-chloro-9,10,11,12-tetrahydro-19-methoxy- (compound 4)

A solution of intermediate 17 (0.001 mol) and tributylphosphine (0.0012 mol) in THF (40 ml) was stirred at RT under N₂ and then a solution of 1,1'-(azodicarbonyl)bis-piperidine (0.0012 mol) in THF (10 ml) was added dropwise. The reaction mixture was stirred for 4 h and then an extra amount of tributylphosphine (1 ml) and 1,1'-(azodicarbonyl)bis-piperidine (1g) were added. The resulting mixture was stirred overnight and the solvent was concentrated under reduced pressure. The residue was purified by RP high-performance liquid chromatography. The product fractions were collected and the organic solvent was evaporated. The resulting precipitate was filtered, washed and dried (vacuum) at 65°C, yielding 0.065 g of compound 4, melting point 213.5-221.2°C.

Example B5

Preparation of 4,6-ethanediylidene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclopentadecine, 8,9,10,11,12,13-hexahydro-20-methoxy- (compound 5)

A solution of intermediate 21 (0.0013 mol) and tributylphosphine (0.002 mol) in THF (50 ml) was stirred at RT and then a solution of 1,1'-(azodicarbonyl)bis-piperidine (0.002 mol) in THF (5 ml) was added. After addition, the reaction mixture was stirred for 6 hours and the reaction was completed. The solvent was evaporated and the residue was purified by RP high-performance liquid chromatography. The product fractions were collected and the organic solvent was evaporated. The aqueous concentrate was filtered and the solid retained washed and dried (vac.) at 65 °C, yielding 0.100 g of compound 5, melting point 243.3-251.2°C.

Example B6

Preparation of 4,6-ethanediylidene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclopentadecine, 17-bromo-8,9,10,11,12,13-hexahydro-20-methoxy- (compound 6)

A solution of intermediate 25 (0.00079 mol) and tributylphosphine (0.00316 mol) in THF, dry (50 ml) was stirred at RT under N₂-atm., then a solution of 1,1'-

-122-

(azodicarbonyl)bis-piperidine (0.00316 mol) in THF, dry (10 ml) was added and the reaction mixture was stirred for 12 hours at RT under N₂-atm. The solvent was evaporated, the residue was stirred in DIPE and the mixture was filtered. The filtrate and the residue were combined and purified by RP high-performance liquid chromatography. The product fractions were collected and concentrated, yielding 0.180 g (51 %) of compound 6, melting point 228.6-234.8°C.

Example B7

Preparation of 4,6-ethanediylidene-8*H*,20*H*-pyrimido[4,5-*b*][6,14,1]benzodioxazacyclohexadecine, 18-chloro-9,10,11,12,13,14-hexahydro-21-methoxy- (compound 7)

Tributylphosphine (0.0017 mol) and 1,1'-(azodicarbonyl)bis-piperidine (0.0017 mol) were added at RT to a solution of intermediate 29 (0.0012 mol) in THF (80 ml) and the reaction mixture was stirred for 2 hours. The solvent was evaporated under reduced pressure and the residue was stirred in boiling DIPE/CH₃CN (20ml/5ml). This mixture was filtered, the solid retained was washed with CH₃CN and purified by RP high-performance liquid chromatography. The product fractions were collected and the organic solvent was evaporated. The aqueous concentrate was filtered, and the solid retained washed and dried (vac.) at 65 °C, yielding 0.145 g (30%) of compound 7, melting point 240.6-243.7°C.

Example B8

Preparation of 4,6-ethanediylidene-21*H*-pyrimido[4,5-*b*][6,15,1]benzodioxazacycloheptadecine, 19-chloro-8,9,10,11,12,13,14,15-octahydro-22-methoxy- (compound 8)

A solution of intermediate 33 (0.0045 mol) in THF (200 ml) was stirred at RT and tributylphosphine (0.0047 mol), then 1,1'-(azodicarbonyl)bis-piperidine (0.0047 mol) were added. The reaction mixture was stirred for 4 hours and the solvent was evaporated until 2/3 of the initial volume. The mixture was filtered and the residue washed with a small amount of THF. The filtrate was concentrated and this residue was suspended in H₂O and stirred. The resulting precipitate was collected by filtration, washed with water and treated with boiling 2-propanol. The mixture was cooled and filtered, the solid retained was washed with 2-propanol and DIPE and dried (vac.) at 60°C, yielding 1.4 g (74%) of compound 8, melting point 147.7-151.1 °C.

-123-

Example B9

Preparation of 4,6-ethanediylidene-8*H*,22*H*-pyrimido[4,5-*b*][6,16,1]benzodioxazacyclooctadecine, 20-chloro-9,10,11,12,13,14,15,16-octahydro-23-methoxy- (compound 9)

A solution of intermediate 37 (0.0022 mol) in THF (100 ml) was stirred at RT and tributylphosphine (0.0023 mol), then 1,1'-(azodicarbonyl)bis-piperidine (0.0023 mol) was added. The reaction mixture was stirred for 4 hours and the solvent was evaporated until 2/3 of the initial volume. The precipitate was filtered and washed with a small amount of THF. The filtrate was concentrated and the residue was stirred in H₂O. The resulting precipitate was collected by filtration, washed with water and treated with boiling 2-propanol. The mixture was cooled and filtered, then the solid retained was washed with 2-propanol and with DIPE and dried (vacuum) at 60°C), yielding 0.6 g (63 %) of compound 9, melting point 177.4-183.8 °C.

Example B10

a) Preparation of 4,6-ethanediylidene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclopentadecine-17-carboxylic acid, 8,9,10,11,12,13-hexahydro-20-methoxy-, methyl ester (compound 10)

A mixture of compound 6 (0.0005 mol), Pd(OAc)₂ (0.022 g), 1,3-propanediylbis[diphenyl-phosphine] (0.088 g) and potassiumacetate (0.100 g) in methanol (q.s.; dry) was reacted under CO-gas (30 atm) for 16 hours at 125 °C. The solvent was evaporated. The residue was taken up into water and this mixture was extracted with DCM. The separated organic layer was dried, filtered and the solvent evaporated. The residue was purified by HPLC over X-Terra (gradient elution with eluent: CH₃CN/CH₃OH/NH₄OAc). The product fractions were collected and the solvent was evaporated. The residue was taken up into water, alkalised with K₂CO₃, then extracted with DCM. The separated organic layer was dried, filtered and the solvent evaporated, yielding 0.057 g of compound 10.

b) Preparation of 4,6-ethanediylidene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclopentadecine-17-carboxylic acid, 8,9,10,11,12,13-hexahydro-20-methoxy- (compound 11)

-124-

A mixture of compound 10 (0.0002 mol) in THF (3 ml), methanol (3 ml), NaOH 1N (1 ml) and H₂O (2 ml) was stirred at 50 °C for 3 hours. The solvent was evaporated. Water (2 ml) was added. HCl (1 N, 1 ml) was added and the mixture was stirred for a while. The precipitate was filtered off, washed with water, then filtered off again and stirred in THF, then filtered off and dried, yielding 0.036 g of compound 11.

Example B11

Preparation of pyrrolidine, 1-[(8,9,10,11,12,13-hexahydro-20-methoxy-4,6-ethanediyldiene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclopentadecin-17-yl)carbonyl]- (compound 12)

A mixture of compound 6 (0.0004 mol), Pd(OAc)₂ (0.011 g), 1,3-propanediylbis[diphenyl-phosphine] (0.044 g) and pyrrolidine (0.100 g) in THF (q.s., dry) was reacted under CO-gas (30 atm) for 16 hours at 125 °C. The solvent was evaporated. The residue was taken up into water and this mixture was extracted with DCM. The separated organic layer was dried, filtered and the solvent evaporated. The residue was purified by HPLC over X-Terra (gradient elution with eluent: CH₃CN/CH₃OH/NH₄OAc). The product fractions were collected and the solvent was evaporated. The residue was taken up into water, alkalisied with K₂CO₃, then extracted with DCM. The separated organic layer was dried, filtered and the solvent evaporated, yielding 0.051 g of compound 12.

Example B12

Preparation of 4,6-ethanediylidene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclopentadecine-17-carbonitrile, 8,9,10,11,12,13-hexahydro-20-methoxy- (compound 13)

A mixture of compound 6 (0.0002 mol), tris[μ-[(1,2-η:4,5-η)-(1*E*,4*E*)-1,5-diphenyl-1,4-pentadien-3-one]]di-palladium, (0.011 g), 1,1'-bis(diphenylphosphino)-ferrocene (0.013 g), Zn (0.005 g) and Zn(CN)₂ (0.045 g) in (2-oxo-1-pyrrolidinyl)-methyl (2 ml) was reacted in the microwave for 30 min at 150 °C. Water (4 ml) was added and this mixture was extracted three times with ethylacetate. The combined organic layers were washed with water (2 x), dried, filtered and the solvent was evaporated. The residue was purified by reversed-phase HPLC over X-Terra (gradient elution with eluent: CH₃CN/CH₃OH/NH₄OAc). The product fractions were collected and the solvent was evaporated. The residue was taken up into water, then alkalisied with K₂CO₃. This mixture was extracted with DCM. The separated organic layer was dried, filtered and the solvent evaporated, yielding 0.063 g (81%) of compound 13.

Example B13

Preparation of morpholine, 4-[(8,9,10,11,12,13-hexahydro-20-methoxy-4,6-ethanediyldiene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclopentadecin-17-yl)carbonyl]- (compound 14)

A mixture of compound 6 (0.0002 mol), Pd(OAc)₂ (0.022 g),
1,3-propanediylbis[diphenyl-phosphine] (0.088 g) and morpholine (0.200 g) in THF
5 (q.s., dry) was reacted under CO-gas (30 atm) for 24 hours at 125 °C. The solvent was
evaporated. The residue was taken up into water and this mixture was extracted with
DCM. The separated organic layer was dried, filtered and the solvent evaporated. The
residue was purified by HPLC over X-Terra (gradient elution with eluent:
CH₃CN/CH₃OH/NH₄OAc). The product fractions were collected and the solvent was
10 evaporated. The residue was taken up into water, alkalisied with K₂CO₃, then extracted
with DCM. The separated organic layer was dried, filtered and the solvent evaporated,
yielding 0.005 g of compound 14.

Example B14

a) Preparation of 4,6-ethanediyldiene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclo-
pentadecine, 17-bromo-8,9,10,11,12,13-hexahydro-20-(phenylmethoxy)- (compound 15)

15 A solution of intermediate 47 (0.0026 mol) in THF (140 ml) was stirred at RT,
tributylphosphine (0.0035 mol) was added and then ADDP (0.0035 mol). The
reaction mixture was stirred for 6 hours and extra ADDP (0.0035 mol) and
tributylphosphine (0.0035 mol) were added. The resulting mixture was stirred for 12
hours. The formed precipitate was removed and the solvent was evaporated under
20 reduced pressure. The residue was dissolved in THF (100 ml) with molecular sieves.
Extra ADDP (0.0035 mol) and tributylphosphine (0.0035 mol) were added and the
mixture was stirred for 2 hours. The resulting precipitate was filtered off and the
solvent was evaporated under reduced pressure. The residue was filtered over silica
gel (eluent: DCM/CH₃OH 98/2). the product fractions were collected and the solvent
25 was evaporated, yielding 0.600 g of compound 15.

b) Preparation of 4,6-ethanediyldiene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclo-
pentadecin-20-ol, 17-bromo-8,9,10,11,12,13-hexahydro- (compound 16)

A solution of compound 15 (0.0006 mol) and (methylthio)- benzene (0.006 mol) in
trifluoroacetic acid (6 ml) was stirred for 3 days at RT and then the solvent was
evaporated. The residue was quenched with H₂O and the aqueous layer was extracted

-126-

with DCM. The precipitate between the two layers was filtered off, washed and dried (vacuum) at 60°C, yielding compound 16.

Example B15

Preparation of 4,6-ethanediylidene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclopentadecine, 17-bromo-8,9,10,11,12,13-hexahydro-20-[3-(4-morpholinyl)propoxy]-acetic acid (compound 17).

- 5 A mixture of compound 16 (0.000065 mol) and K₂CO₃ (0.00013 mol) in DMA (2 ml) was stirred at 60 °C for 30 min., then 4-(3-chloro-propyl)-morpholine (0.000065 mol) was added and the reaction mixture was stirred for 1 day at 60 °C. Extra 4-(3-chloro-propyl)-morpholine (0.000065 mol) was added and the mixture was stirred for 1 day. After the starting material was consumed, the mixture was purified by RP high-
- 10 performance liquid chromatography. The product fractions were collected and the organic solvent was evaporated. The aqueous concentrate was extracted with DCM/CH₃OH (98/2) and the organic layer was dried (MgSO₄), filtered, then the solvent was evaporated, yielding 0.004 g of compound 17.

15 Example B16

a) Preparation of 4,6-ethanediylidene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclopentadecine, 17-bromo-8,9,10,11,12,13-hexahydro- (compound 18)

- A solution of intermediate 49 (0.0012 mol) in THF (50 ml) was stirred at RT under N₂ and tributylphosphine (0.0017 mol) was added, then 1,1'-(azodicarbonyl)bis-piperidine (0.0017 mol) was added and the reaction mixture was stirred for 1 hour. The solvent was evaporated until 1/3 of the initial volume and the formed precipitate
- 20 was filtered off, then washed. The filtrate was evaporated and the residue was quenched with H₂O. The mixture was acidified with HCl (1N) and extracted with DCM/CH₃OH (99/1). The organic layer was dried (MgSO₄), filtered and the solvent was evaporated. The residue was purified by Flash column chromatography (eluent: DCM/CH₃OH 99/1). The product fractions were collected and the solvent was
- 25 evaporated. The residue was stirred in boiling 2-propanol, then the resulting precipitate was filtered off, washed with 2-propanol and with DIPE and dried (vacuum) at 60 °C, yielding 0.111 g of compound 18.

-127-

Example B17

a) Preparation of 4,6-ethanediylidenepyrimido[4,5-*b*][6,1,12]benzoxadiazacyclopentadecine-13(8*H*)-carboxylic acid, 17-bromo-9,10,11,12,14,19-hexahydro-20-methoxy-, 1,1-dimethylethyl ester (compound 19)

A solution of intermediate 55 (0.0021 mol) in THF (dry) (120 ml) was stirred at RT and tributylphosphine (0.0032 mol) was added, then 1,1'-(azodicarbonyl)bis-piperidine, (0.0032 mol) was added and the reaction mixture was stirred for 3 hours.

5 The solvent was evaporated until 1/3 of the initial volume. The resulting precipitate was filtered off and washed. The filtrate was evaporated and used as such in the next reaction step. A part of the residue (0.200 g) was purified by RP high-performance liquid chromatography. The product fractions were collected and the organic solvent was evaporated. The aqueous concentrate was extracted with DCM and the organic
10 layer was dried (MgSO₄), filtered off, then the solvent was evaporated, yielding 0.005 g of compound 19.

b) Preparation of 4,6-ethanediylidenepyrimido[4,5-*b*][6,1,12]benzoxadiazacyclopentadecine, 17-bromo-8,9,10,11,12,13,14,19-octahydro-20-methoxy- (compound 20)

A solution of compound 19 (0.00092 mol) in mono(trifluoroacetate) (20 ml) was stirred for 1 hour at RT, then the solvent was evaporated under reduced pressure and co-evaporated with toluene. The residue was stirred in boiling 2-propanol, then the
15 resulting precipitate was filtered off, washed and dried. The filtrate was evaporated and the residue was purified by RP high-performance liquid chromatography. The product fractions were collected and the organic solvent was evaporated. The aqueous concentrate was filtered off, washed and dried (vac.) at 70 °C, yielding 0.040 g (5 %) of compound 20.

20

Example B18

a) Preparation of 4,6-ethanediylidenepyrimido[4,5-*b*][6,1]benzoxaazacyclopentadecine, 17-bromo-8,9,10,11,12,13,14,19-octahydro-20-methoxy- (compound 21)

A solution of intermediate 60 (0.0011 mol) in THF dry (50 ml) was stirred at RT and tributylphosphine (0.0016 mol) was added, then 1,1'-(azodicarbonyl)bis-piperidine (0.0016 mol) was added and the reaction mixture was stirred for 4 hours. The solvent
25 was evaporated until 1/3 of the initial volume. The resulting precipitate was filtered off and washed. The filtrate was evaporated and the residue was purified by RP high-performance liquid chromatography. The product fractions were collected and the

-128-

organic solvent was evaporated. The aqueous concentrate was filtered off, washed with H₂O and dried (vac.) at 65 °C, yielding 0.037g (7.5 %) of compound 21.

Example B19

a) Preparation of 4,6-ethanediylidenepyrimido[4,5-*b*][6,1,12]benzoxadiazacyclopentadecine, 17-bromo-8,9,10,11,12,13,14,19-octahydro-20-methoxy-13-methyl- (compound 22)

- 5 A solution of intermediate 65 (0.0011 mol) in THF (50 ml) was stirred at RT and tributylphosphine (0.0016 mol) was added, then 1,1'-(azodicarbonyl)bis-piperidine (0.0016 mol) was added and the reaction mixture was stirred for 2 hours. The solvent was evaporated until 1/3 of the initial volume. The resulting precipitate was filtered off and washed. The filtrate was evaporated and the residue was purified by RP high-
- 10 performance liquid chromatography. The product fractions were collected and the organic solvent was evaporated. The aqueous concentrate was extracted 2 times with DCM and the organic layer was dried (MgSO₄), then filtered off. The solvent was evaporated and the residue was dried (vac.) at 50°C, yielding 0.004 g (0.8 %) of compound 22.

15

Example B20

Preparation of 4,6-ethanediylidene-13*H*-pyrimido[4,5-*b*][6,11,1]benzodioxazacyclopentadecine, 17-chloro-8,9,10,11,14,19-hexahydro-20-methoxy- (compound 23)

- A mixture of intermediate 70 (0.0007 mol) in THF (50 ml) was stirred until complete dissolution and tributylphosphine (0.0014 mol) was added, then the mixture was stirred and ADDP (0.0014 mol) was added. The reaction mixture was stirred at RT
- 20 and then extra ADDP (q.s.) and tributylphosphine (q.s.) were added. The resulting mixture was stirred at 60°C for 10 hours and again extra ADDP (q.s.) and tributylphosphine (q.s.) were added. The mixture was stirred at 100°C for 16 hours. The solvent was evaporated and the residue was purified by HPLC. The product fractions were collected and the solvent was evaporated, yielding 0.017 g of
- 25 compound 23.

-129-

Example B21

Preparation of 4,6-ethanediylidene-23*H*-pyrimido[4,5-*b*][6,15,1,16]benzodioxadiazacyclononadecine, 21-chloro-8,9,10,11,12,13,14,15-octahydro-24-methoxy- (compound 24)

A solution of intermediate 75 (0.000355 mol) and tributylphosphine (0.000356 mol) in THF (20 ml) and DMF p.a. dried on molecular sieves (5 ml) was treated with ADDP (0.000353 mol) and the reaction mixture was stirred at RT, then extra ADDP (q.s.) and tributylphosphine (q.s.) were added and the reaction mixture was stirred at RT. The solvent was evaporated and the residue was purified by HPLC. the product fractions were collected and the solvent was evaporated, yielding 0.0274 g (17%) of compound 24, melting point 127.2-132.2 °C.

10 Example B22

Preparation of 22*H*-4,6-ethanediylidene-21,17-methenopyrimido[5,4-*d*][1,12,6]dioxazacycloeicosine, 8,9,10,11,12,13,14,15-octahydro-24-methoxy- (compound 25)

A solution of intermediate 79 (0.0012 mol) in THF (75 ml) was stirred at RT and then ADDP (0.0018 mol) and tributylphosphine (0.0018 mol) were added. The reaction mixture was stirred for 3 hours and extra ADDP (0.0018 mol) and tributylphosphine (0.0018 mol) were added. The resulting mixture was stirred for 2 hours and the solvent was evaporated under reduced pressure. The residue was stirred in 2-propanol and filtered off, then the filtrate was evaporated and the residue was purified by HPLC. The product fractions were collected and the solvent was evaporated, yielding 0.0027 g (72%) of compound 25.

20 Example B23

a) Preparation of 4,6-ethenopyrimido[4,5-*b*][6,1,10]benzoxadiazacyclopentadecine-12(13*H*)-one, 17-chloro-8,9,10,11,14,19-hexahydro-20-methoxy- (compound 26)

ADDP (0.00034 mol) was added to a solution of intermediate 84 (0.00023 mol) and tributylphosphine (0.00042 mol) in THF (20 ml) and DMF (20 ml) at RT and the reaction mixture was stirred at RT for 1 hour. Extra ADDP and tributylphosphine were added at RT and then the resulting mixture was stirred for 1 hour at RT. The

-130-

solvent was evaporated off and again extra ADDP and tributylphosphine were added. The mixture was warmed to 100°C and stirred for 18 hours at 100°C, then the solvent was evaporated under reduced pressure and the residue was purified by HPLC. The product fractions were collected and the solvent was evaporated, yielding 0.0094 g (10 %) of compound 26.

b) Preparation of 4,6-ethanediylidenepyrimido[4,5-
b][6,1,12]benzoxadiazacyclopentadecine-14(19*H*)-one, 17-chloro-8,9,10,11,12,13-hexahydro-20-methoxy- (compound 27)

Compound 27 is made in a similar way as compound 26.

Example B24

Preparation of 4,6-ethanediylidene-19*H*-pyrimido[4,5-
b][6,13,1]benzodioxazacyclopentadecine, 17-bromo-8,9,10,11,12,13-hexahydro-20-(2-methoxyethoxy)- (compound 28)

A mixture of compound 16 (0.00023 mol), 1-bromo-2-methoxy-ethane (0.0046 mol) and K₂CO₃ (0.00046 mol) in DMA (10 ml) was stirred for 18 hours at 60°C and then the reaction mixture was quenched with ice-water. The precipitate was filtered off, washed and stirred in boiling 2-propanol. The resulting precipitate was filtered off, washed and dried (vacuum) at 53°C, yielding 0.030 g (74%) of compound 28.

Example B25

a) Preparation of 4,6-ethanediylidenepyrimido[4,5-
b][6,1,12]benzoxadiazacyclopentadecine-13(8*H*)-carboxylic acid, 17-chloro-16-fluoro-9,10,11,12,14,19-hexahydro-20-methoxy-, 1,1-dimethylethyl ester (compound 29)

Tributylphosphine (0.00044 mol) was added to a solution of intermediate 93 (0.00022 mol) and ADDP (0.00044 mol) in THF (30 ml) and then extra ADDP (0.00044 mol) and tributylphosphine (0.00044 mol) were added. The reaction mixture was stirred was stirred over the weekend and the solvent was evaporated. CH₃OH (5 ml) was added and the resulting mixture was stirred, then filtered and the filtrate evaporated. The residue was purified by reversed phase HPLC. The product fractions were collected and the solvent was evaporated, yielding 0.04 g (35.2%) of compound 29.

-131-

b) Preparation of 4,6-ethanediylidenepyrimido[4,5-
b][6,1,12]benzoxadiazacyclopentadecine, 17-chloro-16-fluoro-8,9,10,11,12,13,14,19-
octahydro-20-methoxy- (compound 30)

A solution of compound 29 (0.000077 mol) in CH₃OH(5 ml) was treated with HCl/2-
propanol (6N) (1ml) and then the reaction mixture was stirred overnight at RT. The
solvent was evaporated and the residue was diluted with DCM/NaHCO₃. After
stirring the mixture for 1 hour, the organic layer was separated, dried (MgSO₄),
5 filtered off and the solvent was evaporated, yielding 0.0089 g (27.7%) of compound
30, melting point 265.9-261.3 °C.

Example B26

a) Preparation of 4,6-ethanediylidenepyrimido[4,5-
b][6,1,11]benzoxadiazacyclopentadecine-12(13*H*)-carboxylic acid, 17-chloro-
8,9,10,11,14,19-hexahydro-20-methoxy-, 1,1-dimethylethyl ester (compound 31)

A solution of intermediate 99 (0.00025 mol), ADDP (0.000375 mol) and
10 tributylphosphine (0.000375 mol) in THF (20 ml) was stirred for 4 hours at RT and
then the solvent was evaporated until 1/3 of the initial volume. The resulting
precipitate was filtered off and washed, then the filtrate was evaporated and the
residue was purified by reversed phase HPLC. The product fractions were collected
and the solvent was evaporated, yielding 0.02 g of compound 31.

b) Preparation of 4,6-ethanediylidenepyrimido[4,5-
b][6,1,11]benzoxadiazacyclopentadecine, 17-chloro-8,9,10,11,12,13,14,19-octahydro-20-
methoxy- (compound 32)

15 A solution of compound 31 (0.00004 mol) in TFA (5 ml) was stirred for 4 hours at
RT and then the solvent was removed under N₂ at 40 °C. The residue was purified by
HPLC. The product fractions were collected and the solvent evaporated, yielding
0.0037g (69%) of compound 32.

20 Example B27

Preparation of 4,6-etheno-19*H*-pyrimido[5,4-
k][1,8,5,13]benzodioxadiazacyclopentadecine, 17-chloro-8,9,10,11,12,13-hexahydro-20-
methoxy-10-[2-(4-morpholinyl)ethyl]- (compound 33)

-132-

ADDP (0.00068 mol) and tributylphosphine (0.00085 mol) were added to a solution of intermediate 105 (0.00047 mol) in THF (30 ml) at RT and then the reaction mixture was stirred for 2 hours at RT. The solvent was evaporated under reduced pressure and the residue was purified by column chromatography over silica gel (eluent: DCM/(CH₃OH/NH₃) 99/1 to 80/20). The pure fractions were collected and the solvent was evaporated under reduced pressure. The residue (0.032g) was then purified by HPLC. The product fractions were collected and the solvent was evaporated, yielding 0.0055 g of compound 33.

10 Example B28

a) Preparation of 4,6-ethanediylidene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclopentadecine, 8,9,10,11,12,13-hexahydro-20-methoxy-17-phenyl- (compound 34)

A mixture of intermediate 88 (0.0001 mol), iodo- benzene (0.0002 mol), dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (cat.quant., 5 %), sodium carbonate 2M in water (0.0003 mol) in DMSO (2 ml) was stirred at 80°C for 3 hours, then the reaction mixture was poured out into ice-water and the aqueous layer was extracted with DCM. The organic layer was separated, dried, filtered off and the solvent was evaporated under reduced pressure. The residue was purified by column chromatography over silica gel (eluent: DCM/CH₃OH 98/2). The product fractions were collected and the solvent was evaporated, yielding 0.016 g (36 %) of compound 34.

b) Preparation of benzonitrile, 3-(8,9,10,11,12,13-hexahydro-20-methoxy-4,6-ethanediylidene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclopentadecin-17-yl)- (compound 35)

20 Compound 35 is made on the same way accordingly compound 34.

Example B29

Preparation of 4,6-ethanediylidene-8*H*-pyrimido[4,5-*b*][6,1,13]benzoxadiazacyclohexadecin-15(20*H*)-one, 18-chloro-9,10,11,12,13,14-hexahydro-21-methoxy- (compound 36)

-133-

ADDP (0.0016 mol) was added to a mixture of intermediate 110 (0.0011 mol) and tributylphosphine (0.0020 mol) in THF (50 ml) and the reaction mixture was stirred for 1 hour at RT. The solvent was evaporated under reduced pressure, then the residue was stirred and refluxed in methanol (80 ml) for 1 hour. The resulting precipitate was filtered off and dissolved in DMF (50 ml). The solution was concentrated again under reduced pressure and the residue was stirred in methanol. Finally, the resulting precipitate was filtered off and dried, yielding 0.242 g (52 %) of compound 36.

Example B30

Preparation of 4,6-ethanediylidene-19*H*-pyrimido[4,5-*b*][6,13,1]benzodioxazacyclopentadecine, 16-chloro-8,9,10,11,12,13-hexahydro-20-methoxy- (compound 37)

10 A solution of intermediate 87 (0.00007 mol) in THF (3ml) was stirred at RT and then ADDP (0.0001 mol) and tributylphosphine (0.0001 mol) were added. The reaction mixture was stirred for 18 hours and extra ADDP (0.0001 mol) and tributylphosphine (0.0001 mol) were added. The resulting mixture was stirred for 18 hours and the solvent was evaporated. The residue was purified by HPLC and the product fractions were collected, then the solvent was evaporated and the residue was dried (vacuum) at 50°C, yielding 0.002g of compound 37.

Example B31

Preparation of 4,6-ethanediylidene-8*H*,14*H*-pyrimido[4,5-*b*][6,12,1]benzodioxazacyclohexadecine, 18-chloro-9,10,11,12,15,20-hexahydro-21-methoxy- (compound 38)

A mixture of intermediate 172 (0.0046 mol) in THF (400 ml) was stirred at RT, then tributylphosphine (0.0092 mol) was added, followed by ADDP (0.0092 mol) and the reaction mixture was stirred for 2 hours. The solvent was evaporated and the residue was purified by RP high-performance liquid chromatography (Hypersil) (eluent: (0.5 % NH₄OAc in water)/CH₃CN 90/10). The product fractions were collected and the solvent was evaporated. The obtained residue was taken up in water and then the mixture was alkalinised with K₂CO₃ and extracted with DCM. The organic layer was separated, dried, filtered off and the solvent was evaporated. The residue (1.1 g) stirred in DIPE and the precipitate was filtered off and then dried, yielding 0.976 g of compound 38.

-134-

Compound 39 is made on the same way accordingly compound 38 .

4,6-ethanediylidene-14H-pyrimido[4,5-b][6,9,12,1]benzotrioxaazacyclohexadecine,
18-chloro-8,9,11,12,15,20-hexahydro-21-methoxy- (compound 39)

Example B32

Preparation of 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,11]benzoxadiazacyclohexadecin-11(12H)-one, 18-chloro-9,10,13,14,15,20-hexahydro-21-methoxy- (compound 40)

- 5 A mixture of intermediate 177 (0.00045 mol), PyBOP (0.00135 mol) and triethylamine (0.00135 mol) was reacted for 3 hours at 60°C and the solvent was evaporated. The residue was purified by RP high-performance liquid chromatography, then the product fractions were collected and the solvent was evaporated, yielding 0.008 g of compound 40.

10

Example B33

a) Preparation of 4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecine-10(11H)-carboxylic acid, 19-chloro-8,9,12,13,14,15,16,21-octahydro-22-methoxy-12-oxo-, 1,1-dimethylethyl ester (compound 41)

- THF p.a. (150 ml) and tributyl-phosphine (0.003 mol) were stirred under N₂ at 50°C and ADDP (0.003 mol) was added, then a mixture of intermediate 181 (0.0009 mol) in THF p.a. (15 ml) was added and the reaction mixture was stirred for 2 hours at
15 60°C. Extra tributyl-phosphine (0.003 mol) and ADDP (0.003 mol) were added and the resulting mixture was stirred for 2 hours at 60°C. Finally, the solvent was evaporated, yielding (used as such in the next reaction step) compound 41.

b) Preparation of 4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-22-methoxy- (compound 42)

A mixture of compound 41 (residue) in methanol (50 ml) and 2-propanol/HCl (5 ml) was stirred for 72 hours at RT and then the solvent was evaporated. The residue was

-135-

taken up in water and washed 3 times with DCM. The aqueous layer was alkalisied with K_2CO_3 and extracted with DCM. The crude mixture was then purified on a glass filter (eluent: DCM(CH_3OH/NH_3) 90/10). The product fractions were collected and the solvent was evaporated, yielding 0.322 g of compound 42.

c) Preparation of 4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-10-[[[2-(hydroxyethyl)methylamino]acetyl]-22-methoxy- (compound 43)

- 5 Compound 42 (0.000045 mol), DMA (2 ml) and DIPEA (0.00013 mol) were stirred and bromo-acetyl chloride (0.00011 mol) was added dropwise, then 2-(methylamino)-ethanol (0.00044 mol) was added and the reaction mixture was stirred for 16 hours at 60°C, yielding 0.013 g of compound 43.

Following compounds were made accordingly:

10

Compound No.	Name
44	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-10-[[[2-(hydroxy-1-(hydroxymethyl)ethyl]amino]acetyl]-22-methoxy-
45	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-10-[[2-(hydroxymethyl)-4-morpholinyl]acetyl]-22-methoxy-
46	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-22-methoxy-10-[[[2-(4-pyridinyl)ethyl]amino]acetyl]-
47	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-12(13H)-one, 19-chloro-10-[[[2-(dimethylamino)ethyl]methylamino]acetyl]-8,9,10,11,14,15,16,21-octahydro-22-methoxy-
48	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-22-methoxy-10-[[[2-methoxyethyl]amino]acetyl]-
49	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-22-methoxy-10-[[[3-methoxypropyl]amino]acetyl]-
50	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-22-methoxy-10-(4-morpholinylacetyl)-

-136-

Compound No.	Name
51	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-22-methoxy-10-[(4-methyl-1-piperazinyl)acetyl]-

Example B34

Preparation of 4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-22-methoxy-10-methyl- (compound 52)

- 4-steps reaction procedure Step (I): intermediate 182 (0.0002 mol), DIPEA (0.0008 mol) and DCM (7 ml) was shaken and bromo-acetyl chloride (0.0008 mol) was added, then the reaction mixture was stirred for 3 hours and washed 3 times with DCM, to give Resin (I). Step (II): Resin (I), 2-(methylamino) ethanol (0.0020 mol) and 1-methyl-2-pyrrolidinone (6 ml) was shaken for 6 hours at 60°C, then the reaction mixture was washed [3 times with DMF and 3 times with DCM] x 2, to give Resin 4-(II). Step -(III): Resin (II), triphenyl-phosphine (0.0020 mol), ADDP (0.0020 mol) and 1-methyl-2-pyrrolidinone (8 ml) was shaken for 6 hours at 60°C, then the reaction mixture was washed 3 times with DMF and 3 times with DCM, to give Resin (III). Step (IV): Resin (III) and DCM/TFA/trisopropylsilane (7 ml) was shaken for 16 hours and filtered, then the filter residue was washed and the solvent was evaporated. The obtained residue was purified by RP high-performance liquid chromatography. The desired product fractions were collected and the solvent was evaporated, yielding 0.001 g of compound 52.

Following compounds are made on the same way accordingly compound 52:

Compound No.	Name
53	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacycloheptadecin-12(13H)-one, 19-chloro-10-ethyl-8,9,10,11,14,15,16,21-octahydro-22-methoxy-
54	1,22-ethanediylidene-5H,17H-pyrimido[4,5-b]pyrrolo[2,1-h][6,1,9,12]benzoxatriazacycloheptadecin-14(15H)-one, 7-chloro-10,11,12,13,18,19,19a,20-octahydro-24-methoxy-

-137-

Compound No.	Name
55	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclooctadecin-13(14H)-one, 20-chloro-9,10,11,12,15,16,17,22-octahydro-23-methoxy-
56	14H-4,6-ethanediylidene-9,13-methano-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacycloeicosin-15(16H)-one, 22-chloro-9,10,11,12,17,18,19,24-octahydro-26-methoxy-
57	13H-4,6-ethanediylidene-9,12-ethanopyrimido[4,5-b][6,1,11,14]benzoxatriazacyclononadecin-14(15H)-one, 21-chloro-8,9,10,11,16,17,18,23-octahydro-26-methoxy-
58	14H-4,6-ethanediylidene-10,13-ethano-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacycloeicosin-15(16H)-one, 22-chloro-9,10,11,12,17,18,19,24-octahydro-27-methoxy-

Example B35

Preparation of 4,6-ethenopyrimido[4,5-b][6,1,12]benzoxadiazacyclopentadecine-13(8H)-carboxylic acid, 17-bromo-9,10,11,12,14,19-hexahydro-20-methoxy-, phenylmethyl ester (compound 59)

A mixture of intermediate 114 (0.005 mol) and K_2CO_3 (0.025 mol) in DMA (25 ml) and water (25 ml) was stirred under microwave conditions for 30 min. at 150°C and then the solvent was evaporated under reduced pressure. The obtained residue was stirred in EtOAc and the precipitate was filtered off. The filtrate was evaporated under reduced pressure and the residue was purified by column chromatography (eluent: DCM/ CH_3OH 98/2 to 96/4). The product fractions were collected and the solvent was evaporated. The obtained residue (1.1 g -38 %) was crystallised from CH_3CN . The resulting precipitate was filtered off and dried. A part of this fraction was extra dried, yielding compound 59.

Example B36

Preparation of 4,6-ethanediylidenepyrimido[4,5-b][6,1,10,13]benzoxatriazacycloheptadecine-12,15-dione, 19-chloro-8,9,10,11,13,14,16,21-octahydro-22-methoxy- (compound 60)

A mixture of intermediate 121 (0.00308 mol) in DMF (300 ml) was dropwise added overnight to a mixture of PyBOP (0.00616 mol) and DIPEA (0.0154 mol) in DMF

-138-

(300 ml), then extra PyBOP (0.00616 mol) and DIPEA (0.0154 mol) were added and the reaction mixture was stirred over the weekend. The solvent was evaporated under reduced pressure and the residue was dissolved in a 10 % solution of methanol in DCM and was then washed with water. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated under reduced pressure. The residue was purified by RP high-performance liquid chromatography. The pure product fractions were collected and then concentrated until precipitation occurred, yielding compound 60.

10 Example B37

Preparation of 4,6-ethenopyrimido[4,5-b][6,1,12]benzoxadiazacyclopentadecine, 17-bromo-16-fluoro-8,9,10,11,12,13,14,19-octahydro-20-methoxy- (compound 61)

TFA (2 ml) was added to a mixture of intermediate 127 (0.00055 mol) in DCM (10 ml), then the reaction mixture was stirred for 3 hours at RT and neutralised with a NaOH solution. The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated. The residue was purified by column chromatography (gradient eluent: DCM/CH₃OH). The pure product fractions were collected and the solvent was evaporated, yielding 0.042 g of compound 61.

Example B38

Preparation of 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-11,14-dione, 18-chloro-9,10,12,13,15,20-hexahydro-21-methoxy-10-[2-(4-morpholinyl)ethyl]- (compound 62)

HBTU (0.00195 mol) was added to a stirred solution of intermediate 130 (0.00069 mol) and DIPEA (0.00324 mol) in DMA (250 ml) at RT, then the reaction mixture was stirred for 3 hours and the solvent was co-evaporated with toluene under reduced pressure. The obtained residue was purified by RP high-performance liquid chromatography (eluent 1: NH₄OAc; eluent 2: NH₄HCO₃). The pure product fractions were collected and the solvent was evaporated under reduced pressure. The obtained residue (0.030 g) was crystallised from 2-propanol, then the resulting precipitate was filtered off and dried (vac.), yielding 0.0165 g of compound 62.

The following compound 63 was made accordingly;

-139-

4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-11,14-dione, 18-chloro-9,10,12,13,15,20-hexahydro-21-methoxy-10-(2-methoxyethyl)- (compound 63).

Example B39

Preparation of benzamide, 4-fluoro-N-(8,9,10,11,12,13-hexahydro-20-methoxy-4,6-ethanediylidene-19H-pyrimido[4,5-b][6,13,1]benzodioxazacyclopentadecin-16-yl)- (compound 64)

- A solution of intermediate 136 (0.0002 mol) in THF (20 ml) was stirred at RT and then ADDP (0.0003 mol) and tributyl- phosphine (0.0003 mol) were added. The
- 5 reaction mixture was stirred for 6 hours at RT and then extra ADDP (0.0003 mol) and tributyl- phosphine (0.0003 mol) were added. The resulting mixture was stirred for 1 hour and the solvent was evaporated under reduced pressure. The residue was stirred in methanol and filtered. The filter residue was stirred in boiling 2-propanol, then the resulting precipitate was filtered off and stirred in a mixture of CH₃OH/HCl(1N)/H₂O.
- 10 After filtration, the filter residue was stirred in a CH₃OH/NH₃ solution and the resulting precipitate was filtered off and dried (vac.) at 60°C, yielding 0.015 g of compound 64.

Example B40

Preparation of 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-11,14-dione, 18-chloro-9,10,12,13,15,20-hexahydro-21-methoxy-12,12-dimethyl- (compound 65)

- 15 A solution of PyBOP (0.0013 mol) and DIPEA (0.0065 mol) in DMA (70 ml) was stirred at RT and then a solution of intermediate 143 (0.0013 mol) in DMA (70 ml) was added dropwise. The reaction mixture was stirred for 18 hours at RT and the solvent was evaporated under reduced pressure. The residue was dissolved in DCM and was washed 2 times with a saturated NaHCO₃ solution and 2 times with water.
- 20 The organic layer was separated, dried (MgSO₄), filtered off and the solvent was evaporated. The dry residue was stirred in boiling 2-propanol, then the formed precipitate was filtered off, washed and dried (vacuum) at 60°C, yielding 0.133 g of compound 65, melting point 285°C.

-140-

Following compounds were made according to the synthesis of compound 65:

Compound No.	Name
66	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-11,14-dione, 18-chloro-9,10,12,13,15,20-hexahydro-21-methoxy-12-(1-methylethyl)- melting point: 335°C
67	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-11,14-dione, 18-chloro-9,10,12,13,15,20-hexahydro-21-methoxy-12-(2-methylpropyl)-
68	4,6-ethanediylidenepyrimido[4,5-b][6,1,10,13]benzoxatriazacycloheptadecine-12,15-dione, 19-chloro-8,9,10,11,13,14,16,21-octahydro-22-methoxy-13-(2-methylpropyl)-
69	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-11,14-dione, 18-chloro-9,10,12,13,15,20-hexahydro-21-methoxy- melting point: 292°C

Example B41

Preparation of 4,6-ethanediylidenepyrimido[4,5-b][6,1,10,14]benzoxatriazacycloheptadecin-12(13H)-one, 19-chloro-8,9,10,11,14,15,16,21-octahydro-22-methoxy- (compound 70)

- 5 DIPEA (0.00930 mol) was added to a solution of intermediate 149 (0.00155 mol) in dry DMF (10 ml) and the mixture was stirred for 15 min. then this solution was cannulated slowly to a solution of HBTU (0.00465 mol) in DMF (40 ml) and the reaction mixture was stirred for 30 min. The solvent was evaporated. The residue was purified by RP high-performance liquid chromatography. The product fractions were
10 collected and the solvent was evaporated, yielding 0.258 g of compound 70, melting point 236.4-237.3°C.

The following compounds were made accordingly

Compound No.	Name
71	4,6-ethanediylidenepyrimido[4,5-b]pyrrolo[2,1-k][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 19-chloro-9,10,11a,12,13,14,16,21-octahydro-22-methoxy- melting point 261.2-265°C

-141-

Compound No.	Name
72	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 17-chloro-9,10,12,13,14,19-hexahydro-20-methoxy-13-methyl-melting point 288.5-290.5°C
73	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 17-chloro-9,10,12,13,14,19-hexahydro-20-methoxy-melting point: 294.2-295.2°C
74	4,6-ethanediylidene-12H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 18-chloro-8,9,10,11,13,14,15,20-octahydro-21-methoxy-14-methyl-melting point: 240.0-240.3°C
75	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,13]benzoxatriazacyclohexadecin-11(12H)-one, 18-chloro-9,10,13,14,15,20-hexahydro-21-methoxy-melting point: 254.4-256.5°C
76	4,6-etheno-8H-pyrimido[4,5-b]pyrrolo[2,1-l][6,1,10,13]benzoxatriazacyclohexadecine-12,15(14H)-dione, 20-chloro-9,10,11,12a,13,17,22-heptahydro-23-methoxy-melting point: 350.5-352.5°C
77	4,6-ethanediylidene-12H-pyrimido[4,5-b]pyrrolo[2,1-l][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 20-chloro-8,9,10,11,12a,13,14,15,17,22-decahydro-23-methoxy-melting point: 129.8-132.8°C
78	4,6-ethanediylidenepyrimido[4,5-b]pyrrolo[2,1-k][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 19-chloro-18-fluoro-9,10,11a,12,13,14,16,21-octahydro-22-methoxy-melting point: 261.4-264.0°C
79	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 17-chloro-16-fluoro-9,10,12,13,14,19-hexahydro-20-methoxy-13-methyl-melting point: 306.3-307.4°C
80	4,6-ethanediylidene-12H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 18-chloro-17-fluoro-8,9,10,11,13,14,15,20-octahydro-21-methoxy-14-methyl-melting point 260.4-261.1°C
81	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 17-chloro-16-fluoro-9,10,12,13,14,19-hexahydro-20-methoxy-melting point 304.2-304.4°C
82	4,6-ethanediylidene-12H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 18-chloro-17-fluoro-8,9,10,11,13,14,15,20-octahydro-21-methoxy-melting point: 311.0-311.9°C
83	4,6-ethanediylidenepyrimido[4,5-b]pyrrolo[2,1-k][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 19-chloro-9,10,11a,12,13,14,16,21-octahydro-22-methoxy-melting point: 262.0-262.8°C

Compound No.	Name
84	4,6-ethanediylidene-12H-pyrimido[4,5-b]pyrrolo[2,1-l][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 20-chloro-8,9,10,11,12a,13,14,15,17,22-decahydro-23-methoxy-melting point: 231.9-232.8°C
85	4,6-ethanediylidenepyrimido[4,5-b]pyrrolo[2,1-k][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 19-chloro-9,10,11a,12,13,14,16,21-octahydro-13-hydroxy-22-methoxy-melting point: 279.4-280.7°C
86	4,6-ethanediylidene-13,16-ethano-8H-pyrimido[4,5-b][6,1,9,12,15]benzoxatetraazacyclooctadecin-11(12H)-one, 20-chloro-9,10,14,15,17,22-hexahydro-25-methoxy-melting point 296.4-297.0°C
87	8H-4,6-ethanediylidene-12,15-ethanopyrimido[4,5-b][6,1,9,14]benzoxatriazacycloheptadecin-11(12H)-one, 19-chloro-9,10,13,14,16,21-hexahydro-24-methoxy-melting point: 246.6-248.2°C
88	4,6-ethanediylidene-12,16-methano-6H-pyrimido[4,5-b][6,1,9,15]benzoxatriazacyclooctadecin-11(8H)-one, 20-chloro-9,10,12,13,14,15,17,22-octahydro-24-methoxy-melting point: 160-170°C
89	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 17-chloro-9,10,12,13,14,19-hexahydro-20-methoxy-12,13-dimethyl-melting point: 265°C
90	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 17-chloro-13-ethyl-9,10,12,13,14,19-hexahydro-20-methoxy-melting point: 261.1-262°C
91	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 17-chloro-9,10,12,13,14,19-hexahydro-12-(hydroxymethyl)-20-methoxy-melting point: 276.3-277.4°C
92	4,6-ethanediylidene-12H-pyrimido[4,5-b]pyrrolo[2,1-l][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 20-chloro-8,9,10,11,12a,13,14,15,17,22-decahydro-14-hydroxy-23-methoxy-melting point: 267.8-268.5°C
93	4,6-ethanediylidene-14,17-ethanopyrimido[4,5-b][6,1,10,13,16]benzoxatetraazacyclononadecin-12(13H)-one, 21-chloro-8,9,10,11,15,16,18,23-octahydro-26-methoxy-melting point: 286.8-287.6°C
94	4,6-ethanediylidene-13,16-ethano-6H-pyrimido[4,5-b][6,1,10,15]benzoxatriazacyclooctadecin-12(13H)-one, 20-chloro-8,9,10,11,14,15,17,22-octahydro-25-methoxy-melting point: 253.1-255.9°C
95	12H-4,6-ethanediylidene-13,17-methanopyrimido[4,5-b][6,1,10,16]benzoxatriazacyclononadecin-12-one, 21-chloro-8,9,10,11,13,14,15,16,18,23-decahydro-25-methoxy-melting point: 240.1-242.8°C

-143-

Compound No.	Name
96	4,6-ethanediylidene-12H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 18-chloro-8,9,10,11,13,14,15,20-octahydro-21-methoxy-13,14-dimethyl-melting point: 241.9-243.0°C
97	4,6-ethanediylidene-12H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 18-chloro-14-ethyl-8,9,10,11,13,14,15,20-octahydro-21-methoxy-melting point: 212.8-214.0°C
98	4,6-ethanediylidene-12H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 18-chloro-8,9,10,11,13,14,15,20-octahydro-13-(hydroxymethyl)-21-methoxy-melting point: 287.6-288.3°C
99	4,6-ethanediylidene-12H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 18-chloro-8,9,10,11,13,14,15,20-octahydro-21-methoxy-melting point: 304.6-304.8°C
166	4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecin-13(8H)-one, 19-chloro-15-ethyl-9,10,11,12,14,15,16,21-octahydro-22-methoxy-
167	4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecin-13(8H)-one, 19-chloro-9,10,11,12,14,15,16,21-octahydro-22-methoxy-14,15-dimethyl-

Example B42

Preparation of 4,6-ethanediylidenepyrimido[4,5-b]pyrrolo[2,1-m][6,1,11,14]benzoxatriazacycloheptadecin-13(8H)-one, 21-chloro-9,10,11,12,13a,14,15,16,18,23-decahydro-24-methoxy- (compound 100)

A solution of intermediate 156 (0.0005 mol) and DIPEA (0.003 mol) was added to a solution of HBTU (0.0015 mol) and 1-hydroxy- 1H-benzotriazole (0.001 mol) in DMF dry (125 ml) and then the reaction mixture was reacted for 1 hour. The solvent was evaporated and the dry residue was purified by RP high-performance liquid chromatography. The product fractions were collected, sodium carbonate was added and the organic solvent was evaporated. DCM was added to the aqueous concentrate and the resulting mixture was extracted 3 times with DCM, then the organic extract was dried and collected, yielding 0.0394 g (16 %) of compound 100, melting point 226.3-227.7°C.

The following compounds were made accordingly;

-144-

4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecin-13(8H)-one, 19-chloro-9,10,11,12,14,15,16,21-octahydro-22-methoxy- (compound 101)
melting point: 286.7-287.2°C
4,6-ethanediylidenepyrimido[4,5-b]pyrrolo[2,1-m][6,1,11,14]benzoxatriazacycloheptadecin-13(8H)-one, 21-chloro-20-fluoro-9,10,11,12,13a,14,15,16,18,23-decahydro-24-methoxy- (compound 102), melting point: 234.7-236.8°C

Example B43

Preparation of 4,6-ethanediylidene-12H-pyrimido[4,5-b]pyrrolo[2,1-l][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 20-chloro-19-fluoro-8,9,10,11,12a,13,14,15,17,22-decahydro-23-methoxy- (compound 103)

A solution of intermediate 162 (0.001 mol) and DIPEA (1.034 ml) in DMF (20 ml) was added to a solution of PyBOP (0.003 mol) and 1-hydroxy-1*H*-benzotriazole (0.001 mol) in DMF (200 ml) and then the reaction mixture was purified by RP high-performance liquid chromatography (CH₃CN/NH₄OAc buffer). The product fractions were collected, sodium carbonate was added and the organic solvent was evaporated (precipitation). The aqueous concentrate was cooled in the fridge, then filtered and washed with water, yielding: 0.2087 g (43 %) of compound 103 melting point 241.6-242.6°C.

The following compound was made accordingly;

4,6-ethanediylidenepyrimido[4,5-b]pyrrolo[2,1-k][6,1,9,12]benzoxatriazacyclopentadecine-11(8H)-thione, 19-chloro-18-fluoro-9,10,11a,12,13,14,16,21-octahydro-22-methoxy- (compound 104)
melting point: 211.3-212.7°C

Example B44

a) preparation of 4,6-ethanediylidenepyrimido[4,5-b][6,1,16]benzoxadiazacyclononadecin-16(17H)-one, 21-chloro-8,9,10,13,14,15,18,23-octahydro-24-methoxy- (B) (compound 105) and

4,6-ethanediylidenepyrimido[4,5-b][6,1,16]benzoxadiazacyclononadecin-16(17H)-one, 21-chloro-8,9,10,13,14,15,18,23-octahydro-24-methoxy- (A) (compound 106)

-145-

A mixture of intermediate 165 (0.000424 mol) and Grubbs II catalyst (0.000042) in DCM (200 ml; degassed) was stirred for 6 hours at RT and under N₂, then the solvent was evaporated and the residue was purified by RP high-performance liquid chromatography. Two product fractions were collected and the solvent was evaporated, yielding 0.046 g (23.3%) of compound 106 (A) and 0.078 g (39.5 %) of compound 105(B).

b) preparation of 4,6-ethanediylidenepyrimido[4,5-b][6,1,16]benzoxadiazacyclononadecin-16(17*H*)-one, 21-chloro-8,9,10,11,12,13,14,15,18,23-decahydro-24-methoxy- (compound 107)

A mixture of compound 105 (0.000064 mol) in THF (15 ml) and methanol (15 ml) was hydrogenated for 3 hours with Pt/C 5% (0.03 g) as a catalyst. After uptake of H₂ (1 equiv.), the catalyst was filtered off and the filtrate was evaporated, yielding compound 107.

The following compounds were made accordingly;

4,6-ethanediylidene-8*H*-pyrimido[4,5-b][6,1,15]benzoxadiazacyclooctadecin-15(16*H*)-one, 20-chloro-9,12,13,14,17,22-hexahydro-23-methoxy- (compound 108).

Example B45

a) Preparation of carbamic acid, (20-chloro-9,10,13,14,15,16,17,22-octahydro-23-methoxy-15-oxo-4,6-ethanediylidene-8*H*-pyrimido[4,5-b][6,1,15]benzoxadiazacyclooctadecin-14-yl)-, 1,1-dimethylethyl ester (A) (compound 109) and

Preparation of carbamic acid, (20-chloro-9,10,13,14,15,16,17,22-octahydro-23-methoxy-15-oxo-4,6-ethanediylidene-8*H*-pyrimido[4,5-b][6,1,15]benzoxadiazacyclooctadecin-14-yl)-, 1,1-dimethylethyl ester (B) (compound 110)

A solution of intermediate 169 (0.0015 mol) and Grubbs II catalyst (0.00015) in DCM (150 ml) was stirred overnight at RT, then the solvent was evaporated and the residue was purified by RP high-performance liquid chromatography. Two product fractions were collected and the solvent was evaporated, yielding 0.110 g of compound 109 (A) and 0.064 g of compound 110 (B).

-146-

b) Preparation of carbamic acid, (20-chloro-9,10,11,12,13,14,15,16,17,22-decahydro-23-methoxy-15-oxo-4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,15]benzoxadiazacyclooctadecin-14-yl)-, 1,1-dimethylethyl ester (compound 111)

A mixture of compound 109 (0.00025 mol) in THF (15 ml) and methanol (15 ml) was hydrogenated for 3 hours with Pt/C 5% (0.1 g) as a catalyst. After uptake of H₂ (1 equiv.), the catalyst was filtered off and the filtrate was evaporated. The residue was filtered over silica gel with DCM/CH₃OH (10/1) and the filtrate was evaporated, then
5 the obtained residue was crystallised from methanol and the resulting solids were collected, yielding compound 111.

Following compound was made accordingly:

carbamic acid, (18-chloro-11,12,13,14,15,20-hexahydro-21-methoxy-13-oxo-4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,13]benzoxadiazacyclohexadecin-12-yl)-, 1,1-dimethylethyl ester (compound 165)

c) Preparation of 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,15]benzoxadiazacyclooctadecin-15(16H)-one, 14-amino-20-chloro-9,10,11,12,13,14,17,22-octahydro-23-methoxy- .HCl (1:2) (compound 112)

6N HCl in 2-propanol (5 ml) was added to a solution of compound 111 (0.000088 mol) in THF (q.s.) and the reaction mixture was stirred for 1 hour at RT and then the
10 solvent was evaporated, yielding 0.050 g of compound 112, isolated as a hydrochloric acid salt.

d) Preparation of 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,15]benzoxadiazacyclooctadecin-15(16H)-one, 20-chloro-14-(dimethylamino)-9,10,11,12,13,14,17,22-octahydro-23-methoxy- (compound 113)

A mixture of compound 112 (0.000085 mol) and formaldehyde (0.00052 g) in methanol (20 ml) was hydrogenated with Pt/C 5% (0.04 g) as a catalyst in the presence of thiophene solution (0.04 ml). After uptake of H₂ (2 equiv.), the catalyst
15 was filtered off and the filtrate was evaporated. The obtained residue was purified by RP high-performance liquid chromatography, then the product fractions were collected and the solvent was evaporated, yielding compound 113.

-147-

Example B46

Preparation of 4,6-ethanediylidene-8*H*-pyrimido[4,5-
b][6,1,10,13]benzoxatriazacyclohexadecine, 18-chloro-9,10,11,12,13,14,15,20-octahydro-
21-methoxy-14-methyl- (compound 114)

A solution of intermediate 186 (0.00095 mol) in dioxane (10 ml), water (5 ml) and HCl concentrated (5 ml) was stirred for 27 hours at 50°C and then the reaction mixture was poured out into a saturated aqueous NaHCO₃ solution and extracted with
5 DCM. The organic extract was then dried and filtered over potassium carbonate. NaBH(OAc)₃ (0.00095 mol) was immediately added. The reaction mixture was stirred for 1 hour at RT. The mixture was purified by RP high-performance liquid chromatography, then the product fractions were collected and the solvent was evaporated, yielding 0.0576 g of compound 114, melting point 202.8-203.6°C.

10 The following compounds were made accordingly;

Compound No.	Name
115	4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine, 19-chloro-8,9,10,11,12,13,14,15,16,21-decahydro-22-methoxy-15-methyl-melting point: 196.9-197.8°C
116	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine, 17-chloro-8,9,10,11,12,13,14,19-octahydro-20-methoxy-13-methyl-melting point: 195.8-196.6°C
117	4,6-ethanediylidene-8 <i>H</i> -pyrimido[4,5-b][6,1,12,15]benzoxatriazacyclooctadecine, 20-chloro-9,10,11,12,13,14,15,16,17,22-decahydro-23-methoxy-16-methyl-

Example B47

Preparation of 4,6-ethanediylidenepyrimido[4,5-
b][6,1,12]benzoxadiazacyclopentadecine, 17-chloro-8,9,12,13,14,19-hexahydro-20-
methoxy-13-methyl- ±75% E and ± 25% Z (compound 118)

15 Grubbs II catalyst (a total of 0.0012 mol) was added in several portions to a solution of intermediate 191 (0.0016 mol) in DCM (100 ml) and the reaction mixture was stirred and refluxed for a total of 4 days. The obtained mixture was purified 2 times by RP high-performance liquid chromatography. The product fractions were collected and the solvent was evaporated, yielding 0.0116 g of compound 118.

-148-

Example B48

Preparation of 4,6-ethanediylidene-8*H*-pyrimido[4,5-*b*][6,1,9,12]benzoxatriazacyclohexadecine-9,12-dione, 18-chloro-10,11,13,14,15,20-hexahydro-21-methoxy-11-methyl- hydrate (1:1) (compound 119)

A mixture of intermediate 197 (0.0010 mol) and DIPEA (0.0040 mol) in DMF dry (50 ml) was slowly added to a solution of 1-[bis(dimethylamino)methylene]- 1*H*-

- 5 benzotriazolium,, hexafluorophosphate(1-), 3-oxide (0.0025 mol) in DMF dry (200 ml) at RT, then the reaction mixture was quenched with water (5 ml) and the solvent was evaporated. The obtained residue was purified by RP high-performance liquid chromatography. The product fractions were collected and the organic solvent was evaporated, yielding 0.024 g of compound 119.

- 10 The following compounds were made accordingly;

Compound No.	Name
120	4,6-ethanediylidenepyrimido[4,5- <i>b</i>][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8 <i>H</i> ,13 <i>H</i>)-dione, 17-chloro-10,11,14,19-tetrahydro-20-methoxy-11-methyl-
121	4,6-ethanediylidenepyrimido[4,5- <i>b</i>][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8 <i>H</i> ,13 <i>H</i>)-dione, 17-chloro-10,11,14,19-tetrahydro-20-methoxy-11-(1-methylethyl)-
122	4,6-ethanediylidenepyrimido[4,5- <i>b</i>][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8 <i>H</i> ,13 <i>H</i>)-dione, 17-chloro-10,11,14,19-tetrahydro-20-methoxy-11-(phenylmethyl)-
123	4,6-ethanediylidenepyrimido[4,5- <i>b</i>][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8 <i>H</i> ,13 <i>H</i>)-dione, 10,11,14,19-tetrahydro-20-methoxy-11-methyl-
124	4,6-ethanediylidenepyrimido[4,5- <i>b</i>][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8 <i>H</i> ,13 <i>H</i>)-dione, 10,11,14,19-tetrahydro-20-methoxy-11-(1-methylpropyl)-
125	9,11-ethanediylidenepyrimido[4,5- <i>b</i>]pyrrolo[1,2- <i>i</i>][6,1,9,12]benzoxatriazacyclopentadecine-14,19(5 <i>H</i> ,13 <i>H</i>)-dione, 16,17,18,18a,20,21-hexahydro-22-methoxy-
126	9,11-ethanediylidenepyrimido[4,5- <i>b</i>]pyrrolo[1,2- <i>i</i>][6,1,9,12]benzoxatriazacyclopentadecine-14,19(5 <i>H</i> ,13 <i>H</i>)-dione, 3-chloro-16,17,18,18a,20,21-hexahydro-22-methoxy-
127	4,6-ethanediylidenepyrimido[4,5- <i>b</i>][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8 <i>H</i> ,13 <i>H</i>)-dione, 17-chloro-10,11,14,19-tetrahydro-11-(1-hydroxyethyl)-20-methoxy-

-149-

Compound No.	Name
128	4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine-11,14(8H,15H)-dione, 19-chloro-9,10,12,13,16,21-hexahydro-22-methoxy-13-(1-methylpropyl)-
129	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8H,13H)-dione, 17-chloro-10,11,14,19-tetrahydro-11-(hydroxymethyl)-20-methoxy-
130	4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine-11,14(8H,15H)-dione, 19-chloro-9,10,12,13,16,21-hexahydro-13-(hydroxymethyl)-22-methoxy-
131	4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine-11,14(8H,15H)-dione, 19-chloro-9,10,12,13,16,21-hexahydro-22-methoxy-13-methyl-
132	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8H,13H)-dione, 17-chloro-10,11,14,19-tetrahydro-20-methoxy-11,11-dimethyl-
133	9,11-ethanediylidenepyrimido[4,5-b]pyrrolo[1,2-i][6,1,9,12]benzoxatriazacyclopentadecine-14,19(5H,13H)-dione, 3-chloro-16,17,18,18a,20,21-hexahydro-17-hydroxy-22-methoxy-
134	4,6-ethanediylidene-8H-pyrimido[4,5-b]pyrrolo[1,2-i][6,1,12,15]benzoxatriazacyclooctadecine-12,17(18H)-dione, 22-chloro-9,10,11,14,15,16,16a,19,24-nonahydro-25-methoxy-
135	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-9,12-dione, 18-chloro-10,11,13,14,15,20-hexahydro-21-methoxy-11,11-dimethyl-
136	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacyclooctadecine-12,15(16H)-dione, 20-chloro-9,10,11,13,14,17,22-heptahydro-23-methoxy-14-(2-methylpropyl)-
137	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacyclooctadecine-12,15(16H)-dione, 20-chloro-9,10,11,13,14,17,22-heptahydro-23-methoxy-14,14-dimethyl-
138	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacyclooctadecine-12,15(16H)-dione, 20-chloro-9,10,11,13,14,17,22-heptahydro-23-methoxy-14-(phenylmethyl)-
139	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacyclooctadecine-12,15(16H)-dione, 20-chloro-9,10,11,13,14,17,22-heptahydro-23-methoxy-14-methyl-
140	1,21-ethanediylidene-5H-pyrimido[4,5-b]pyrrolo[1,2-i][6,1,9,12]benzoxatriazacyclohexadecine-13,18(19H)-dione, 7-chloro-10,11,12,13a,14,15,16-heptahydro-23-methoxy-
141	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-9,12-dione, 18-chloro-10,11,13,14,15,20-hexahydro-21-methoxy-11-(2-methylpropyl)-
142	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-9,12-dione, 18-chloro-10,11,13,14,15,20-hexahydro-11-(1-hydroxyethyl)-21-methoxy-
143	4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine-11,14(8H,15H)-dione, 19-chloro-9,10,12,13,16,21-hexahydro-22-methoxy-13-(2-methylpropyl)-

-150-

Compound No.	Name
144	4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine-11,14(8H,15H)-dione, 19-chloro-9,10,12,13,16,21-hexahydro-22-methoxy-13,13-dimethyl-
145	4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine-11,14(8H,15H)-dione, 19-chloro-9,10,12,13,16,21-hexahydro-22-methoxy-13-(phenylmethyl)-
146	4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine-11,14(8H,15H)-dione, 19-chloro-9,10,12,13,16,21-hexahydro-13-(1-hydroxyethyl)-22-methoxy-
147	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacyclooctadecine-12,15(16H)-dione, 20-chloro-9,10,11,13,14,17,22-heptahydro-14-(1-hydroxyethyl)-23-methoxy-
148	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-9,12-dione, 18-chloro-10,11,13,14,15,20-hexahydro-11-(hydroxymethyl)-21-methoxy-
149	1,21-ethanediylidene-5H-pyrimido[4,5-b]pyrrolo[1,2-i][6,1,9,12]benzoxatriazacyclohexadecine-13,18(19H)-dione, 7-chloro-10,11,12,13a,14,15,16-heptahydro-15-hydroxy-23-methoxy-

Example B49

Preparation of 4,6-ethenopyrimido[4,5-b][6,1,9,14]benzoxatriazacycloheptadecine-9,14(8H,15H)-dione, 19-chloro-10,11,12,13,16,21-hexahydro-22-methoxy- (compound 150)

Intermediate 122 (0.001 mol) and DIPEA (0.004 mol) were added to a mixture of PyBOP (0.003 mol) in DMF (250 ml) and the reaction mixture was stirred for 2 hours, then water was added and the solvent was evaporated. The obtained residue was purified by RP high-performance liquid chromatography. The product fractions were collected and the organic solvent was evaporated. The aqueous concentrate was allowed to precipitate overnight in the fridge and the resulting solids were then filtered off, yielding 0.093 g (20 %) of compound 150.

10

The following compounds were made accordingly;

Compound No.	Name
--------------	------

-151-

Compound No.	Name
151	4,6-ethenopyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8H,13H)-dione, 17-chloro-10,11,14,19-tetrahydro-20-methoxy-
152	4,6-etheno-8H-pyrimido[4,5-b][6,1,9,13]benzoxatriazacyclohexadecine-9,13(10H,14H)-dione, 18-chloro-11,12,15,20-tetrahydro-21-methoxy-
153	4,6-ethenopyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine-11,14(8H,15H)-dione, 19-chloro-9,10,12,13,16,21-hexahydro-22-methoxy-
154	4,6-ethenopyrimido[4,5-b][6,1,11,16]benzoxatriazacyclononadecine-11,16(8H,17H)-dione, 21-chloro-9,10,12,13,14,15,18,23-octahydro-24-methoxy-
155	4,6-etheno-8H-pyrimido[4,5-b][6,1,11,15]benzoxatriazacyclooctadecine-11,15(12H,16H)-dione, 20-chloro-9,10,13,14,17,22-hexahydro-23-methoxy-
156	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8H,13H)-dione, 10,11,14,19-tetrahydro-20-methoxy-
157	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,13]benzoxatriazacyclohexadecine-9,13(10H,14H)-dione, 11,12,15,20-tetrahydro-21-methoxy-
158	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,14]benzoxatriazacycloheptadecine-9,14(8H,15H)-dione, 10,11,12,13,16,21-hexahydro-22-methoxy-
159	4,6-ethanediylidenepyrimido[4,5-b][6,1,9,12]benzoxatriazacyclopentadecine-9,12(8H,13H)-dione, 17-chloro-10,11,14,19-tetrahydro-20-methoxy-10-methyl-
160	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-9,12-dione, 18-chloro-10,11,13,14,15,20-hexahydro-21-methoxy-
161	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,14]benzoxatriazacyclooctadecine-9,14-dione, 20-chloro-10,11,12,13,15,16,17,22-octahydro-23-methoxy-
162	4,6-ethanediylidenepyrimido[4,5-b][6,1,12,16]benzoxatriazacyclononadecine-12,16(13H,17H)-dione, 21-chloro-8,9,10,11,14,15,18,23-octahydro-24-methoxy-
163	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,12,17]benzoxatriazacycloeicosine-12,17(18H)-dione, 22-chloro-9,10,11,13,14,15,16,19,24-nonahydro-25-methoxy-
164	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,9,12]benzoxatriazacyclohexadecine-9,12-dione, 18-chloro-10,11,13,14,15,20-hexahydro-21-methoxy-10-methyl-
168	4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,12,15]benzoxatriazacyclooctadecine-12,15(16H)-dione, 20-chloro-9,10,11,13,14,17,22-heptahydro-23-methoxy-13-methyl-

Compound identification

-152-

The compounds were identified by LC/MS using a gradient elution system on a reversed phase HPLC. The compounds are identified by their specific retention time and their protonated molecular ion MH^+ peak. The HPLC gradient was supplied by a Waters Alliance HT 2790 system with a column heater set at 40°C. Flow from the column was split to a Waters 996 photodiode array (PDA) detector and a Waters-Micromass ZQ mass spectrometer with an electrospray ionization source operated in positive and negative ionization mode. Reversed phase HPLC was carried out on a Xterra MS C18 column (3.5 μm , 4.6 x 100 mm) with a flow rate of 1.6 ml/min. Three mobile phases (mobile phase A 95% 25mM ammoniumacetate + 5% acetonitrile; mobile phase B: acetonitrile; mobile phase C: methanol) were employed to run a gradient condition from 100 % A to 50% B and 50% C in 6.5 minutes, to 100 % B in 1 minute, 100% B for 1 minute and reequilibrate with 100 % A for 1.5 minutes. An injection volume of 10 μL was used.

Mass spectra were acquired by scanning from 100 to 1000 in 1s using a dwell time of 0.1 s. The capillary needle voltage was 3kV and the source temperature was maintained at 140°C. Nitrogen was used as the nebulizer gas. Cone voltage was 10 V for positive ionization mode and 20 V for negative ionization mode. Data acquisition was performed with a Waters-Micromass MassLynx-Openlynx data system.

Table : retention time (RT in minutes) and molecular weight as the MH^+

Compound No.	Rt	MH^+
64	6.38	503
38	6.64	414
40	4.93	427
71	5.66	454
150	3.84	454
153	3.89	546
120	3.77	442

Compound No.	Rt	MH^+
121	4.32	470
122	4.84	518
60	4.53	456
123	3.05	408
124	4.08	450
125	3.56	434
156	2.83	394

-153-

Compound No.	Rt	MH ⁺
157	3.01	408
158	3.3	422
126	4.33	468
159	4.02	442
127	3.62	472
128	5.34	512
129	3.47	458
130	3.93	486
131	4.34	470
160	4.04	442
161	3.86	470
106	8.73	467
105	9.6	467
132	3.74	456
133	3.63	484
108	6.09	453
117	5.95	456
63	4.75	500
162	4	484
163	3.91	498
134	4.51	510
135	4.56	470

Compound No.	Rt	MH ⁺
168	4.17	484
136	4.62	526
137	4.18	498
138	4.62	560
139	3.88	456
119	4.5	456
140	5.03	482
164	4.74	456
141	5.41	498
142	4.17	486
143	5.16	512
144	4.42	484
145	5.18	546
147	4.3	514
110	6.2	568
109	6.14	568
148	3.92	472
149	4.45	498
111	6.26	570
165	5.82	540

-154-

C. Pharmacological examplesExample C.1 : in vitro inhibition of EGFR using a Scintillant Proximity Assay

In the present EGFR SPA kinase reaction assay, a kinase substrate consisting of biotinylated poly(L-glutamic acid-L-tyrosine) (poly(GT)biotin), is incubated with the aforementioned protein in the presence of (³³P) radiolabeled ATP. (³³P) phosphorylation of the substrate is subsequently measured as light energy emitted using streptavidin-coated SPA beads (Amersham Pharmacia Biotech) by trapping and quantifying the binding of the biotin tagged and radiolabeled substrate.

Detailed description

The EGFR SPA kinase reaction is performed at 30°C for 60 minutes in a 96-well microtiter plate. For each of the tested compounds a full dose response 1.10^{-6} M to 1.10^{-10} M has been performed. IRESSA[®] and Tarceva[™] (erlotinib) were used as reference compounds. The 100 µl reaction volume contains 54.5 mM TrisHCl pH 8.0, 10 mM MgCl₂, 100µM Na₃VO₄, 5.0 µM unlabeled ATP, 1mM DTT, 0.009% BSA, 0.8 µCi ³³P-ATP, 0.35 µg/well poly(GT)biotin and 0.5 µg EGFR-kinase domain/well.

The reaction is stopped by adding to each well 100 µl of the streptavidin beads (10 mg/ml in PBS + 100 mM EDTA + 100 µM ATP). The plates are then shaken at 300 rpm for 30 min to allow binding of the biotinylated substrate to the streptavidin coated beads. Then the beads are allowed to settle at the bottom of the plate for 30 minutes. The microtiterplates are centrifuged at 800 rpm for 10 minutes and the amount of phosphorylated (³³P) Poly(GT)biotin is determined by counting (30sec/well) in a microtiterplate scintillation counter.

Example C.2 : in vitro inhibition of EGFR

The *in vitro* inhibition of EGFR was assessed using either the Flash Plate technology or the glass-fiber filter technology as described by Davies, S.P. et al., Biochem J. (2000), 351; p.95-105. The Flash Plate technology is generally described by B.A. Brown *et al.* in High Throughput Screening (1997), p.317-328. Editor(s): Devlin, John P. Publisher: Dekker, New York, N. Y.

In the Flash Plate EGFR kinase reaction assay, a kinase substrate consisting of biotinylated poly(L-glutamic acid-L-tyrosine) (poly(GT)biotin), is incubated with the aforementioned protein in the presence of (³³P) radiolabeled ATP. (³³P) phosphorylation

-155-

of the substrate is subsequently measured as light energy emitted using a streptavidin-coated Flash Plate (PerkinElmer Life Sciences) by trapping and quantifying the binding of the biotin tagged and radiolabeled substrate.

5 *Detailed description*

The EGFR kinase reaction is performed at 30°C for 60 minutes in a 96-well microtiter FlashPlate (PerkinElmer Life Sciences). For each of the tested compounds a full dose response 1.10^{-6} M to 1.10^{-10} M has been performed. IRESSA® and Tarceva™ (erlotinib) were used as reference compounds. The 100 µl reaction volume contains 54.5 mM
10 TrisHCl pH 8.0, 10 mM MgCl₂, 100µM Na₃VO₄, 5.0 µM unlabeled ATP, 1mM DTT, 0.009% BSA, 0.8 µCi AT³³P, 0.35 µg/well poly(GT)biotin and 0.5 µg EGFR-kinase domain/well.

The reaction is stopped by aspirating the reaction mixture and washing the plate 3x with 200 µl wash/stop buffer (PBS + 100 mM EDTA). After the final wash step 200 µl
15 of wash/stop buffer was added to each well and the amount of phosphorylated (³³P) Poly(GT)biotin determined by counting (30 sec/well) in a microtiterplate scintillation counter.

In the glass-fiber filter technology EGFR kinase reaction assay, a kinase substrate
20 consisting of poly(L-glutamic acid-L-tyrosine) (poly(GT)), is incubated with the aforementioned protein in the presence of (³³P) radiolabeled ATP. (³³P) Phosphorylation of the substrate is subsequently measured as radioactivity bound on a glassfiber-filter.

Detailed description

25 The EGFR kinase reaction is performed at 25°C for 10 minutes in a 96-well microtiterplate. For each of the tested compounds a full dose response 1.10^{-6} M to 1.10^{-10} M has been performed. IRESSA® and Tarceva™ (erlotinib) were used as reference compounds. The 25 µl reaction volume contains 60 mM TrisHCl pH 7.5, 3 mM MgCl₂, 3 mM Mn Cl₂, 3 µM Na₃VO₄, 50 µg/ml PEG20000, 5.0 µM unlabeled ATP,
30 1mM DTT, 0.1 µCi AT³³P, 62.5 ng/well poly(GT) and 0.5 µg EGFR-kinase domain/well.

The reaction is stopped by adding 5 µl of a 3% phosphoric acid solution. 10 µl of the reaction mixture is then spotted onto a Filtermat A filter (Wallac) and washed 3 times for 5 min. in 75 mM phosphoric acid and 1 time for 5 min. in methanol prior to drying
35 and quantification on the Typhoon (Amersham) using a LE phosphorage storage screen.

Example C.3: Serum starved proliferation assay on the ovarian carcinoma SKOV3 cells

5 The ovarian carcinoma cell line (SKOV3) was used in an epidermal growth factor stimulated cell proliferation assay, to assess the inhibitory effect of the compounds on EGF in whole cells.

In a first step the SKOV3 cells were incubated for 24 hours in the presence of 10% FCS serum. In the second step the cells were incubated with the compounds to be tested in a serum free condition (37 °C and 5% (v/v) CO₂) and subsequently stimulated for 72
10 hours with EGF at a final concentration of 100 ng/ml. The effect of the compounds on the EGF stimulation was finally assessed in a standard MTT cell viability assay.

Alternatively, the SKOV3 cells were incubated for 24 hours in the presence of 10% FCS serum. In the second step the cells were incubated for 72 hours with the
15 compounds to be tested and the effect of the compounds on cell proliferation was finally assessed in a standard MTT cell viability assay.

Example C.4: ELISA assay of EGFR tyrosine kinase activity

20 The EGFR ELISA is generally described by Yang, E.B. *et al.*, 2001, Biochimica et Biophysica Acta, 1550; 144.

For the determination of EGFR tyrosine kinase activity, 100 µl of 0.4 µg/ml poly(Glu, Tyr) in PBS was coated per well on a 96-well microplate at 37°C overnight. The non-specific binding sites were subsequently blocked by incubation for 30 minutes at room
25 temperature with 200 µl BSA dilution (10 mg/ml in PBS) per well. After washing three times with PBS the plates were either used immediately or stored at 4°C.

Prior to the determination of the EGFR tyrosine kinase activity, the coated plates were washed two times with PBS. Next, to each well 88 µl of an ATP dilution (50 mM Tris HCl pH 8.0, 10 mM MgCl₂, 100 µM Na₃VO₄, 1 mM DTT, 5 µM ATP) and 2 µl with
30 various concentrations of the compounds to be tested, were added. The EGFR tyrosine kinase-catalysed reaction was started by the addition of 10 µl of diluted EGFR (dilution → 0.05 µg per well enzyme diluted in 50 mM TrisHCl pH 8.0 + 0.1% BSA).

After incubation at room temperature for 10 min, the reaction was stopped by washing five times with PBS with 0.1% Tween 20. Subsequently 100 µl of recombinant anti-phosphotyrosine horseradish peroxidase conjugate (1:2500) in BSA (10mg/ml in PBS)
35 was added. After incubation at room temperature for 1 h, the microplate was washed five times with PBS/Tween 20. After the microplate was incubated with 100 µl of

-157-

TMB-ELISA (1-step Ultra TMB-ELISA, Pierce) until colour development, 100 µl of 0.5 M H₂SO₄ was added to stop the reaction, and it was read in a microplate reader at 450-655 nm.

5 Example C.5: Proliferation assay on the squamous carcinoma cell line A431 cells

The squamous carcinoma cell line (A431) was used in a cell proliferation assay, to assess the inhibitory effect of the compounds in whole cells.

10 In a first step the A431 cells were incubated for 24 hours in the presence of 10% FCS serum. In the second step the cells were incubated for 72 hours with the compounds to be tested at a final concentration of 100 ng/ml. The effect of the compounds on the cell proliferation was finally assessed in a standard MTT cell viability assay.

-158-

The following tables provides the IC₅₀ values of the compounds according to the invention, obtained using the above mentioned kinase assays.

Compound number	EGFR SPA (C1) : IC ₅₀ in nM	Kinase activity-(C2) : pIC ₅₀	SKOV3 cell (C3) : pIC ₅₀	A431 cell (C5) : pIC ₅₀	EGFR ELISA (C4) : IC ₅₀ in nM
1	>100	< 5.0	< 5.7	< 6.0	>1000
2	>100	7.0	< 5.7	< 6.0	>1000
3	3.61	7.8	5.87	< 6.0	>1000
4	32.58	7.3	5.54	NT	556
5	81.10	6.4	5.32	NT	>1000
6	4.40	7.6	5.74	NT	359
7	6.64	7.1	< 5	NT	>1000
8	3.97	7.5	< 5	NT	329
9	6.79	7.4	NT	NT	>1000
16	NT	5.6	NT	NT	>1000
18	NT	7.1	< 5	NT	>1000
17	NT	7.4	5.08	5.6	269
10	NT	7.3	< 5	< 5.5	>1000
12	NT	< 5	< 5	< 5.5	>1000
14	NT	< 5	< 5	< 5.5	>1000
11	NT	< 5	< 5	< 5.5	>1000
13	NT	6.5	< 5	< 5.5	>1000
20	NT	7.6	6.64	5.9	NT
19	NT	7.2	5.3	< 5.5	NT
21	NT	8.0	6.09	< 5.5	158
22	NT	7.8	6.59	5.8	38.4
23	NT	8.0	7.22	6.3	NT
24	NT	7.7	5.5	< 5.5	NT

-159-

Compound number	EGFR SPA (C1) : IC50 in nM	Kinase activity.(C2) : pIC50	SKOV3 cell (C3) : pIC50	A431 cell (C5) : pIC50	EGFR ELISA (C4) : IC50 in nM
25	NT	6.2	< 5	< 5.5	NT
26	NT	8.6	6.92	5.7	NT
28	NT	7.6	5.84	5.8	NT
29	NT	6.1	< 5	< 5.5	NT
30	NT	7.5	5.99	< 6.0	NT
32	NT	8.4	6.54	NT	NT
33	NT	7.6	5.76	5.8	NT
34	NT	7.2	< 5	< 5.5	NT
35	NT	5.4	< 5	< 5.5	NT
36	NT	7.8	6.78	< 5.5	NT
37	NT	< 5	< 5	< 5.5	NT
38	NT	7.7	6.9	5.9	NT
39	NT	7.7	6.7	5.8	NT
40	NT	7.5	7.3	6.2	NT

-160-

Compound number	Kinase activity.(C2) : pIC50	SKOV3 cell (C3) : pIC50	Example number
64	6.0	6.4	B39
71	7.5	5.5	B41
72	6.8	5.0	B41
73	7.4	5.4	B41
74	7.6	6.0	B41
75	7.2	5.2	B41
70	8.0	5.5	B41
151	5.5	5.0	B49
76	7.6	6.7	B41
77	7.8	6.1	B41

Compound number	Kinase activity.(C2) : pIC50	SKOV3 cell (C3) : pIC50	Example number
150	7.7	5.1	B49
152	7.5	5.0	B49
153	7.8	6.1	B49
154	7.4	5.6	B49
155	6.5	6.0	B49
61	7.8	6.4	B39
120	5.7	6.0	B48
121	5.6	5.0	B48
122	6.4	5.0	B48
66	7.1	8.0	B40

-161-

Compound number	Kinase activity.(C2) : pIC50	SKOV3 cell (C3) : pIC50	Example number
65	7.9	7.7	B40
67	7.0	6.3	B40
68	7.9	8.2	B40
69	7.3	7.2	B40
60	7.8	8.7	B36
123	5.2	5.0	B48
124	5.0	5.0	B48
125	5.1	5.0	B48
156	5.0	5.0	B49
157	5.0	5.0	B49

Compound number	Kinase activity.(C2) : pIC50	SKOV3 cell (C3) : pIC50	Example number
158	5.3	5.0	B49
78	7.5	6.5	B41
114	8.0	7.4	B46
79	6.2	5.0	B41
80	7.7	7.2	B41
81	6.6	6.6	B41
82	7.7	7.1	B41
99	7.6	6.7	B41
126	5.7	5.6	B48
159	6.5	5.7	B49

-162-

Compound number	Kinase activity (C2) : pIC50	SKOV3 cell (C3) : pIC50	Example number
127	6.6	5.8	B48
128	8.0	6.7	B48
129	5.8	5.4	B48
130	7.6	6.9	B48
131	7.6	7.0	B48
160	7.2	6.2	B49
161	7.2	5.9	B49
83	6.3	6.1	B41
84	6.2	6.5	B41
103	7.2	7.3	B43

Compound number	Kinase activity (C2) : pIC50	SKOV3 cell (C3) : pIC50	Example number
101	6.6	7.5	B42
52	7.1	7	B34
53	6.7	6.3	B34
54	6.9	5.9	B34
55	6.3	6.7	B34
56	5.8	5.0	B34
57	5.2	6.4	B34
58	5.7	5.7	B34
44	6.2	5.1	B33
45	6.8	5.0	B33

-163-

Compound number	Kinase activity.(C2) : pIC50	SKOV3 cell (C3) : pIC50	Example number
42	6.3	7.2	B33
43	7.1	6.3	B33
46	6.7	5.1	B33
47	6.9	5.8	B33
48	6.5	5.8	B33
49	6.4	6.1	B33
50	7.1	6.8	B33
51		6.4	B33
100	8.1	6.9	B42
102	6.7	7.3	B42

Compound number	Kinase activity.(C2) : pIC50	SKOV3 cell (C3) : pIC50	Example number
106	7.0	7.2	B44
105	6.8	7.6	B44
104	6.0	6.5	B43
115	8.2	6.9	B46
116	6.7	7.2	B46
132	5.0	5.0	B48
133	5.8	5.1	B48
108	6.4	6.8	B44
117	6.7	5.4	B46
63	6.9	5.7	B38

-164-

Compound number	Kinase activity (C2) : pIC50	SKOV3 cell (C3) : pIC50	Example number
162	6.8	7.3	B49
163	7.0	6.4	B49
134	6.4	6.1	B48
135	5.4	6.0	B48
136	6.2	6.6	B48
137	6.6	6.6	B48
138	5.7	6.2	B48
139	6.4	6.8	B48
119	6.0	5.8	B48
140	6.7	6.7	B48

Compound number	Kinase activity (C2) : pIC50	SKOV3 cell (C3) : pIC50	Example number
164	6.3	6.8	B49
141	6.1	5.8	B48
142	7.0	6.2	B48
143	6.3	6.7	B48
144	6.8	6.9	B48
145	6.7	6.6	B48
146	6.7	6.2	B48
147	6.9	6.6	B48
62	6.4	5.3	B48
110	5.4	5.0	B45

-165-

Compound number	Kinase activity.(C2) : pIC50	SKOV3 cell (C3) : pIC50	Example number
109	5.7	6.5	B45
148	6.6	5.0	B48
149	6.9	5.8	B48

Compound number	Kinase activity.(C2) : pIC50	SKOV3 cell (C3) : pIC50	Example number
111	5.8	5.0	B45

D. Composition examples

- 5 The following formulations exemplify typical pharmaceutical compositions suitable for systemic administration to animal and human subjects in accordance with the present invention.

"Active ingredient" (A.I.) as used throughout these examples relates to a compound of formula (I) or a pharmaceutically acceptable addition salt thereof.

10 Example D.1 : film-coated tablets

Preparation of tablet core

- A mixture of A.I. (100 g), lactose (570 g) and starch (200 g) was mixed well and thereafter humidified with a solution of sodium dodecyl sulfate (5 g) and polyvinylpyrrolidone (10 g) in about 200 ml of water. The wet powder mixture was sieved, dried
15 and sieved again. Then there was added microcrystalline cellulose (100 g) and hydrogenated vegetable oil (15 g). The whole was mixed well and compressed into tablets, giving 10.000 tablets, each comprising 10 mg of the active ingredient.

Coating

- To a solution of methyl cellulose (10 g) in denaturated ethanol (75 ml) there was added a
20 solution of ethyl cellulose (5 g) in DCM (150 ml). Then there were added DCM (75 ml)


-166-

and 1,2,3-propanetriol (2.5 ml). Polyethylene glycol (10 g) was molten and dissolved in dichloromethane (75 ml). The latter solution was added to the former and then there were added magnesium octadecanoate (2.5 g), polyvinyl-pyrrolidone (5 g) and concentrated color suspension (30 ml) and the whole was homogenated. The tablet cores were coated
5 with the thus obtained mixture in a coating apparatus.

-167-

Claims

1. A compound having the formula



the *N*-oxide forms, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

Z represents O, CH₂, NH or S; in particular Z represents NH;

Y represents -C₃₋₉alkyl-, -C₃₋₉alkenyl-, -C₃₋₉alkynyl-, -C₃₋₇alkyl-CO-NH- optionally substituted with amino, mono- or di(C₁₋₄alkyl)amino or C₁₋₄alkyloxycarbonylamino-, -C₃₋₇alkenyl-CO-NH- optionally substituted with amino, mono- or di(C₁₋₄alkyl)amino or C₁₋₄alkyloxycarbonylamino-, -C₃₋₇alkynyl-CO-NH- optionally substituted with amino, mono- or di(C₁₋₄alkyl)amino or C₁₋₄alkyloxycarbonylamino-, -C₁₋₅alkyl-oxy-C₁₋₅alkyl-, -C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-, -C₁₋₅alkyl-NR¹⁴-CO-C₁₋₅alkyl-, -C₁₋₅alkyl-CO-NR¹⁵-C₁₋₅alkyl-, -C₁₋₆alkyl-CO-NH-, -C₁₋₆alkyl-NH-CO-, -C₁₋₃alkyl-NH-CS-Het²⁰-, -C₁₋₃alkyl-NH-CO-Het²⁰-, C₁₋₂alkyl-CO-Het²¹-CO-, -Het²²-CH₂-CO-NH-C₁₋₃alkyl-, -CO-NH-C₁₋₆alkyl-, -NH-CO-C₁₋₆alkyl-, -CO-C₁₋₇alkyl-, -C₁₋₇alkyl-CO-, -C₁₋₆alkyl-CO-C₁₋₆alkyl-, -C₁₋₂alkyl-NH-CO-CR¹⁶R¹⁷-NH-, -C₁₋₂alkyl-CO-NH-CR¹⁸R¹⁹-CO-, -C₁₋₂alkyl-CO-NR²⁰-C₁₋₃alkyl-CO-, -C₁₋₂alkyl-NR²¹-CH₂-CO-NH-C₁₋₃alkyl-, or -NR²²-CO-C₁₋₃alkyl-NH-;

X¹ represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO- C₁₋₂alkyl-, NR¹¹, -NR¹¹-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or -C₁₋₂alkyl-;

X² represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO- C₁₋₂alkyl-, NR¹², -NR¹²-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or -C₁₋₂alkyl-;

R¹ represents hydrogen, cyano, halo, hydroxy, formyl, C₁₋₆alkoxy-, C₁₋₆alkyl-, halo-phenyl-carbonylamino-, C₁₋₆alkoxy- substituted with halo,

-168-

C₁₋₄alkyl substituted with one or where possible two or more substituents selected from hydroxy or halo;

R² represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-, Het¹⁶-carbonyl-, C₁₋₄alkyloxycarbonyl-, C₁₋₄alkylcarbonyl-, aminocarbonyl-, mono-or
 5 di(C₁₋₄alkyl)aminocarbonyl-, Het¹, formyl, C₁₋₄alkyl-, C₂₋₆alkynyl-, C₃₋₆cycloalkyl-, C₃₋₆cycloalkyloxy-, C₁₋₆alkoxy-, Ar⁵, Ar¹-oxy-, dihydroxyborane , C₁₋₆alkoxy- substituted with halo,

C₁₋₄alkyl substituted with one or where possible two or more substituents selected from halo, hydroxy or NR⁵R⁶,

10 C₁₋₄alkylcarbonyl- wherein said C₁₋₄alkyl is optionally substituted with one or where possible two or more substituents selected from hydroxy or C₁₋₄alkyl-oxy-;

R³ represents hydrogen, C₁₋₄alkyl, or C₁₋₄alkyl substituted with one or more substituents selected from halo, C₁₋₄alkyloxy-, amino-, mono-or di(C₁₋₄alkyl)amino-,
 15 C₁₋₄alkyl-sulfonyl- or phenyl;

R⁴ represents hydrogen, hydroxy, Ar³-oxy, Ar⁴-C₁₋₄alkyloxy-, C₁₋₄alkyloxy-, C₂₋₄alkenyloxy- optionally substituted with Het¹² or R⁴ represents C₁₋₄alkyloxy substituted with one or where possible two or more substituents selected from C₁₋₄alkyloxy-, hydroxy, halo, Het²-, -NR⁷R⁸-, -carbonyl- NR⁹R¹⁰ or Het³-carbonyl-;

20 R⁵ and R⁶ are each independently selected from hydrogen or C₁₋₄alkyl;

R⁷ and R⁸ are each independently selected from hydrogen, C₁₋₄alkyl, Het⁸, aminosulfonyl-, mono- or di (C₁₋₄alkyl)-aminosulfonyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl-, hydroxycarbonyl-C₁₋₄alkyl-, C₃₋₆cycloalkyl, Het⁹-carbonyl-C₁₋₄alkyl-, Het¹⁰-carbonyl-, polyhydroxy-C₁₋₄alkyl-, Het¹¹-C₁₋₄alkyl- or Ar²-C₁₋₄alkyl-;

25 R⁹ and R¹⁰ are each independently selected from hydrogen, C₁₋₄alkyl, C₃₋₆cycloalkyl, Het⁴, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl- or polyhydroxy-C₁₋₄alkyl-;

R¹¹ represents hydrogen, C₁₋₄alkyl, Het⁵, Het⁶-C₁₋₄alkyl-, C₂₋₄alkenylcarbonyl- optionally substituted with Het⁷-C₁₋₄alkylaminocarbonyl-, C₂₋₄alkenylsulfonyl-,
 30 C₁₋₄alkyloxyC₁₋₄alkyl- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;

R¹² represents hydrogen, C₁₋₄alkyl, C₁₋₄alkyl-oxy-carbonyl-, Het¹⁸-C₁₋₄alkyl-, phenyl-C₁₋₄alkyl-oxy-carbonyl-, Het¹⁷, C₂₋₄alkenylcarbonyl- optionally substituted
 35 with Het¹⁹-C₁₋₄alkylaminocarbonyl-, C₂₋₄alkenylsulfonyl-, C₁₋₄alkyloxyC₁₋₄alkyl- or R¹² represents phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;

-169-


- R^{13} represents hydrogen, C_{1-4} alkyl, Het^{13} , $Het^{14}-C_{1-4}$ alkyl- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C_{1-4} alkyloxy-;
- R^{14} and R^{15} are each independently selected from hydrogen, C_{1-4} alkyl, $Het^{15}-C_{1-4}$ alkyl- or C_{1-4} alkyloxy C_{1-4} alkyl-;
- R^{16} and R^{17} each independently represents hydrogen or C_{1-4} alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl, aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;
- R^{18} and R^{19} each independently represents hydrogen or C_{1-4} alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl, aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;
- R^{20} and R^{22} each independently represents hydrogen or C_{1-4} alkyl optionally substituted with hydroxy or C_{1-4} alkyloxy;
- R^{21} represents hydrogen, C_{1-4} alkyl, $Het^{23}-C_{1-4}$ alkylcarbonyl- or
- R^{21} represents mono- or di(C_{1-4} alkyl)amino- C_{1-4} alkyl-carbonyl- optionally substituted with hydroxy, pyrimidinyl, dimethylamine or C_{1-4} alkyloxy;
- Het^1 represents a heterocycle selected from piperidinyl, morpholinyl, piperazinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het^1 is optionally substituted with amino, C_{1-4} alkyl, hydroxy- C_{1-4} alkyl-, phenyl, phenyl- C_{1-4} alkyl-, C_{1-4} alkyl-oxy- C_{1-4} alkyl- mono- or di(C_{1-4} alkyl)amino- or amino-carbonyl-;
- Het^2 represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl, thiomorpholinyl or dithianyl wherein said Het^2 is optionally substituted with one or where possible two or more substituents selected from hydroxy, halo, amino, C_{1-4} alkyl-, hydroxy- C_{1-4} alkyl-, C_{1-4} alkyl-oxy- C_{1-4} alkyl-, hydroxy- C_{1-4} alkyl-oxy- C_{1-4} alkyl-, mono- or di(C_{1-4} alkyl)amino-, mono- or di(C_{1-4} alkyl)amino- C_{1-4} alkyl-, amino C_{1-4} alkyl-, mono- or di(C_{1-4} alkyl)amino-sulfonyl-, aminosulfonyl-;
- Het^3 , Het^4 and Het^8 each independently represent a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het^3 , Het^4 or Het^8 is optionally substituted with one or where possible two or more substituents selected from hydroxy-, amino-, C_{1-4} alkyl-, C_{3-6} cycloalkyl- C_{1-4} alkyl-, aminosulfonyl-, mono- or di(C_{1-4} alkyl)aminosulfonyl or amino- C_{1-4} alkyl-;
- Het^5 represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from

-170-

C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het⁶ and Het⁷ each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het⁹ and Het¹⁰ each independently represent a heterocycle selected from furanyl, piperidinyl, morpholinyl, piperazinyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het⁹ or Het¹⁰ is optionally substituted C₁₋₄alkyl, C₃₋₆cycloalkyl-C₁₋₄alkyl- or amino-C₁₋₄alkyl-;



Het¹¹ represents a heterocycle selected from indolyl or

Het¹² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl, thiomorpholinyl or dithianyl wherein said Het¹² is optionally substituted with one or where possible two or more substituents selected from hydroxy, halo, amino, C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-oxy-C₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino- or mono- or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-;

Het¹³ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het¹⁴ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het¹⁵ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het¹⁶ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl, 1,3,2-dioxaborolane or piperidinyl wherein said heterocycle is optionally substituted with one or more substituents selected from C₁₋₄alkyl; and

Het¹⁷ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from

-171-

C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het¹⁸ and Het¹⁹ each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het²⁰, Het²¹ and Het²² each independently represent a heterocycle selected from pyrrolidinyl, 2-pyrrolidinonyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from hydroxy, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl- or polyhydroxy-C₁₋₄alkyl-;

Het²³ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Ar¹, Ar², Ar³, Ar⁴ and Ar⁵ each independently represent phenyl optionally substituted with cyano, C₁₋₄alkylsulfonyl-, C₁₋₄alkylsulfonylamino-, aminosulfonylamino-, hydroxy-C₁₋₄alkyl, aminosulfonyl-, hydroxy-, C₁₋₄alkyloxy- or C₁₋₄alkyl.

2. A compound according to claim 1 wherein;

Z represents O, NH or S;

Y represents -C₃₋₉alkyl-, -C₃₋₉alkenyl-, -C₁₋₅alkyl-oxy-C₁₋₅alkyl-,

-C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-, -C₁₋₅alkyl-NR¹⁴-CO-C₁₋₅alkyl-,

-C₁₋₅alkyl-CO-NR¹⁵-C₁₋₅alkyl-, -C₁₋₆alkyl-CO-NH-, -C₁₋₆alkyl-NH-CO-,

-CO-NH-C₁₋₆alkyl-, -NH-CO-C₁₋₆alkyl-, -CO-C₁₋₇alkyl-, -C₁₋₇alkyl-CO-,

-C₁₋₆alkyl-CO-C₁₋₆alkyl-, -C₁₋₂alkyl-NH-CO-CH₂R¹⁶-NH-;

X¹ represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹¹,

-NR¹¹-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or -C₁₋₂alkyl-;

X² represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹²,

NR¹²-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or -C₁₋₂alkyl-;

R¹ represents hydrogen, cyano, halo, hydroxy, formyl, C₁₋₆alkoxy-, C₁₋₆alkyl, C₁₋₆alkoxy- substituted with halo,

C₁₋₄alkyl substituted with one or where possible two or more substituents selected from hydroxy or halo;

R² represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-, Het¹⁶-carbonyl-, C₁₋₄alkyloxycarbonyl-, C₁₋₄alkylcarbonyl-, aminocarbonyl-, mono-or


-172-

- di(C₁₋₄alkyl)aminocarbonyl-, Het¹, formyl, C₁₋₄alkyl-, C₂₋₆alkynyl-, C₃₋₆cycloalkyl-, C₃₋₆cycloalkyloxy-, C₁₋₆alkoxy-, Ar⁵, Ar¹-oxy-, dihydroxyborane, C₁₋₆alkoxy- substituted with halo, C₁₋₄alkyl substituted with one or where possible two or more substituents selected from halo, hydroxy or NR⁵R⁶, C₁₋₄alkylcarbonyl- wherein said C₁₋₄alkyl is optionally substituted with one or where possible two or more substituents selected from hydroxy or C₁₋₄alkyl-oxy-;
- R³ represents hydrogen, C₁₋₄alkyl or C₁₋₄alkyl substituted with one or more substituents selected from halo, C₁₋₄alkyloxy-, amino-, mono- or di(C₁₋₄alkyl)amino-, C₁₋₄alkyl-sulfonyl- or phenyl;
- R⁴ represents hydrogen, hydroxy, Ar³-oxy, Ar⁴-C₁₋₄alkyloxy-, C₁₋₄alkyloxy-, C₂₋₄alkenyloxy- optionally substituted with Het¹² or R⁴ represents C₁₋₄alkyloxy substituted with one or where possible two or more substituents selected from C₁₋₄alkyloxy-, hydroxy, halo, Het²-, -NR⁷R⁸-, -carbonyl- NR⁹R¹⁰ or Het³-carbonyl-;
- R⁵ and R⁶ are each independently selected from hydrogen or C₁₋₄alkyl;
- R⁷ and R⁸ are each independently selected from hydrogen, C₁₋₄alkyl, Het⁸, aminosulfonyl-, mono- or di (C₁₋₄alkyl)-aminosulfonyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl-, hydroxycarbonyl-C₁₋₄alkyl-, C₃₋₆cycloalkyl, Het⁹-carbonyl-C₁₋₄alkyl-, Het¹⁰-carbonyl-, polyhydroxy-C₁₋₄alkyl-, Het¹¹-C₁₋₄alkyl- or Ar²-C₁₋₄alkyl-;
- R⁹ and R¹⁰ are each independently selected from hydrogen, C₁₋₄alkyl, C₃₋₆cycloalkyl, Het⁴, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl- or polyhydroxy-C₁₋₄alkyl-;
- R¹¹ represents hydrogen, C₁₋₄alkyl, Het⁵, Het⁶-C₁₋₄alkyl-, C₂₋₄alkenylcarbonyl- optionally substituted with Het⁷-C₁₋₄alkylaminocarbonyl-, C₂₋₄alkenylsulfonyl-, C₁₋₄alkyloxyC₁₋₄alkyl- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;
- R¹² represents hydrogen, C₁₋₄alkyl, C₁₋₄alkyl-oxy-carbonyl-, Het¹⁷, Het¹⁸-C₁₋₄alkyl-, C₂₋₄alkenylcarbonyl- optionally substituted with Het¹⁹-C₁₋₄alkylaminocarbonyl-, C₂₋₄alkenylsulfonyl-, C₁₋₄alkyloxyC₁₋₄alkyl- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;
- R¹³ represents hydrogen, C₁₋₄alkyl, Het¹³, Het¹⁴-C₁₋₄alkyl- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;

-173-

- R¹⁴ and R¹⁵ are each independently selected from hydrogen, C₁₋₄alkyl, Het¹⁵-C₁₋₄alkyl- or C₁₋₄alkyloxyC₁₋₄alkyl-;
- R¹⁶ represents hydrogen or C₁₋₄alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl, aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;
- Het¹ represents a heterocycle selected from piperidinyl, morpholinyl, piperazinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het¹ is optionally substituted with amino, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl-, phenyl, phenyl-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl- mono- or di(C₁₋₄alkyl)amino- or amino-carbonyl-;
- Het² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl, thiomorpholinyl or dithianyl wherein said Het² is optionally substituted with one or where possible two or more substituents selected from hydroxy, halo, amino, C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-oxy-C₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino-, mono- or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-, aminoC₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino-sulfonyl-, aminosulfonyl-;
- Het³, Het⁴ and Het⁸ each independently represent a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het³, Het⁴ or Het⁸ is optionally substituted with one or where possible two or more substituents selected from hydroxy-, amino-, C₁₋₄alkyl-, C₃₋₆cycloalkyl-C₁₋₄alkyl-, aminosulfonyl-, mono- or di(C₁₋₄alkyl)aminosulfonyl or amino-C₁₋₄alkyl-;
- Het⁵ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;
- Het⁶ and Het⁷ each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;
- Het⁹ and Het¹⁰ each independently represent a heterocycle selected from furanyl, piperidinyl, morpholinyl, piperazinyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het⁹ or Het¹⁰ is optionally substituted C₁₋₄alkyl, C₃₋₆cycloalkyl-C₁₋₄alkyl- or amino-C₁₋₄alkyl-;

-174-



Het¹¹ represents a heterocycle selected from indolyl or

Het¹² represents a heterocycle selected from morpholiny, piperaziny, piperidinyl, pyrrolidinyl, thiomorpholiny or dithianyl wherein said Het¹² is optionally substituted with one or where possible two or more substituents selected from hydroxy, halo, amino, C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-oxy-C₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino- or mono- or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-;

Het¹³ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het¹⁴ represent a heterocycle selected from morpholiny, pyrrolidinyl, piperaziny or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het¹⁵ represent a heterocycle selected from morpholiny, pyrrolidinyl, piperaziny or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het¹⁶ represent a heterocycle selected from morpholiny, pyrrolidinyl, piperaziny, 1,3,2-dioxaborolane or piperidinyl wherein said heterocycle is optionally substituted with one or more substituents selected from C₁₋₄alkyl; and

Het¹⁷ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het¹⁸ and Het¹⁹ each independently represent a heterocycle selected from morpholiny, pyrrolidinyl, piperaziny or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Ar¹, Ar², Ar³, Ar⁴ and Ar⁵ each independently represent phenyl optionally substituted with cyano, C₁₋₄alkylsulfonyl-, C₁₋₄alkylsulfonylamino-, aminosulfonylamino-, hydroxy-C₁₋₄alkyl, aminosulfonyl-, hydroxy-, C₁₋₄alkyloxy- or C₁₋₄alkyl.

3. A compound according to claims 1 or 2 wherein;

Z represents NH;

-175-

- Y represents -C₃₋₉alkyl-, -C₂₋₉alkenyl-, -C₁₋₅alkyl-oxy-C₁₋₅alkyl-,
 -C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-, -C₁₋₅alkyl-NR¹⁴-CO-C₁₋₅alkyl-, -C₁₋₆alkyl-NH-CO-,
 -NH-CO-C₁₋₆alkyl-, -CO-C₁₋₇alkyl-, -C₁₋₇alkyl-CO-, C₁₋₆alkyl-CO-C₁₋₆alkyl,
 -C₁₋₂alkyl-NH-CO-CR¹⁶R¹⁷-NH-, -C₁₋₂alkyl-CO-NH-CR¹⁸R¹⁹-CO-,
 5 -C₁₋₂alkyl-CO-NR²⁰-C₁₋₃alkyl-CO-, -C₁₋₂alkyl-NR²¹-CH₂-CO-NH-C₁₋₃alkyl-,
 -NR²²-CO-C₁₋₃alkyl-NH-, -C₁₋₃alkyl-NH-CO-Het²⁰-, C₁₋₂alkyl-CO-Het²¹-CO-, or
 -Het²²-CH₂-CO-NH-C₁₋₃alkyl-;
- X¹ represents O, -O-C₁₋₂alkyl-, -O-N=CH-, NR¹¹ or -NR¹¹-C₁₋₂alkyl-;
- X² represents a direct bond, -C₁₋₂alkyl-, O, -O-C₁₋₂alkyl-, -O-N=CH-, NR¹² or
 10 NR¹²-C₁₋₂alkyl-;
- R¹ represents hydrogen, cyano, halo or hydroxy;
- R² represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-,
 C₁₋₄alkyloxycarbonyl-, Het¹⁶-carbonyl-, C₁₋₄alkyl-, C₂₋₆alkynyl-, Ar⁵ or Het¹;
- R³ represents hydrogen;
- 15 R⁴ represents hydrogen, hydroxy, C₁₋₄alkyloxy- or R⁴ represents C₁₋₄alkyloxy
 substituted with one or where possible two or more substituents selected from
 C₁₋₄alkyloxy- or Het²;
- R¹² represents hydrogen, C₁₋₄alkyl- or C₁₋₄alkyl-oxy-carbonyl-;
- R¹³ represents hydrogen or Het¹⁴-C₁₋₄alkyl;
- 20 R¹⁴ and R¹⁵ represent hydrogen;
- R¹⁶ represents hydrogen or C₁₋₄alkyl substituted with hydroxy;
- R¹⁷ represents hydrogen or C₁₋₄alkyl, in particular hydrogen or methyl;
- R¹⁸ represents hydrogen or C₁₋₄alkyl optionally substituted with hydroxy or phenyl;
- R¹⁹ represents hydrogen or C₁₋₄alkyl;
- 25 R²⁰ represents hydrogen or C₁₋₄alkyl;
- R²¹ represents hydrogen, C₁₋₄alkyl, Het²³-C₁₋₄alkylcarbonyl- or
 R²¹ represents mono- or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-carbonyl- optionally
 substituted with hydroxy, pyrimidinyl, dimethylamine or C₁₋₄alkyloxy;
- R²² represents hydrogen or C₁₋₄alkyl optionally substituted with hydroxy or C₁₋₄alkyloxy;
- 30 Het¹ represents thiazolyl optionally substituted with amino, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl-,
 phenyl, phenyl-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl- mono- or di(C₁₋₄alkyl)amino- or amino-carbonyl-;
- Het² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or
 35 pyrrolidinyl wherein said Het² is optionally substituted with one or where possible
 two or more substituents selected from hydroxy, amino or C₁₋₄alkyl-;

-176-

Het³ represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or pyrrolidinyl wherein said Het³ is optionally substituted with one or where possible two or more substituents selected from hydroxy, amino or C₁₋₄alkyl-;

5 Het¹² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or pyrrolidinyl wherein said Het¹² is optionally substituted with one or where possible two or more substituents selected from hydroxy, amino or C₁₋₄alkyl-;

Het¹⁶ represents a heterocycle selected from piperidinyl or pyrrolidinyl;

Het²⁰ represents pyrrolidinyl, 2-pyrrolidinonyl, piperidinyl or hydroxy-pyrrolidinyl, preferably pyrrolidinyl or hydroxy-pyrrolidinyl;

10 Het²¹ represents pyrrolidinyl or hydroxy-pyrrolidinyl;

Het²² represents pyrrolidinyl, piperazinyl or piperidinyl.

4. A compound according to claims 1 or 2 wherein;

Z represents NH;

15 Y represents -C₃₋₉alkyl-, -C₂₋₉alkenyl-, -C₁₋₅alkyl-oxy-C₁₋₅alkyl-,

-C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-, -C₁₋₆alkyl-NH-CO-, -CO-C₁₋₇alkyl-, -C₁₋₇alkyl-CO- or C₁₋₆alkyl-CO-C₁₋₆alkyl;

X¹ represents O, -O-C₁₋₂alkyl-, -O-N=CH-, NR¹¹ or -NR¹¹-C₁₋₂alkyl-;

X² represents a direct bond, O, -O-C₁₋₂alkyl-, -O-N=CH-, NR¹² or NR¹²-C₁₋₂alkyl-;

20 R¹ represents hydrogen, cyano, halo or hydroxy, preferably halo;

R² represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-,

C₁₋₄alkyloxycarbonyl-, Het¹⁶-carbonyl-, C₂₋₆alkynyl-, Ar⁵ or Het¹;

R³ represents hydrogen;

R⁴ represents hydroxy, C₁₋₄alkyloxy- or R⁴ represents C₁₋₄alkyloxy substituted with one

25 or where possible two or more substituents selected from C₁₋₄alkyloxy- or Het²-;

R¹² represents hydrogen, C₁₋₄alkyl- or C₁₋₄alkyl-oxy-carbonyl-;

R¹³ represents Het¹⁴-C₁₋₄alkyl-;

Het¹ represents thiazolyl optionally substituted with amino, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl-, phenyl, phenyl-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl- mono- or di(C₁₋

30 ₄alkyl)amino- or amino-carbonyl-;

Het² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or pyrrolidinyl wherein said Het² is optionally substituted with one or where possible two or more substituents selected from hydroxy, amino or C₁₋₄alkyl-;

35 Het³ represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or pyrrolidinyl wherein said Het³ is optionally substituted with one or where possible two or more substituents selected from hydroxy, amino or C₁₋₄alkyl-;

-177-

Het¹² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or pyrrolidinyl wherein said Het¹² is optionally substituted with one or where possible two or more substituents selected from hydroxy, amino or C₁₋₄alkyl-;

Het¹⁶ represents a heterocycle selected from piperidinyl or pyrrolidinyl.

5

5. A compound according to claim 1 wherein;

Z represents NH;

Y represents -C₃₋₉alkyl-, -CO-C₁₋₇alkyl- or -C₁₋₇alkyl-CO-;

X¹ represents -NR¹¹-, -O- or -O-CH₂-;

10 X² represents a direct bond, -O- or -O-CH₂-;

R¹ represents halo;

R² represents hydrogen, cyano, halo, hydroxy or C₂₋₆alkynyl-;

R³ represents hydrogen;

15 R⁴ represents C₁₋₄alkyloxy substituted with one or where possible two or more substituents selected from C₁₋₄alkyloxy- or Het²-;

R¹² represents C₁₋₄alkyl or R¹² represents C₁₋₄alkyl-oxy-carbonyl;

10 Het² represents a heterocycle selected from morpholinyl or piperidinyl optionally substituted with C₁₋₄alkyl-;

20 Het³ represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or pyrrolidinyl wherein said Het³ is optionally substituted with one or where possible two or more substituents selected from hydroxy, amino or C₁₋₄alkyl-;

Het¹² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl or pyrrolidinyl wherein said Het¹² is optionally substituted with one or where possible two or more substituents selected from hydroxy, amino or C₁₋₄alkyl-.

25

6. A compound according to claim 1 wherein;

Z represents NH;

Y represents -C₃₋₉alkyl-, -C₂₋₉alkenyl-, -C₃₋₇alkyl-CO-NH optionally substituted with amino, mono- or di(C₁₋₄alkyl)amino or C₁₋₄alkyloxycarbonylamino-,
 30 -C₃₋₇alkenyl-CO-NH- optionally substituted with amino, mono- or di(C₁₋₄alkyl)amino- or C₁₋₄alkyloxycarbonylamino-,
 C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-, -C₁₋₅alkyl-NR¹⁴-CO-C₁₋₅alkyl-, -C₁₋₆alkyl-CO-NH-,
 -C₁₋₅alkyl-CO-NR¹⁵-C₁₋₅alkyl-, -C₁₋₃alkyl-NH-CO-Het²⁰-,
 -C₁₋₂alkyl-CO-Het²¹-CO-, -C₁₋₂alkyl-NH-CO-CR¹⁶R¹⁷-NH-,
 35 -C₁₋₂alkyl-CO-NH-CR¹⁸R¹⁹-CO-, -C₁₋₂alkyl-CO-NR²⁰-C₁₋₃alkyl-CO-, or
 -NR²²-CO-C₁₋₃alkyl-NH-;

X¹ represents a direct bond, O or -O-C₁₋₂alkyl-;

-178-

X² represents a direct bond, -CO-C₁₋₂alkyl-, NR¹², -NR¹²-C₁₋₂alkyl-, -O-N=CH- or -C₁₋₂alkyl-;

R¹ represents hydrogen or halo;

R² represents hydrogen or halo;

5 R³ represents hydrogen;

R⁴ represents hydrogen or C₁₋₄alkyloxy;

R¹² represents hydrogen or C₁₋₄alkyl;

R¹³ represents hydrogen or C₁₋₄alkyl;

R¹⁴ represents hydrogen;

10 R¹⁵ represents hydrogen;

R¹⁶ and R¹⁷ each independently represent hydrogen or C₁₋₄alkyl;

R¹⁸ and R¹⁹ each independently represent hydrogen or C₁₋₄alkyl optionally substituted with phenyl or hydroxy;

15 R²⁰ and R²¹ each independently represent hydrogen or C₁₋₄alkyl optionally substituted with C₁₋₄alkyloxy;

Het²⁰, Het²¹ and Het²² each independently represent a heterocycle selected from the group consisting pyrrolidinyl, 2-pyrrolidinonyl or piperidinyl optionally substituted with hydroxy.

20 7. A compound according to formula (I) wherein said compound is selected from the group consisting of;

4,6-ethanediylidenepyrimido[4,5-b][6,1,12]benzoxadiazacyclopentadecine, 17-bromo-8,9,10,11,12,13,14,19-octahydro-20-methoxy-13-methyl-,

25 4,6-ethanediylidenepyrimido[4,5-b][6,1,12]benzoxadiazacyclopentadecine, 17-bromo-8,9,10,11,12,13,14,19-octahydro-20-methoxy-,

4,6-ethanediylidenepyrimido[4,5-b][6,1,10,13]benzoxatriazacycloheptadecine-12,15-dione, 19-chloro-8,9,10,11,13,14,16,21-octahydro-22-methoxy-13-(2-methylpropyl)-,

30 4,6-ethanediylidenepyrimido[4,5-b][6,1,10,13]benzoxatriazacycloheptadecine-12,15-dione, 19-chloro-8,9,10,11,13,14,16,21-octahydro-22-methoxy-,

4,6-ethanediylidenepyrimido[4,5-b]pyrrolo[2,1-k][6,1,9,12]benzoxatriazacyclopentadecin-11(8H)-one, 19-chloro-18-fluoro-9,10,11a,12,13,14,16,21-octahydro-22-methoxy-,

35 4,6-ethanediylidene-8H-pyrimido[4,5-b][6,1,10,13]benzoxatriazacyclohexadecine, 18-chloro-9,10,11,12,13,14,15,20-octahydro-21-methoxy-14-methyl-,

4,6-ethanediylidenepyrimido[4,5-b][6,1,11,14]benzoxatriazacycloheptadecine, 19-chloro-8,9,10,11,12,13,14,15,16,21-decahydro-22-methoxy-15-methyl-,

-179-

4,6-ethanediylidenepyrimido[4,5-*b*][6,1,9,12]benzoxatriazacyclopentadecine, 17-chloro-8,9,10,11,12,13,14,19-octahydro-20-methoxy-13-methyl-,
12H-4,6-ethanediylidene-13,17-methanopyrimido[4,5-
b][6,1,10,16]benzoxatriazacyclonadecin-12-one, 21-chloro-
5 8,9,10,11,13,14,15,16,18,23-decahydro-25-methoxy-,
4,6-ethanediylidene-12H-pyrimido[4,5-*b*][6,1,10,13]benzoxatriazacyclohexadecin-12-one, 18-chloro-8,9,10,11,13,14,15,20-octahydro-21-methoxy-13,14-dimethyl-,
4,6-ethanediylidenepyrimido[4,5-*b*][6,1,11,14]benzoxatriazacycloheptadecin-13(8*H*)-one, 19-chloro-15-ethyl-9,10,11,12,14,15,16,21-octahydro-22-methoxy-, or
10 4,6-ethanediylidenepyrimido[4,5-*b*][6,1,11,14]benzoxatriazacycloheptadecin-13(8*H*)-one, 19-chloro-9,10,11,12,14,15,16,21-octahydro-22-methoxy-14,15-dimethyl-,

8. A compound according to any one of claims 1 to 6 wherein the X² substituent is at
15 position 2', the R¹ substituent is at position 4', the R² substituent is at position 5',
the R³ substituent is at position 3 and the R⁴ substituent at position 7 of the
structure of formula (I).

9. A kinase inhibitor of formula (I).

20

10. A compound as claimed in any one of claims 1 to 7 for use as a medicine.


11. Use of a compound as claimed in any one of claims 1 to 7 in the manufacture of a
medicament for treating cell proliferative disorders such as atherosclerosis,
25 restenosis and cancer.

12. A pharmaceutical composition comprising a pharmaceutically acceptable carrier
and, as active ingredient, an effective kinase inhibitory amount of a compound as
described in any one of the claims 1 to 7.


30

13. A process for preparing a compound as claimed in claims 1 to 7, comprising
a) coupling the 6-acetoxy-quinazolines of formula (II) with the suitable substituted
anilines of formula (III) to furnish the intermediates of formula (IV), and
deprotecting the intermediates of formula (IV) followed by ring closure under
35 suitable conditions.

-180-




b) Deprotecting the intermediates of formula (IV^b) followed by formation of the corresponding ether using the appropriate aminated alcohol under standard conditions provides the intermediates of formula (XXVIII). Next, deprotection followed by ring closure provides the target compounds of formula (I^b).



V = protective group such as for example, methylcarbonyl, t-butyl, methyl, ethyl, benzyl or trialkylsilyl groups, or in case of solid phase chemistry V represents the resin to which the remainder of the molecule is being attached.

R¹⁷ represents Ar³, Ar⁴-C₁₋₄alkyl, C₁₋₄alkyl, C₂₋₆alkenyl optionally substituted with Het¹² or R¹⁶ represents C₁₋₄alkyl substituted with one or where possible two or more substituents selected from C₁₋₄alkoxy, hydroxy, halo, Het², NR⁷R⁸, NR⁹R¹⁰-carbonyl or Het³-carbonyl, wherein Ar³, Ar⁴, Het¹², Het², R⁷, R⁸, R⁹, R¹⁰ and Het³ are defined as for the compounds of formula (I)

Y₁ and Y₂ each independently represent a C₁₋₅alkyl, CO-C₁₋₅alkyl or CO-CH₂R¹⁶-NH-



the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

V represents hydrogen or a protective group preferably selected from the group consisting of methylcarbonyl, t-butyl, methyl, ethyl, benzyl or trialkylsilyl;

Y represents -C₃₋₉alkyl-, -C₃₋₉alkenyl-, -C₃₋₇alkyl-CO-NH- optionally substituted with amino, mono- or di(C₁₋₄alkyl)amino or C₁₋₄alkyloxycarbonylamino-, -C₃₋₇alkenyl-CO-NH- optionally substituted with amino, mono- or di(C₁₋₄alkyl)amino or C₁₋₄alkyloxycarbonylamino-;

-C₁₋₅alkyl-oxy-C₁₋₅alkyl-, -C₁₋₅alkyl-NR¹³-C₁₋₅alkyl-, -C₁₋₅alkyl-NR¹⁴-CO-C₁₋₅alkyl-,
 -C₁₋₅alkyl-CO-NR¹⁵-C₁₋₅alkyl-, -C₁₋₆alkyl-CO-NH-, -C₁₋₆alkyl-NH-CO-,
 -C₁₋₃alkyl-NH-CS-Het²⁰-, -C₁₋₃alkyl-NH-CO-Het²⁰-, C₁₋₂alkyl-CO-Het²¹-CO-,
 -Het²²-CH₂-CO-NH-C₁₋₃alkyl-, -CO-NH-C₁₋₆alkyl-, -NH-CO-C₁₋₆alkyl-,
 -CO-C₁₋₇alkyl-, -C₁₋₇alkyl-CO-, -C₁₋₆alkyl-CO-C₁₋₆alkyl-,
 -CO-Het²⁰-, -C₁₋₂alkyl-NH-CO-CR¹⁶R¹⁷-NH-, -C₁₋₂alkyl-CO-NH-CR¹⁸R¹⁹-CO-,
 -C₁₋₂alkyl-CO-NR²⁰-C₁₋₃alkyl-CO-, -C₁₋₂alkyl-NR²¹-CH₂-CO-NH-C₁₋₃alkyl-, or
 -NR²²-CO-C₁₋₃alkyl-NH-;

X² represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹²-,
 -NR¹²-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or -C₁₋₂alkyl-;

R¹ represents hydrogen, cyano, halo, hydroxy, formyl, C₁₋₆alkoxy-, C₁₋₆alkyl-,
 halo-phenyl-carbonylamino-,
 C₁₋₆alkoxy- substituted with halo,
 C₁₋₄alkyl substituted with one or where possible two or more substituents selected from hydroxy or halo;

R² represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-, Het¹⁶-carbonyl-,
 C₁₋₄alkyloxycarbonyl-, C₁₋₄alkylcarbonyl-, aminocarbonyl-, mono-or
 di(C₁₋₄alkyl)aminocarbonyl-, Het¹, formyl, C₁₋₄alkyl-, C₂₋₆alkynyl-, C₃₋₆cycloalkyl-,
 C₃₋₆cycloalkyloxy-, C₁₋₆alkoxy-, Ar⁵, Ar¹-oxy-, dihydroxyborane ,
 C₁₋₆alkoxy- substituted with halo,

C₁₋₄alkyl substituted with one or where possible two or more substituents selected from halo, hydroxy or NR⁵R⁶,
 C₁₋₄alkylcarbonyl- wherein said C₁₋₄alkyl is optionally substituted with one or where possible two or more substituents selected from hydroxy or C₁₋₄alkyl-oxy-;

-182-

- R⁵ and R⁶ are each independently selected from hydrogen or C₁₋₄alkyl;
- R¹² represents hydrogen, C₁₋₄alkyl, C₁₋₄alkyl-oxy-carbonyl-, Het¹⁸-C₁₋₄alkyl-, phenyl-C₁₋₄alkyl-oxy-carbonyl-, Het¹⁷, C₂₋₄alkenylcarbonyl- optionally substituted with Het¹⁹-C₁₋₄alkylaminocarbonyl-, C₂₋₄alkenylsulfonyl-, C₁₋₄alkyloxyC₁₋₄alkyl- or R¹² represents phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;
- R¹³ represents hydrogen, C₁₋₄alkyl, Het¹³, Het¹⁴-C₁₋₄alkyl- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;
- R¹⁴ and R¹⁵ are each independently selected from hydrogen, C₁₋₄alkyl, Het¹⁵-C₁₋₄alkyl- or C₁₋₄alkyloxyC₁₋₄alkyl-;
- R¹⁶ and R¹⁷ each independently represents hydrogen or C₁₋₄alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl, aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;
- R¹⁸ and R¹⁹ each independently represents hydrogen or C₁₋₄alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl, aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;
- R²⁰ and R²² each independently represents hydrogen or C₁₋₄alkyl optionally substituted with hydroxy or C₁₋₄alkyloxy;
- R²¹ represents hydrogen, C₁₋₄alkyl, Het²³-C₁₋₄alkylcarbonyl- or R²¹ represents mono-or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-carbonyl- optionally substituted with hydroxy, pyrimidinyl, dimethylamine or C₁₋₄alkyloxy;
- Het¹ represents a heterocycle selected from piperidinyl, morpholinyl, piperazinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het¹ is optionally substituted with amino, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl-, phenyl, phenyl-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl- mono- or di(C₁₋₄alkyl)amino- or amino-carbonyl-;
- Het¹³ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;
- Het¹⁴ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;
- Het¹⁵ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more

-183-

substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het¹⁶ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl, 1,3,2-dioxaborolane or piperidinyl wherein said heterocycle is optionally substituted with one or more substituents selected from C₁₋₄alkyl; and

Het¹⁷ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;


Het¹⁸ and Het¹⁹ each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het²⁰, Het²¹ and Het²² each independently represent a heterocycle selected from pyrrolidinyl, 2-pyrrolidinonyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from hydroxy, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl- or polyhydroxy-C₁₋₄alkyl-;

Ar¹, Ar², Ar³, Ar⁴ and Ar⁵ each independently represent phenyl optionally substituted with cyano, C₁₋₄alkylsulfonyl-, C₁₋₄alkylsulfonylamino-, aminosulfonylamino-, hydroxy-C₁₋₄alkyl, aminosulfonyl-, hydroxy-, C₁₋₄alkyloxy- or C₁₋₄alkyl.

15. Use of an intermediate of formula (III) in the synthesis of a compound of formula (I).

25 16. An intermediate of formula (XXX)



the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

Y1 and Y2 each independently represent a C₁₋₅alkyl, C₁₋₆alkyl, CO-C₁₋₆alkyl, CO-C₁₋₅alkyl, Het²²-CH₂-CO, CO-CR¹⁶R¹⁷-NH-, Het²⁰, CR¹⁸R¹⁹-CO-, CH₂-CO-NH-C₁₋₃alkyl-, -C₁₋₂alkyl-NR²¹-CH₂-CO- or CO-C₁₋₃alkyl-NH-;

-184-

- X^1 represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹¹,
-NR¹¹-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or -C₁₋₂alkyl-;
- X^2 represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹²,
-NR¹²-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or -C₁₋₂alkyl-;
- 5 R^1 represents hydrogen, cyano, halo, hydroxy, formyl, C₁₋₆alkoxy-, C₁₋₆alkyl-,
halo-phenyl-carbonylamino-,
C₁₋₆alkoxy- substituted with halo,
C₁₋₄alkyl substituted with one or where possible two or more substituents selected
from hydroxy or halo;
- 10 R^2 represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-, Het¹⁶-carbonyl-,
C₁₋₄alkyloxycarbonyl-, C₁₋₄alkylcarbonyl-, aminocarbonyl-, mono- or
di(C₁₋₄alkyl)aminocarbonyl-, Het¹, formyl, C₁₋₄alkyl-, C₂₋₆alkynyl-, C₃₋₆cycloalkyl-,
C₃₋₆cycloalkyloxy-, C₁₋₆alkoxy-, Ar⁵, Ar¹-oxy-, dihydroxyborane ,
C₁₋₆alkoxy- substituted with halo,
- 15 C₁₋₄alkyl substituted with one or where possible two or more substituents selected
from halo, hydroxy or NR⁵R⁶,
C₁₋₄alkylcarbonyl- wherein said C₁₋₄alkyl is optionally substituted with one or
where possible two or more substituents selected from hydroxy or
C₁₋₄alkyl-oxy-;
- 20 R^3 represents hydrogen, C₁₋₄alkyl, or C₁₋₄alkyl substituted with one or more substituents
selected from halo, C₁₋₄alkyloxy-, amino-, mono- or di(C₁₋₄alkyl)amino-,
C₁₋₄alkyl-sulfonyl- or phenyl;
- R^5 and R^6 are each independently selected from hydrogen or C₁₋₄alkyl;
 R^7 and R^8 are each independently selected from hydrogen, C₁₋₄alkyl, Het⁸,
- 25 aminosulfonyl-, mono- or di (C₁₋₄alkyl)-aminosulfonyl, hydroxy-C₁₋₄alkyl-,
C₁₋₄alkyl-oxy-C₁₋₄alkyl-, hydroxycarbonyl-C₁₋₄alkyl-, C₃₋₆cycloalkyl,
Het⁹-carbonyl-C₁₋₄alkyl-, Het¹⁰-carbonyl-, polyhydroxy-C₁₋₄alkyl-, Het¹¹-C₁₋₄alkyl-
or Ar²-C₁₋₄alkyl-;
- R^9 and R^{10} are each independently selected from hydrogen, C₁₋₄alkyl, C₃₋₆cycloalkyl,
30 Het⁴, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl- or polyhydroxy-C₁₋₄alkyl-;
- R^{11} represents hydrogen, C₁₋₄alkyl, Het⁵, Het⁶-C₁₋₄alkyl-, C₂₋₄alkenylcarbonyl-
optionally substituted with Het⁷-C₁₋₄alkylaminocarbonyl-, C₂₋₄alkenylsulfonyl-,
C₁₋₄alkyloxyC₁₋₄alkyl- or phenyl optionally substituted with one or where possible
two or more substituents selected from hydrogen, hydroxy, amino or
- 35 C₁₋₄alkyloxy-;
- R^{12} represents hydrogen, C₁₋₄alkyl, C₁₋₄alkyl-oxy-carbonyl-, Het¹⁷, Het¹⁸-C₁₋₄alkyl-,
C₂₋₄alkenylcarbonyl- optionally substituted with Het¹⁹-C₁₋₄alkylaminocarbonyl-,

-185-

C₂₋₄alkenylsulfonyl-, C₁₋₄alkyloxyC₁₋₄alkyl- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C₁₋₄alkyloxy-;

R¹⁶ and R¹⁷ each independently represents hydrogen or C₁₋₄alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl, aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;

R¹⁸ and R¹⁹ each independently represents hydrogen or C₁₋₄alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl, aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;

R²¹ represents hydrogen, C₁₋₄alkyl, Het²³-C₁₋₄alkylcarbonyl- or R²¹ represents mono- or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-carbonyl- optionally substituted with hydroxy, pyrimidinyl, dimethylamine or C₁₋₄alkyloxy;

R²³ represents Ar³, Ar⁴-C₁₋₄alkyl, C₁₋₄alkyl, C₂₋₆alkenyl optionally substituted with

Het¹² or R¹⁷ represents C₁₋₄alkyl substituted with one or where possible two or more substituents selected from C₁₋₄alkyloxy, hydroxy, halo, Het², NR⁷R⁸, NR⁹R¹⁰-carbonyl or Het³-carbonyl;

Het¹ represents a heterocycle selected from piperidinyl, morpholinyl, piperazinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het¹ is optionally substituted with amino, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl-, phenyl, phenyl-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl- mono- or di(C₁₋₄alkyl)amino- or amino-carbonyl-;

Het² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl, thiomorpholinyl or dithianyl wherein said Het² is optionally substituted with one or where possible two or more substituents selected from hydroxy, halo, amino, C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-oxy-C₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino-, mono- or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-, aminoC₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino-sulfonyl-, aminosulfonyl-;

Het³, Het⁴ and Het⁸ each independently represent a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het³, Het⁴ or Het⁸ is optionally substituted with one or where possible two or more substituents selected from hydroxy-, amino-, C₁₋₄alkyl-, C₃₋₆cycloalkyl-C₁₋₄alkyl-, aminosulfonyl-, mono- or di(C₁₋₄alkyl)aminosulfonyl or amino-C₁₋₄alkyl-;


Het⁵ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from

-186-

C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het⁶ and Het⁷ each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het⁹ and Het¹⁰ each independently represent a heterocycle selected from furanyl, piperidinyl, morpholinyl, piperazinyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het⁹ or Het¹⁰ is optionally substituted C₁₋₄alkyl, C₃₋₆cycloalkyl-C₁₋₄alkyl- or amino-C₁₋₄alkyl-;



Het¹¹ represents a heterocycle selected from indolyl or

Het¹² represents a heterocycle selected from morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl, thiomorpholinyl or dithianyl wherein said Het¹² is optionally substituted with one or where possible two or more substituents selected from hydroxy, halo, amino, C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyl-oxy-C₁₋₄alkyl-, hydroxy-C₁₋₄alkyl-oxy-C₁₋₄alkyl-, mono- or di(C₁₋₄alkyl)amino- or mono- or di(C₁₋₄alkyl)amino-C₁₋₄alkyl-;

Het¹⁶ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl, 1,3,2-dioxaborolane or piperidinyl wherein said heterocycle is optionally substituted with one or more substituents selected from C₁₋₄alkyl; and

Het¹⁷ represent a heterocycle selected from pyrrolidinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het¹⁸ and Het¹⁹ each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-, C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;

Het²⁰, Het²¹ and Het²² each independently represent a heterocycle selected from pyrrolidinyl, 2-pyrrolidinonyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from hydroxy, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl- or polyhydroxy-C₁₋₄alkyl-;

Het²³ represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more

-187-


substituents selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, hydroxy-C₁₋₄alkyl-,
 C₁₋₄alkyloxyC₁₋₄alkyl or polyhydroxy-C₁₋₄alkyl-;
 Ar¹, Ar³, Ar⁴ and Ar⁵ each independently represent phenyl optionally substituted with
 cyano, C₁₋₄alkylsulfonyl-, C₁₋₄alkylsulfonylamino-, aminosulfonylamino-,
 5 hydroxy-C₁₋₄alkyl, aminosulfonyl-, hydroxy-, C₁₋₄alkyloxy- or C₁₋₄alkyl.

17. An intermediate of formula (XXX) for use as a medicine.

18. Use of an intermediate of formula (XXX) in the manufacture of a medicament for
 10 treating cell proliferative disorders such as atherosclerosis, restenosis and cancer.

19. Use of an intermediate of formula (XXX) in the synthesis of a compound of
 formula (I).

15 20. An intermediate of formula (XXXIII)



the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

m represents 1, 2, 3 or 4;

20 X² represents a direct bond, O, -O-C₁₋₂alkyl-, CO, -CO-C₁₋₂alkyl-, NR¹²,
 -NR¹²-C₁₋₂alkyl-, -CH₂-, -O-N=CH- or C₁₋₂alkyl;

Y₃ represents a C₁₋₅alkyl, CO-C₁₋₅alkyl or CO-CR¹⁶R¹⁷-NH- or C₁₋₅alkyl-CO-
 optionally substituted with amino, mono- or di(C₁₋₄alkyl)amino or
 C₁₋₄alkyloxycarbonylamino;

25 R¹ represents hydrogen, cyano, halo, hydroxy, formyl, C₁₋₆alkoxy-, C₁₋₆alkyl-,
 halo-phenyl-carbonylamino-,
 C₁₋₆alkoxy- substituted with halo,
 C₁₋₄alkyl substituted with one or where possible two or more substituents selected
 from hydroxy or halo;

-188-

- R^2 represents hydrogen, cyano, halo, hydroxy, hydroxycarbonyl-, Het^{16} -carbonyl-, C_{1-4} alkyloxycarbonyl-, C_{1-4} alkylcarbonyl-, aminocarbonyl-, mono- or di(C_{1-4} alkyl)aminocarbonyl-, Het^1 , formyl, C_{1-4} alkyl-, C_{2-6} alkynyl-, C_{3-6} cycloalkyl-, C_{3-6} cycloalkyloxy-, C_{1-6} alkoxy-, Ar^5 , Ar^1 -oxy-, dihydroxyborane, C_{1-6} alkoxy- substituted with halo, C_{1-4} alkyl substituted with one or where possible two or more substituents selected from halo, hydroxy or NR^5R^6 , C_{1-4} alkylcarbonyl- wherein said C_{1-4} alkyl is optionally substituted with one or where possible two or more substituents selected from hydroxy or C_{1-4} alkyl-oxy-;
- R^3 represents hydrogen, C_{1-4} alkyl, or C_{1-4} alkyl substituted with one or more substituents selected from halo, C_{1-4} alkyloxy-, amino-, mono- or di(C_{1-4} alkyl)amino-, C_{1-4} alkyl-sulfonyl- or phenyl;
- R^5 and R^6 are each independently selected from hydrogen or C_{1-4} alkyl;
- R^{12} represents hydrogen, C_{1-4} alkyl, C_{1-4} alkyl-oxy-carbonyl-, Het^{18} - C_{1-4} alkyl-, phenyl- C_{1-4} alkyl-oxy-carbonyl- Het^{17} , C_{2-4} alkenylcarbonyl- optionally substituted with Het^{19} - C_{1-4} alkylaminocarbonyl-, C_{2-4} alkenylsulfonyl-, C_{1-4} alkyloxy C_{1-4} alkyl- or phenyl optionally substituted with one or where possible two or more substituents selected from hydrogen, hydroxy, amino or C_{1-4} alkyloxy-;
- R^{16} and R^{17} each independently represents hydrogen or C_{1-4} alkyl optionally substituted with phenyl, indolyl, methylsulfide, hydroxy, thiol, hydroxyphenyl, aminocarbonyl, hydroxycarbonyl, amine, imidazolyl or guanidino;
- Het^1 represents a heterocycle selected from piperidinyl, morpholinyl, piperazinyl, furanyl, pyrazolyl, dioxolanyl, thiazolyl, oxazolyl, imidazolyl, isoxazolyl, oxadiazolyl, pyridinyl or pyrrolidinyl wherein said Het^1 is optionally substituted with amino, C_{1-4} alkyl, hydroxy- C_{1-4} alkyl-, phenyl, phenyl- C_{1-4} alkyl-, C_{1-4} alkyl-oxy- C_{1-4} alkyl- mono- or di(C_{1-4} alkyl)amino- or amino-carbonyl-;
- Het^{16} represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl, 1,3,2-dioxaborolane or piperidinyl wherein said heterocycle is optionally substituted with one or more substituents selected from C_{1-4} alkyl;
- Het^{18} and Het^{19} each independently represent a heterocycle selected from morpholinyl, pyrrolidinyl, piperazinyl or piperidinyl optionally substituted with one or where possible two or more substituents selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, hydroxy- C_{1-4} alkyl-, C_{1-4} alkyloxy C_{1-4} alkyl or polyhydroxy- C_{1-4} alkyl-; and
- Ar^1 and Ar^5 each independently represent phenyl optionally substituted with cyano, C_{1-4} alkylsulfonyl-, C_{1-4} alkylsulfonylamino-, aminosulfonylamino-, hydroxy- C_{1-4} alkyl, aminosulfonyl-, hydroxy-, C_{1-4} alkyloxy- or C_{1-4} alkyl.

-189-

21. An intermediate of formula (XXXIII) for use as a medicine.
22. Use of an intermediate of formula (XXXIII) in the manufacture of a medicament
5 for treating cell proliferative disorders such as atherosclerosis, restenosis and cancer.
23. Use of an intermediate of formula (XXXIII) in the synthesis of a compound of formula (I).

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/005621

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/529 C07D498/08 A61P35/00 C07D239/88 C07D239/93
C07D239/94

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96/33980 A (ZENECA LTD ; GIBSON KEITH HOPKINSON (GB)) 31 October 1996 (1996-10-31) cited in the application	1-13,15, 19,23
Y	page 1, line 1 - page 4, line 11; claims 1,16,19	16-18, 20-22
A	WO 96/09294 A (WELLCOME FOUND ; HUDSON ALAN THOMAS (GB); VILE SADIE (GB); BARRACLOUGH) 28 March 1996 (1996-03-28)	1-13,15, 19,23
Y	claims 1,19,22	16-18, 20-22
X	GB 1 542 514 A (JANSSEN PHARMACEUTICA NV) 21 March 1979 (1979-03-21) page 8, formula XI; examples III-V,XIX	14
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

12 October 2004

Date of mailing of the international search report

27/10/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Seymour, L

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/005621

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 1 465 451 A (BAYER AG) 23 February 1977 (1977-02-23) page 5, formula II; example 49	14
X	GB 807 899 A (ICI LTD) 21 January 1959 (1959-01-21) claim 1 page 2, line 73 - line 75 page 3, line 3 - line 4 page 3, line 19 - line 20	14
X	FURUTA K ET AL: "Molecular Design of Glutathione-derived Biochemical Probes targeting the GS-X Pump" TETRAHEDRON, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 55, no. 24, 11 June 1999 (1999-06-11), pages 7529-7540, XP004167533 ISSN: 0040-4020 compound 14	14
X	KAWATO H C ET AL.: "Novel peptidomimetics of the antifungal cyclic peptide Rhodopeptin: synthesis of mimetics and their antifungal activity" ORGANIC LETTERS., vol. 3, no. 22, 2001, pages 3451-3454, XP002300247 USACS, WASHINGTON, DC. compound 28	14
X	DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; 2001, XP002300248 Database accession no. 6269642 (CNR) BRN 1104891, 3260032, 8686727, 8687097 & RUSSIAN JOURNAL OF ORGANIC CHEMISTRY., vol. 36, no. 5, 2000, pages 674-678, USCONSULTANTS BUREAU.	14
X	DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; 2000, XP002300249 Database accession no. 6164062 (CNR) BRN 7476906, 7477251, 7477808, 8203983, 8205486, 8205933, 8207525 & JOURNAL OF THE CHEMICAL SOCIETY, PERKIN TRANSACTIONS 1., no. 15, 1998, pages 2331-2340, GBCHEMICAL SOCIETY. LETCHWORTH.	14
-/-		

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/005621

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE BEILSTEIN BEILSTEIN INSTITUTE FOR ORGANIC CHEMISTRY, FRANKFURT-MAIN, DE; 1994, XP002300250 Database accession no. 160121 (CNR) BRN 2807833, 2809124, 2813863, 2816816, 6308537, 6310017 & JOURNAL OF THE CHEMICAL SOCIETY., 1961, pages 1863-1879, GBCHEMICAL SOCIETY. LETCHWORTH.</p>	14

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/005621

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9633980	A	31-10-1996	AT 198329 T	15-01-2001
			AU 699163 B2	26-11-1998
			AU 5343396 A	18-11-1996
			BG 62730 B1	30-06-2000
			BG 102052 A	31-08-1998
			BR 9608082 A	26-01-1999
			CA 2215732 A1	31-10-1996
			CN 1182421 A , B	20-05-1998
			CZ 9703396 A3	18-02-1998
			DE 69611361 D1	01-02-2001
			DE 69611361 T2	26-04-2001
			DK 823900 T3	02-04-2001
			EE 9700252 A	15-04-1998
			EP 0823900 A1	18-02-1998
			ES 2153098 T3	16-02-2001
			WO 9633980 A1	31-10-1996
			GR 3035211 T3	30-04-2001
			HK 1005371 A1	16-11-2001
			HR 960204 A1	31-08-1997
			HU 9802839 A2	29-03-1999
			IL 118045 A	31-10-2001
			JP 3040486 B2	15-05-2000
			JP 11504033 T	06-04-1999
			NO 974940 A	24-10-1997
			NZ 305444 A	29-03-1999
			PL 323066 A1	02-03-1998
			PT 823900 T	30-04-2001
			RO 117849 B1	30-08-2002
			RU 2153495 C2	27-07-2000
			SI 823900 T1	30-06-2001
			SK 145497 A3	04-02-1998
			TW 436486 B	28-05-2001
			US 5770599 A	23-06-1998
			ZA 9603358 A	28-10-1996
WO 9609294	A	28-03-1996	AU 3482495 A	09-04-1996
			EP 0782570 A1	09-07-1997
			WO 9609294 A1	28-03-1996
			JP 10505600 T	02-06-1998
			TR 960233 A2	21-06-1996
			ZA 9507853 A	18-03-1997
GB 1542514	A	21-03-1979	KR 8100334 A	20-04-1981
			AT 364358 B	12-10-1981
			AT 534776 A	15-03-1981
			AU 510027 B2	05-06-1980
			AU 511027 B2	24-07-1980
			AU 1596476 A	19-01-1978
			BE 844087 A2	14-01-1977
			BG 26526 A3	12-04-1979
			CA 1085852 A1	16-09-1980
			CH 623820 A5	30-06-1981
			CS 202550 B2	30-01-1981
			DE 2632870 A1	10-02-1977
			DK 327576 A , B,	22-01-1977
			ES 449740 A1	16-02-1978
			ES 461630 A1	16-06-1978
			ES 461631 A1	01-12-1978

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/005621

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 1542514	A		FI 762086 A ,B,	22-01-1977
			FI 813935 A ,B,	08-12-1981
			FR 2361889 A1	17-03-1978
			HK 14180 A	28-03-1980
			IE 43496 B1	11-03-1981
			IT 1062614 B	20-10-1984
			JP 1364895 C	09-02-1987
			JP 52017475 A	09-02-1977
			JP 61031109 B	17-07-1986
			JP 1419185 C	14-01-1988
			JP 61093183 A	12-05-1986
			JP 62025669 B	04-06-1987
			KE 3215 A	09-07-1982
			LU 75354 A1	28-02-1977
			MY 1781 A	31-12-1981
			NL 7608023 A ,B,	25-01-1977
			NZ 181256 A	28-04-1978
			PH 13710 A	09-09-1980
			PT 65387 A ,B	01-08-1976
			RO 70566 A1	11-10-1982
			SE 426490 B	24-01-1983
			SE 7608265 A	22-01-1977
			SE 435621 B	08-10-1984
			SE 8008713 A	11-12-1980
			US 4066772 A	03-01-1978
			US 4175129 A	20-11-1979
			YU 179476 A1	21-01-1983
			ZA 7604322 A	29-03-1978
			HU 173220 B	28-03-1979
			IL 50073 A	30-11-1979
			NO 762397 A ,B,	24-01-1977
			PL 191309 A1	30-07-1979
			SU 701534 A3	30-11-1979
			SU 795466 A3	07-01-1981
GB 1465451	A	23-02-1977	DE 2423536 A1	27-11-1975
			AT 365375 A	15-11-1977
			BE 829051 A1	14-11-1975
			DD 120116 A5	05-06-1976
			DK 211575 A	16-11-1975
			FI 751400 A	16-11-1975
			FR 2271201 A1	12-12-1975
			JP 51001643 A	08-01-1976
			LU 72461 A1	17-03-1976
			NL 7505606 A	18-11-1975
			SE 7505526 A	17-11-1975
			US 4067726 A	10-01-1978
			ZA 7503124 A	28-04-1976
GB 807899	A	21-01-1959	BE 558251 A	14-04-1959
			FR 1176645 A	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☒ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.